


UMASS/AMHERST




312066 0033 0256 2



8-16  
2  
C. Funds  
12/62











# BULLETIN

OF THE

## NUTTALL ORNITHOLOGICAL CLUB:

A Quarterly Journal of Ornithology.

VOLUME II.

---

Editor,

J. A. ALLEN.

Associate Editors,

S. F. BAIRD AND ELLIOTT COUES.

CAMBRIDGE, MASS. :  
PUBLISHED BY THE CLUB.  
1877.

UNIVERSITY PRESS : WELCH, BIGELOW, & Co.,  
CAMBRIDGE.



## CONTENTS OF VOLUME II.

### NUMBER I.

	PAGE
THE BLACK-AND-YELLOW WARBLER. By <i>William Brewster</i> . . . . .	1
GEOGRAPHICAL VARIATION IN <i>TURDUS MIGRATORIUS</i> . By <i>Robert Ridgway</i> . . . . .	8
UNUSUAL ABUNDANCE OF THE SNOWY OWL IN NEW ENGLAND. By <i>Ruthven Deane</i> . . . . .	9
DISTRIBUTION OF NEW ENGLAND BIRDS. — A REPLY TO DR. T. M. BREWER. By <i>H. A. Purdie</i> . . . . .	11
OCCURRENCE OF THE BARNACLE GOOSE ( <i>Bernicla leucopsis</i> ) ON LONG ISLAND, N. Y. By <i>G. N. Lawrence</i> . . . . .	18
CAPTURE OF A SECOND SPECIMEN OF <i>HELMINTHOPHAGA LAWRENCEI</i> . By <i>Harold Herrick</i> . . . . .	19
NOTICE OF A FEW BIRDS OF RARE OR ACCIDENTAL OCCURRENCE IN NEW ENGLAND. By <i>H. A. Purdie</i> . . . . .	20

#### RECENT LITERATURE.

Notices of five recent Ornithological Papers, 23. — Vennor's Rapacious Birds of Canada, 24.

#### GENERAL NOTES.

A Humming-Bird new to the Fauna of the United States, 26. — Note on *Podiceps dominicus*, 26. — Eastward Range of the Ferruginous Buzzard (*Archibuteo ferrugineus*), 26. — Occurrence of Leconte's Bunting (*Coturniculus lecontei*) in Iowa, 26. — Audubon's Warbler in Massachusetts, 27. — Occurrence of the Sooty Tern in Massachusetts, 27. — The Black Gyr-Falcon (*Falco sacer* var. *labradoria*) in Massachusetts, 27. — Notes on Birds new to the Fauna of Maine, etc., 27. — Northern Range of the Sharp-tailed Finch (*Ammodramus caudacutus*), 28.

### NUMBER II.

CORRECTIONS OF NOMENCLATURE IN THE GENUS <i>SIRUS</i> . By <i>Dr. Elliott Coues</i> , U. S. A. . . . .	25
NOTES ON THE BREEDING OF THE BLACK TERN ( <i>Hydrochelidon lariformis</i> ). By <i>T. S. Roberts</i> . . . . .	34
TWO UNDESCRIBED NESTS OF CALIFORNIAN BIRDS. By <i>William Brewster</i> . . . . .	37
A CONTRIBUTION TO THE BIOGRAPHY OF WILSON'S PHALAROPE. By <i>E. W. Nelson</i> . . . . .	38
A DEFENCE OF HIS CATALOGUE OF NEW ENGLAND BIRDS. By <i>Dr. T. M. Brewer</i> . . . . .	44

#### RECENT LITERATURE.

Burroughs's "Wake-Robin," 48. — Minot's "Land and Game Birds of New England," 49.

#### GENERAL NOTES.

Western Range of *Conurus carolinensis*, 50. — Fecundity of the Carolina Wren, 50. — The Louisiana Heron in Indiana, 51. — Note on the Cinnamon Teal (*Querquedula cyanoptera*), 51. — *Egithus exilipes* in Europe, 51. — Note on *Cupidonia cupido* var. *pallidicinctus*, Ridgway, 52. — Capture of the Egyptian Goose on Long Island, N. Y., 52. — MacCown's Longspur in Illinois, 52.

## NUMBER III.

DESCRIPTION OF A NEW SPECIES OF HUMMING-BIRD FROM CALIFORNIA. By <i>H. W. Henshaw</i> . . . . .	53
THE BIRDS OF GUADALOUPE ISLAND DISCUSSED WITH REFERENCE TO THE PRESENT GENESIS OF SPECIES. By <i>Robert Ridgway</i> . . . . .	58
AN UNDESCRIBED HYBRID BETWEEN TWO NORTH AMERICAN GROUSE. By <i>William Brewster</i> . . . . .	66

## RECENT LITERATURE.

Nelson's "Birds of Northeastern Illinois," 68. — Salvin on the *Procellariidæ*, 69. — Catalogue of the Birds of the Islands of Malta and Gozo, 70. — Ridgway's "Studies of the American *Falconidæ*," 70. — Recent Ornithological Articles in American Journals, 73. — Californian Ornithology, 76. — McCauley's "Notes on Texan Ornithology," 76.

## GENERAL NOTES.

Nest and Eggs of Townsend's Flycatcher, 77. — Persistency at Nest-Building in a House-Wren, 78. — A New Bird to Massachusetts, 78. — A New Form of *Surnia* to New England, 78. — Capture of the Philadelphia Vireo in New Hampshire, 78. — Occurrence of *Passerculus princeps* in New York, 78. — The Pigeon-Hawk (*Falco columbarius*) at Sea, 79. — Capture of a Second Specimen of *Helminthophaga leucobranchialis*, 79. — The Mottled Owl as a Fisherman, 80. — Breeding of Leach's Petrel on the Coast of Maine, 80. — Nest and Eggs of the Alaskan Wren, 82. — *Junco oregonus* in Illinois, 82. — *Leptoptili albifrons*, a Pigeon new to the United States Fauna, 82. — *Melospiza leucoptera* in Colorado, 83. — The Ruff and the Purple Gallinule in Ohio, 83. — Notes on *Nyctale acadica*, 83. — Probable Breeding of the Saw-Whet Owl in Massachusetts, 84.

## NUMBER IV.

NOTES ON MOLOTHRUS ÆNEUS. By <i>J. C. Merrill</i> , Assist. Surg., U. S. A. . . . .	85
ON SEVENTY-FIVE DOUBTFUL WEST-COAST BIRDS. By <i>J. G. Cooper</i> , M. D. . . . .	88
REMARKS ON SELASPHORUS ALLENI, Henshaw. By <i>D. G. Elliot</i> . . . . .	97
THE YELLOW-THROATED WARBLER ( <i>Dendroica dominica</i> ). By <i>William Brewster</i> . . . . .	102

## RECENT LITERATURE.

D'Hamonville's Enumeration of the Birds of Europe, 106. — Merriam's "Review of the Birds of Connecticut," 107.

## GENERAL NOTES.

Note on *Doricha enicura*, 108. — Occurrence of the Black Vulture or Carrion Crow in Ohio, 109. — Occurrence of the Western Nonpariel and Berlandier's Wren at Fort Brown, Texas, 109. — A Cuckoo's Egg in a Cedar-Bird's Nest, 110. — Occurrence for the first Time in England of the Robin (*Turdus migratorius*), 110.

[Photographic reprint.]

BULLETIN  
OF THE  
NUTTALL ORNITHOLOGICAL CLUB.

---

Vol. II.

JANUARY, 1877.

No. I.

---

THE BLACK-AND-YELLOW WARBLER (*DENDRECA  
MACULOSA*).

BY WILLIAM BREWSTER.

FIRST impressions are apt to be most lasting, and in many cases are engraved upon the memory with a vividness that defies the effacing influence of time. Thus the Black-and-Yellow Warbler was one of my earliest bird acquaintances, and I shall not soon forget our introduction.

My family was spending a few days in a quiet little country town in New Hampshire, when, one hot summer afternoon, finding time hang heavily on my hands, I borrowed an old gun, and at the country store, where everything was sold from a patent coffee-mill to the latest specific for rheumatism, I purchased a supply of ammunition, and, thus equipped, took to the woods and searched a long time in vain for game. At length, entering a grove of thickly growing young spruces, I sat down to rest on a mossy log. I had been there but a short time when I became conscious of faint sounds in the trees above and around me, — chirpings, twitterings, and occasionally a modest little effort at song. Watching attentively, I soon spied a movement among the branches, and a tiny bird hopped out into the light, presenting a bright yellow breast and throat for just a moment before flying into the next tree. Here was a revelation! I already knew a few of the most familiar birds, — the Robin, the Bluebird, the Sparrow, the Oriole, and some others; but it had never occurred to me that dark forests like these might be tenanted by such delicate and beautiful forms.

Only the tropics surely could boast such gems. With enthusiasm now fairly aroused, and animated with the spirit of an explorer, I went at once to work to investigate, and in the course of an hour or two more my ammunition was nearly exhausted, and quite a line of poor, lifeless, mutilated little birds lay arranged along the old log. Resting my gun against a neighboring tree, I examined long and carefully the results of my work. Scarcely any two of my specimens were alike, and as I contemplated in amazement their varied forms and coloring, I felt like the discoverer of a new world, and doubted whether human eyes had ever beheld the like before. Finally, the deepening twilight brought an end to my reveries, and, collecting my prizes, I took my way homeward. Taxidermy being to me then a sealed book, I had recourse to pepper and salt as preservatives, but a few hot days settled the matter and proved the ruin of my collection. I can recall with sufficient distinctness for identification but a single bird of them all,— a fine adult male Black-and-Yellow Warbler, which at the time I considered the handsomest, and which I still think cannot be surpassed in beauty by any New England representative of the family. That afternoon was an unlucky one for the birds. It laid the foundation for a taste that has since caused the destruction of thousands of their tribe.

The Black-and-Yellow Warbler arrives in Massachusetts from the South about the 15th of May. During the next two or three weeks they are abundant everywhere in congenial localities. Willow thickets near streams, ponds, and other damp places, suit them best, but it is also not unusual to find many in the upland woods, especially where young pines or other evergreens grow thickly. Their food at this season is exclusively insects, the larger part consisting of the numerous species of *Diptera*. The males sing freely, especially on warm bright mornings. They associate indifferently with all the migrating warblers, but not unfrequently I have found large flocks composed entirely of members of their own species, and in this way have seen at least fifty individuals collected in one small tract of woodland. By the first of June all excepting a few stragglers have left. If we follow them northward, we find a few pairs passing the summer on the mountains of Southern Maine and New Hampshire. In July, 1875, I found them breeding, in company with the Blackburnian Warbler (*Dendroica blackburniæ*), the Snowbird (*Junco hyemalis*), the Golden-crested Kinglet (*Regulus satrapa*), and

several other birds of the Canadian Fauna, on Mount Monadnock, New Hampshire, within fifteen miles of the Massachusetts State line. Throughout the White Mountains of New Hampshire they are everywhere common during the summer, but it is not until we reach the latitude of the Umbagog Lakes, in Western Maine, that we find them evenly distributed over high and low country alike. In this region summer succeeds winter so quickly that there is almost no spring. Thus when I reached Upton on the 25th of May, 1876, I found that the lakes had broken up but four days previously; not a leaf had unfolded, even in the most sheltered places, and snow lay in large masses everywhere in the hollows and on northern exposures. Yet many species of Warblers had already arrived, and among them the subject of the present sketch was well represented.

They kept closely about the buildings, and although the day was warm, maintained an almost perfect silence. Dozens at a time were hopping about the manure heap behind the stables and around the sink-spout, while all showed a certain apprehensiveness of manner, as if they feared the issue of their temerity in penetrating into so bleak and dreary a region. Taking a short walk into the woods, I found them untenanted, save by a few Titmice, Woodpeckers, and some of the earlier Sparrows. But in the course of the next week wonderful changes took place. The birches first, afterwards the maples, beech-trees, and poplars, put on a feathery drapery of the most delicate green. The shad-bush, (*Amelanchier canadensis*) and the "moose-wood" (*Cornus circinata*) became white with clustering blossoms, and looked at a distance like fleecy summer-clouds entangled among the trees. Underfoot, beautiful trilliums of both the purple (*Trillium erectum*) and white (*T. grandiflorum*) species, were conspicuous among a host of other wild-flowers. Bees hummed among the blossoms, and butterflies flitted airily through the forest glades. Everything was fresh, lovely, and suggestive of the calm, peaceful security of summer. Thus in one week were consummated changes that, farther south, are often extended through nearly thrice the time. All this while the birds had kept ample pace with the advance of the season. Hundreds were daily arriving, passing on, or settling into their accustomed summer-haunts, and the woods fairly rang with the first burst of their melody. During the next week all the Warblers, and most of the smaller birds generally, were occupied in pairing and constructing their nests. Then came

the harvest-time for the oölogist, and rarities were in order. But how brief it all was! A dozen or so days only, and the young were hatched out; the woods swarmed with mosquitoes, black flies, and other bloodthirsty insects, and "the season" was at an end. Nothing remained but to pack up the accumulated treasures, and get them safely home for future comparison and investigation.

Before taking out our cabinet specimens, however, and diving into the dry details of description, let us return to the woods, and contemplate for a few moments the undisturbed nest. We shall be most likely to find one along this old wood-road, for the removal of the taller trees has let in the sunlight a little, and birds love such places.

You will rarely find the interior of a forest so well peopled as the edges and little openings, and the birds are not singular in this respect. Men always choose the shores of rivers, ponds, or the sea, for their first settlements in a new country, and I fancy it is not entirely from considerations of utility, but partly because they crave an adjacent breathing-space, where the sun and wind may have fair sweep. There are some exceptions to the rule among the birds, of course, there being some morbidly disposed individuals that can find no place too dark or too secluded.

As we follow the old wood-path, you shall take one side while I make good the other. These little clumps of fir and spruce shrubs are the likely places, and, judging from the numbers of Black-and-Yellow Warblers that I hear singing, our chances are good, but you must remember that not above one male in three or four of this species is blessed with a mate, so do not let your hopes rise too high. They are a gay lot of bachelors, though, are they not? chasing one another through the branches, more in sport than anger apparently, and uttering their queer, emphatic little songs on all sides. *She knew she was right; yes, she knew she was right*, they seem to say; but what all this means I never could imagine. Some idle gossip of theirs probably, which it will not profit us to inquire into. Ha! I have it, even so soon. I thought you fellow singing so gayly in the fallen tree-top had more the air of a Benedict than any we have previously seen, and here, almost under my hand, sits his modest little wife on her nest. Be careful how you shake that branch, for I would have you take a good long look ere we disturb her. See how her dark little eye glistens, and note the rapid pulsating motion of her back. Underneath those puffed-up feathers a poor little heart

is beating wildly with fear and apprehension, but still she sits bravely on her trust. She would say, if she could, with the Roman mother, "These are my jewels," and would entreat us to spare them. Now I will advance my hand cautiously. See! I almost touch her tail with my finger-tips; but the next instant she is gone. How quietly at the last moment she slid over the edge of the nest, barely eluding my grasp! A faint cry or two, and there comes the male; but he, gaudy little braggart! is far better at singing brave deeds than performing them, and will not trust himself very near, though he keeps up a constant chirping. His mate, however, is bold enough for both, and in her anxiety almost comes within reach of our hands. Now look into the nest! Beauties, are they not! Four of them; rosy white, spotted prettily with amber, lilac, and a few scattered dashes of black. Observe how cunningly the whole affair is concealed, — built close to the stem of the little fir, resting on the flat horizontally disposed rows of "needles," and arched over by the flake-like layer of twigs above. One long rootlet alone hangs down in full view, and had it not caught my eye I might have passed without discovering the nest. It seems, indeed, a pity to disturb it, but we shall regret it next winter if we leave it behind. Naturalists are probably not hard-hearted by inclination, but of necessity. I dare say the female will commence another nest before we pass here on our way back, and the male will be singing as joyously as ever in an hour or two. Birds' grief, like their average lives, is short, though apparently intense for the time. It is only the end, however, that can ever justify the destruction of a nest, and unthinking persons might, in many cases, be benefited by contemplating a little more closely the suffering which they inflict.

As the published descriptions of the nesting of this species are meagre and more or less conflicting, I shall go somewhat fully into the matter.

*Location of Nest.* — The nest is usually placed in a small fir or spruce, and rarely at a greater elevation than five or six feet. The average height would probably not exceed four feet, and I have found some barely twelve inches above the ground. It is usually laid somewhat loosely among the horizontal twigs, from which it can in most cases be lifted intact. Favorite localities are the edges of wood-paths, or roads bordered by woods, and clearings grown up to small evergreens. Exceptional situations are the interior of

the woods, where, in some cases, the nest is placed in the top of a young hemlock ten or fifteen feet up. In one instance I found a nest on a horizontal spruce limb in the very heart of the forest, and at least *thirty-five feet* above the ground. This nest contained four eggs, and the female bird, which was sitting, established its identity beyond question.

*Composition of Nest.* — The framework is wrought somewhat loosely of fine twigs, those of the hemlock being apparently preferred. Next comes a layer of coarse grass or dry weed-stalks; while the interior is lined invariably with fine black roots, which closely resemble horse-hairs. In an examination of more than thirty examples I have found not one in which these black roots were not used. One specimen has, indeed, a few *real* horse-hairs in the lining, but the roots predominate. This uniform coal-black lining shows in strong contrast with the lighter aspect of the outer surface of the nest. The whole structure is loosely put together, and bears a no distant resemblance to the nest of the Chipping Sparrow (*Spizella socialis*).

Among nests of the *Sylvicolidae*, it finds, perhaps, its nearest approach in that of the Chestnut-sided Warbler (*Dendroica pennsylvanica*).

Audubon, describing a nest from Labrador, affirms that it was lined with "a great quantity of feathers." As regards this statement, I can only say that it is entirely at variance with my own observations, and the employment of feathers in a nest of this character seems to me almost *mal a propos* as it would in that of a Heron or Cuckoo.

*Description of Eggs, etc.* — The time of laying with this species varies, in relation to the season, from June 8 to June 15. Four eggs commonly constitute a set, though in some cases but three are laid; and I know of an instance where five were found in one nest. They measure about .62 of an inch in length by .50 in breadth. The usual shape is a rounded oval, and the ground-color almost invariably creamy white after the removal of the contents. The markings are most commonly blotches of rich, warm umber, with smaller dottings of pale lilac or brown, disposed about the larger end. Some specimens are, however, thickly sprinkled over their entire surface with fine brownish spots. One set of four eggs differs from any of the others in having a decided tinge of bluish in the ground-color; while upon the large patches of umber which en-



circle the greater ends are drawn numerous wavy lines of black, precisely like the characteristic pen-markings of some of the Oriole's eggs. With an extensive series of specimens before me, I am led to the inevitable conviction that eggs of *D. maculosa* are in many cases indistinguishable from those of *D. virens*, *D. pennsylvanica*, and *D. discolor*; and an examination of an equal number of authentic eggs of the other *Dendroica* would, I am satisfied, result in adding many more to this list. In the eggs of each of the above-named species there is an almost endless variation, and many sets are consequently quite unique, but the type — if, indeed, any can be established — finds equally near approaches among them all. Nests may, however, in most cases be relied upon, especially when procured from proximate localities.

In the case of the young, both before and after they leave the nest, this bird displays no exceptional traits. Both old and young, when the latter have become able to take care of themselves, join the immense congregations of mingled Warblers, Wrens, Titmice, Sparrows, and Woodpeckers, which collect in the northern forests in early August, to be dispersed — most of them southward — by the first frosts of September.

In Eastern Massachusetts this species occurs as a fall migrant from September 21 to October 30, but it is never seen at this season in anything like the numbers which pass through the same section in spring, and the bulk of the migration must follow a more westerly route. Its haunts while with us in the autumn are somewhat different from those which it affects during its northward journey. We now find it most commonly on hillsides, among scrub-oaks and scattered birches, and in company with such birds as the Yellow-Rump (*Dendroica coronata*) and the Black-Poll (*D. striata*). A dull, listless troop they are, comparatively sombre of plumage, totally devoid of song, and apparently intent only upon the gratification of their appetites. It seems, at first thought, strange that the birds, at a season when all the rest of Nature puts on its most gorgeous coloring, should array themselves in their dullest; but it must be borne in mind that many of them played their part before these brilliant leaves had burst their buds, and now, like ushers and orchestra, whose duty has been performed, they stand aside among the audience, and watch the shifting glories of the final transformation-scene. So let us leave them until, attired in fullest costume, they come again to herald, with overtures of joyous song, the rising of the curtain on a new year.

ON GEOGRAPHICAL VARIATION IN *TURDUS*  
*MIGRATORIUS*.

BY ROBERT RIDGWAY.

CERTAIN differences between Eastern specimens of the common Robin and those from the Rocky Mountains were first pointed out by Professor Baird, in his "Review of American Birds" (1864, pp. 28, 29), in the following words: "In highly plumaged specimens from the East the feathers of the interseapular region are frequently, even generally, tinged with blackish in their centres, passing gradually into ash on the edges, and the black of the head ceases to be abruptly defined. There is also usually a well-defined whitish tip, half an inch long, to the outer tail-feathers. In Rocky Mountain skins the tail is either black, except a very narrow whitish edge, or the white tips of Eastern specimens are replaced by a dull gray. The black of the head, too, is better defined, the interseapular feathers more uniformly ash, and the upper parts without the faint brownish wash so frequently seen in Eastern specimens. There are, however, some exceptions to these features in specimens from each locality. The colors generally of Western birds appear to be paler." Again, in the "History of North American Birds" (Vol. I, p. 25), the same and additional differences are alluded to, as follows: "There are some variations, both of color and proportions, between Eastern and Western specimens of the Robin. In the latter there is a tendency to a longer tail, though the difference is not marked, and, as a rule, they slightly exceed Eastern specimens in size. The broad white tip to the lateral tail-feather — so conspicuous a mark of Eastern birds — is scarcely to be found at all in any Western ones; and in the latter the black of the head is very sharply defined against the lighter, clearer ash of the back, there hardly ever being a tendency in it to continue backward in the form of central spots to the feathers, as is almost constantly seen in Eastern examples; of Western specimens, the rufous, too, is appreciably lighter than in Eastern."

Very extensive material received at the National Museum since the above was written tends to confirm the constancy of most of these differences between Eastern and Western Robins, while other

points of diversity, previously overlooked, have been detected, the most important being the much blacker tail of Eastern birds, and their decidedly shorter wing.

Upon the whole, the two forms seem to constitute two very strongly marked geographical races, which may be distinguished as follows:—

*T. migratorius*.—Wing, 4.85–5.33; tail, 4.10–4.60; bill, from nostril, .48–.51; tarsus, 1.20–1.35; middle toe, .83–.92.\* Inner web of outer tail-feather with a distinct white terminal spot. Tail-feathers of adult male dusky black, with slight edging of plumbeous. *Habitat*. Eastern region, including the whole of Alaska, Eastern Mexico, and the eastern border of the Missouri Plains.

*T. propinquus*, RIDGW. (MSS.).—Wing, 5.35–5.60; tail, 4.60–4.70; bill, from nostril, .50–.55; tarsus, 1.30–1.35; middle toe, .90. Inner web of lateral tail-feather with merely a narrower terminal edging of white, or with no white whatever. Tail-feathers of adult male dusky slate, without distinctly paler edges. *Habitat*. Western region, including eastern base of Rocky Mountains.

We find the character of blackish centres to the interscapulars in Eastern specimens to be too inconstant a feature to serve as a character. No specimens of the Western series are so marked, but many Eastern ones, otherwise typical, have no trace of these markings. It is a well-known fact that the eggs of the Western Robins average considerably larger in size than those of Eastern birds.

---

### UNUSUAL ABUNDANCE OF THE SNOWY OWL (*NYCTEA SCANDIACA*) IN NEW ENGLAND.

BY RUTHVEN DEANE.

IN a recent number of this Bulletin (Vol. I, p. 95), Mr. N. C. Brown gave some interesting notes respecting the variable abundance of birds at the same locality in different seasons. I do not think a more forcible illustration of his remarks can be cited than the recent great abundance of the Snowy Owl in New England. This bird is regarded as not a rare winter visitor to New England,

---

\* Eight specimens.

where it is confined mostly to the coast, although occasionally taken throughout the interior. About the first of November, 1876, however, large numbers suddenly appeared along our coast. This being the season when sportsmen and the market gunners were in pursuit of water-fowl on the sea-shore, dozens of Snowy Owls were shot by them and sent to the markets and to taxidermists, so that during the three following weeks it was a common thing to see them hanging with other game in the markets, or confined alive.

I first heard of them on our Massachusetts coast as frequenting the islands off Rockport, where numbers were taken. One gunner spoke of seeing fifteen at once on a small island one foggy morning, nearly half of which he procured. As the Owls flew around over the rocks uttering their weird cries, they presented a scene of rare occurrence in New England. Specimens were soon after captured in nearly every town in this vicinity (Boston), and were sent to the city from various other parts of the country. Several were shot in the very heart of the city of Boston, where they were occasionally seen perched upon the house-tops or church-spires.

I learn from Mr. George A. Boardman, of Milltown, Me., that they were at this time very abundant in his locality, where they appeared as early as September. Mr. Simeon F. Cheney, of Grand Menan, also informs me that they were never before so abundant there as during the present season, arriving there about October 20. He reports that eight were seen together at one time, and that on another occasion a flock of fifteen was noticed.

Mr. N. C. Brown, of Portland, Me., reports that about one hundred and fifty were shot in the immediate vicinity of that city, and that five flew about the buildings of the city for a week unmolested. Mr. J. M. Le Moyne also writes me that the unusual abundance of these birds about Quebec, Canada, has been the subject of general remark.

The migration seems also to have extended far to the southward of New England, as I learn from Mr. Boardman that specimens have been taken as far south as Philadelphia, Baltimore, and Washington. In Philadelphia Mr. John Krider, the well-known taxidermist, had forty sent to him for preparation during October and November. One was taken near Baltimore during the last of September. I have heard of some five hundred specimens that have been seen, the majority of which have been shot.

Many of the specimens were in exceedingly poor condition. Of

some two hundred of these Owls examined by me, nearly all were in very dark plumage, and none wore that almost spotless dress which we occasionally see.

The cause of the sudden visit of such an unusual multitude of these boreal birds, coming as they did when the weather for a few days was unusually warm for the season, the thermometer standing at 75° at noonday, is a question not easily solved. Scarcity of food would seem the most probable solution, or perhaps an early severe cold snap started them on their southward flight. If so, it seems strange that other less hardy species should not be affected in a similar way, as but few Geese and Brant had passed south when the Owls had been with us for a week.

About ten years since there was a somewhat similar migration of this species into the British Provinces and New England, but the birds appeared later in the season, and not in such great numbers as in the present instance.

---

---

DISTRIBUTION OF NEW ENGLAND BIRDS. — A REPLY TO  
DR. T. M. BREWER.

BY H. A. PURDIE.

WHEN asked, some time since, to review a "Catalogue of the Birds of New England," by Dr. T. M. Brewer, I at first declined, feeling that if I expressed myself conscientiously, I should give some displeasure to its author. But I finally consented, and penned the short article in the third number of this Bulletin, bearing the signature "H. A. P." I intended to give the writer of this Catalogue all the credit due him, but in this, according to his reply in the following number, I have signally failed. The tone of Dr. Brewer's article, and the demand he makes that I must produce something of more weight than "unsupported assertions," "sweeping generalizations," opinions and conjectures unsupported by facts, and "positive dogmas given out quite *ex cathedra*," renders it necessary for me, in defence, to reply somewhat in detail. He must, however, be aware that reviews are generally limited as to space, and especially so was the case in this instance, so that full citations in support of my differences with him were out of place. He now

says he gave the list for what it was worth, expecting and desiring to have it amended and improved; and elsewhere he has said, "however lenient we may be even towards errors and incorrect statements that apparently might have been avoided, we should also, all of us, never hesitate to expose and to correct whatever we know to be wrong." Unhappily such an attempt on my part has not been very graciously received. He also says "it was but an initiative towards a complete and reliable list of the birds of New England," and that it was "at the last moment, and when it could only be done briefly," that "the character of the presence of the species" was added, and that of course the additions were "never exhaustive." Now I will respectfully ask how the general reader was to know this. I *received* the list "for what it was worth." There was nothing in the introduction to show that it was not considered complete or correct.

Of course, after all that has been written upon the subject, I was aware that generally a species is not resident *individually* in a given section of territory. His remark respecting the Robin, that the birds found with us in winter are not the same as those that pass the summer here, but "are of a very different *race*," is not at all to the point at issue. As a species, I say *Turdus migratorius* is a resident of New England. If, however, as he holds, the birds found in winter are another and very different race, — as race is now understood, — he should have so indicated it in the Catalogue, perhaps as "*Turdus hyperboreus*. Arctic Robin. Winter visitant." I intended in reply to cite at length all the facts that bear upon the points in dispute, but found that to do so would require quite too much of the valuable space of the Bulletin; but I trust that I have brought forward sufficient evidence to show that my statements were not altogether "conjectures" and "unsupported assertions."

The following five species, among others, Dr. Brewer claimed had never been taken in New England, and therefore should be excluded from the list of New England birds. In respect to this, I simply asked "if previous record did not show that these at least could be retained," intending thereby to imply that I considered this to be the case. The following are the birds and their record:—

**Quiscalus major.** BOAT-TAILED GRACKLE. — "*Q. baritus*, Bonaparte. Thrush Blackbird. New Haven. Of the Thrush Blackbird one specimen only has been observed, by Dr. Whelpley at New Haven, and of

course is rare in Connecticut." (Rev. J. H. Linsley, Cat. of the Birds of Conn., in Am. Jour. Sci. and Arts, Vol. XLIV, 1843, p. 249.) "Accidental. Have heard of one that was killed in Cambridge a few years since. Mr. E. A. Samuels tells me that a pair bred in Cambridge in 1861." (J. A. Allen, Proc. Ess. Inst., IV, p. 85, 1864.) Both these and the Connecticut bird are cited by Dr. Coues (Proc. Ess. Inst., V, p. 285, 1868) as valid. But I understand that more recently the authenticity of the specimens taken is doubted, they being referred to the Crow Blackbird (*Q. purpureus*). As Mr. Linsley also gives *Q. purpureus* as common, I see no reason for doubting his record. Of Mr. Samuels's birds, I have always understood him, and he still avers that two of them, in the flesh, were brought to him by Professor Jeffries Wyman, and that to his best knowledge and belief they were shot in the Cambridge salt marshes; that their rarity was commented on at the time, and that they were not *Q. purpureus*.

**Corvus ossifragus.** FISH CROW. — "Stratford," Conn., Linsley (l. c.). "An occasional visitor along the southern coast of the State of Massachusetts." (J. A. Allen, l. c.) "Very rare visitor in summer" to Massachusetts. (E. A. Samuels, Descriptive Catalogue of the Birds of Mass., in Rept. of Sec. Board of Agriculture of Mass. for 1863.) "A rare summer visitor, chiefly along the more southern portions of the coast" of New England. (Coues, Proc. Ess. Inst., V, 1868.) "Coast of the United States, from New England to Florida." (Coues, Birds of the Northwest.) And now, as confirmatory of the above, Mr. Brewster gives an instance of his seeing it in Cambridge, March 16, 1875. (See this Bulletin, Vol. I, p. 19.)

**Ægialitis wilsonius.** WILSON'S PLOVER. — Allowing that this species has not yet been found in Massachusetts, we have: "Stratford," Conn., Linsley (l. c.). "Appears to be rather rare, and perhaps only occasional, as far north as Massachusetts. (Coues, l. c.) "Dr. Wood informs me that Wilson's Plover is abundant in August on Long Island, and Mr. Linsley has recorded it from Stratford, Conn. It hence seems unquestionable that they sometimes occur in Southern New England, and it would not be strange if they should occasionally reach the coast of Massachusetts." (Allen, Am. Nat., Vol. III, 1869.) "North to Long Island and Connecticut, probably to Massachusetts, but rare beyond New Jersey." (Coues, Birds of the Northwest.)

**Nettion crecca.** EUROPEAN TEAL. — At a meeting of the Boston Society of Natural History, April 18, 1855, Dr. Bryant remarked that a specimen of the European Green-winged Teal had been sent to Mr. E. Samuels to be mounted. It was shot in Massachusetts, the first he had ever seen in the State. (Proc. Bos. Soc. Nat. His., Vol. V, p. 195.) This instance is cited by Allen (Proc. Ess. Inst., Vol. IV, 1864) and by C. J. Maynard (Naturalist's Guide, 1870). "A European species, but so often

taken on the coast as to be fairly considered as more than an accidental visitor" (Coues, Proc. Ess. Inst., Vol. V, 1868.)

**Sula fiber** BOOBY GANNET. — "Guilford," Conn., Linsley (l. c.). "September. Rare." Essex Co., F. W. Putnam. (Proc. Ess. Inst., Vol. I, 1856.) Both these cases are cited by Dr. Coues in his Birds of New England, and the last by C. J. Maynard. (Naturalist's Guide, 1870.)

I think that the above-named five birds have as much right to be included among those that have occurred with us as have the Mealy Red-Poll (*Egypthos canescens*), Small-headed Flycatcher (*Megodytes minimus*), Willow Ptarmigan (*Lagopus albus*), White-fronted Goose (*Anser gambeli*), and Hutchin's Goose (*Bernicla hutchinsi*), which are all retained by Dr. Brewer. There are also two other birds, namely, the Blue-gray Gnatcatcher (*Poliophtila carulea*) and the Blue Warbler (*Dendroica cerulea*), expunged by Dr. Brewer, whose record of occurrence in New England is as good as any just cited, which I am prepared to show have recently been taken in Connecticut and Rhode Island. (See this number of the Bulletin, p. 20.)

In regard to the Robin (*Turdus migratorius*), the Crow (*Corvus americanus*), the Hairy Woodpecker (*Picus villosus*), the Long-eared Owl (*Otus wilsonianus*), the Short-eared Owl (*Brachyotus cassini*), the Acadian Owl (*Nyctale acadica*), and the Sharp-shinned Hawk (*Nisus fuscus*), which Dr. Brewer classed as summer residents of New England (he saying of the Crow that a few winter), but which I stated must be considered as constant residents, at least of the southern portions, and some of them also of the northern, I will say that a reference to the often cited local lists, and to other writings on New England birds, quite fully confirms my remark, these species being given as residents, some of them occurring in smaller numbers in winter, as is to be expected, while again others are found more frequently in winter than at any other season. I notice a slight exception in the case of *Nisus fuscus*, a few writers regarding it as only a summer visitant, even to Massachusetts; but that it, as well as the others, remains in greater or less numbers in certain sections the year round, is well known to collectors.

In the "History of North American Birds," by Messrs. Baird, Brewer, and Ridgway, we find the following respecting the winter distribution of the Robin: "In the winter months it is most abundant in the Southern States, while in the Middle and even the Northern States, in favorable localities, it may be found throughout the year; its migrations being influenced more by the question of food than of climate. In the valleys among the White Mountains, where snow covers the ground from October to June, and where the cold reaches the freezing-point of mercury, flocks of Robins remain during the entire winter, attracted by the abundance of berries. In Massachusetts a few Robins remain throughout the year, but the greater proportion leave early in November, returning late in February or early in March."



Also in respect to the Hairy Woodpecker, we read in the same work (Vol. II, p. 506) : "It is a resident and not a migratory species, and wherever found it also breeds."

Also (in Vol. III, p. 46) of *Nyctale acadica* : "Mr. Boardman and Professor Verrill both give it as resident, and as common in Maine."

Other species, namely, the Short-billed Marsh Wren (*Cistothorus stellaris*), the Warbling Vireo (*Vireo gilvus*), the White-eyed Vireo (*V. noveboracensis*), the Field Sparrow (*Spizella pusilla*), the Carolina Dove (*Zenaidura carolinensis*), and the Quail (*Ortyx virginianus*), which were given as summer residents, presumably of all New England, I said seldom reached Northern New England. With the exception of *Ortyx virginianus*, I did not make the positive statement that the above-named species never did so, knowing that one or two of them had been found sparingly at certain localities in that section. Here again, the published record, with but slight exceptions, supports me in my assertion. As to *Vireo gilvus* and *Spizella pusilla*, though given by Mr. Verrill as summer visitants at Norway, Me., and by Dr. Coues as summer visitants to all New England, the former is rare, and the latter does not occur at all, at Calais, Me., nor does C. J. Maynard give either as found in Coos County, N. H., or Oxford County, Me. He considers the White Mountain range as forming their northern limit of distribution. Mr. William Brewster did not find them at Franconia, N. H.

Respecting *V. noveboracensis*, I quote the following from the "History of North American Birds" (Vol. II, pp. 385, 386) : "In the last-named State [Massachusetts] it becomes exceedingly rare, and beyond it is apparently not found, none having been met with either by Messrs. Verrill or Boardman in any part of Maine. Mr. Audubon states that he himself found them along the coast in Maine, Nova Scotia, and Labrador. This, however, I am inclined to consider a misstatement, as they have not since been detected north of the 42d parallel."

From the same work (Vol. II, p. 5), respecting *S. pusilla*, is the following : "In the summer it breeds from Virginia to Maine, as far as the central and western portions. It is not found near Calais, but occurs and breeds near Norway, Oxford County."

Again of *Z. carolinensis* (Vol. III, p. 384) : "It is found in the southern part of Maine as far to the eastward as Calais, but was not collected by Mr. Verrill at Norway, and is not known to occur in the northern part of that State." I said it was not "rare," meaning of course in Southern New England, and by looking up the matter, such will be found to be the case. Citing again from our standard work on North American birds, we find this of *Ortyx virginianus* : "This bird is probably found in all the New England States, though its presence in Maine is not certain, and if found there at all, is only met with in the extreme southwestern portion. It is also rare in Vermont and New Hampshire, and only found in the southern portions. It is not given by Mr. Boardman, nor by Professor Verrill."

Observations made in the nesting-season during the last five or six years in New Hampshire and Maine, by such experienced collectors as Messrs. Brewster and Maynard, and, to a less extent, by Bailey, Deane, and myself, show the absence of the foregoing species from the Fauna, and the presence of the five following, namely, the Cape May Warbler (*Perissoglossa tigrina*), the Mourning Warbler (*Geothlypis philadelphia*), the Olive-sided Flycatcher (*Contopus borealis*), and, in less abundance, the Black-backed and the Banded Three-toed Woodpeckers (*Picoules arcticus* and *P. americanus*).

What had already appeared in print respecting the distribution of these birds, added to the observations of the above-named gentlemen, I thought warranted me in saying that at least *P. tigrina*, *G. philadelphia*, and *C. borealis*, were "generally," that is, usually, if, perhaps, not universally, common, and bred regularly in Northern New England.

The Worm-eating Warbler (*Helmitherus vermivorus*), the Blue-winged Yellow Warbler (*Helminthophaga pinus*), the Yellow-breasted Chat (*Icteria virens*), the Hooded Warbler (*Myiodiocetes mitratus*), and the Great-crested Flycatcher (*Myiarchus crinitus*) have generally been considered rare birds in any part of New England, but in the "American Naturalist" (Vol. VII, 1873, p. 692) I mentioned, on the authority of Mr. J. N. Clark, of Saybrook, Conn., that they were found at that locality regularly in numbers and breeding, though he had not actually found the nest of *H. vermivorus*. I also spoke of a Water-Thrush that occurred there, and inferred that it was probably the Long-billed Water-Thrush (*Siurus ludovicianus*). Subsequent correspondence, and a visit to Saybrook in June, 1875, confirmed my suspicion, proving that *ludovicianus* was the species that summered there, and that it was common, as were all the others, with the exception of *H. vermivorus*. Mr. C. M. Jones, now of Eastford, Conn., has written me that he observed all but *S. ludovicianus* and *H. vermivorus*, at Madison, in that State, where he formerly resided.

The Golden-winged Warbler (*Helminthophaga chrysoptera*) and the Yellow-winged Sparrow (*Coturniculus passerinus*) are two species that Dr. Brewer still denies can be considered as breeding regularly, or in numbers, in any portion of New England. As far back as June, 1869 (Am. Nat., Vol. III, p. 497), and again in 1870 (Samuel's Orn. and Oöl. of New England, revised edition, 1870, Appendix), I showed that the Golden-winged Warbler was far from uncommon in Massachusetts. Observations made every year since have not altered my opinion. I find it in the proper places from May to August. It is apparently less plenty after the first of June, but is still not a bird of the Canadian Fauna.

In "History of North American Birds" (Vol. I, p. 193) we read: "Occasionally specimens have been obtained in Massachusetts, and, of late, these occurrences have become more common or more observed. . . . Mr. J. A. Allen has known of several specimens taken within the State. Mr. Jillson has observed it spending the summer in Bolton, and evidently

breeding, as has also Mr. Allen at Springfield, and Mr. Bennett at Holyoke." (See also Am. Nat., Vol. III, 1869, p. 575; Maynard's Naturalist's Guide, 1870; and this Bulletin, Vol. I, p. 6, for accounts of the nesting of this species in Massachusetts.)

I had no idea that any one acquainted at all with New England birds could say that *C. passerinus* was rare, or even uncommon, in Southern New England. Why, it absolutely swarms, so to speak, on Nantucket. I presume Dr. Brewer will allow that island to be included within our limits. On Cape Cod, and, indeed, in various portions of Massachusetts, Rhode Island, and Connecticut, and even northward to Concord, New Hampshire, it may be found in plenty at all suitable localities. At Saybrook, Conn., its notes were to be heard in every field. (See History North American Birds, Vol. I, p. 554, and local lists of New England birds, south of Northern sections, in confirmation of this statement.)

The Long-billed Curlew (*Numenius longirostris*), the Yellow Rail (*Porzana norboracensis*), and the Coot (*Fulica americana*) I considered spring and fall migrants, rather than as summer residents. The lists show this statement also to be true, while the gunners and collectors further confirm it. Perhaps a few may summer on the extreme northeastern coast of Maine.

But my space is becoming limited. That the Golden-crested Kinglet (*Regulus satrapa*) winters in numbers in Southern New England, that the Snowbird (*Junco hyemalis*) does not do so in Northern New England, that the Titlark (*Anthus ludovicianus*) does not winter (perhaps with rare exceptions in the southernmost parts), and that *Ectopistes migratoria* regularly summers in different portions of New England, are all statements demonstrable by facts already in print, and by the observations of those who speak of that which they do know.

A word about the Stilt Sandpiper (*Micropalma himantopus*), and I am done. In the "American Naturalist" (Vol. III, p. 639) is recorded the first supposed instance of its occurrence in New England. In the same periodical (Vol. VII, p. 727) is given the first supposed\* instance for Massachusetts. Again (in Vol. VI, p. 307) Mr. Brewster says: "The Stilt Sandpiper (*Micropalma himantopus*), which I see was recorded in a recent number of the 'Naturalist' as new to our Fauna, I consider by no means rare in its migrations. Indeed, I have seen as many as six or seven sent into Boston market at one time, from Cape Cod, and, in the course of a few weeks' shooting in August, at Rye Beach, N. H. (just north of our State limits), secured no less than ten specimens." Not only has he since shot it, but he, as well as myself and others, find it frequently in the Boston markets.

---

\* Mr. F. C. Browne, of Framingham, has a specimen taken at Plymouth in 1852.

OCCURRENCE OF THE BARNACLE GOOSE (*BERNICLE LEUCOPSIS*) ON LONG ISLAND, N. Y.

BY GEO. N. LAWRENCE.

I WAS recently informed, by Mr. Harold Herrick, that a specimen of this species could be seen at the store of Mr. Conway, taxidermist, in Carmine Street, said to have been killed on Long Island. I called there and was shown a nicely mounted example of this Goose in perfect plumage. Mr. Conway said that it was brought to him in the flesh, in good condition, and was eaten by his family; he spoke very favorably of its edible qualities.

I learned from him that its possessor was Mr. J. K. Kendall of this city. I had an interview with this gentleman, and requested that he would ascertain all the facts possible as to its capture, and send me the information. I received from him the following letter giving the result of his inquiries:—

NEW YORK, November 29, 1876.

DEAR SIR,—About October 20 I saw a specimen of the Barnacle Goose hanging in a restaurant in this city,—bought it and had it stuffed. I questioned the proprietor, and learned from him the place where he bought it,—from a produce-dealer near Washington Market. Afterwards I interviewed the marketman, and he recollected the bird well, although he had no idea what it was. He told me he bought it from a Long Island farmer, who brought it to the city in his wagon, and who said that it was killed by a boy in Jamaica Bay. Unfortunately he did not know the farmer,—never saw him before nor since, so I was unable to trace the bird any farther, but I am fully satisfied the story was true.

Yours truly,

J. K. KENDALL.

This is the second instance of this species having been procured on the Atlantic coast; the first was obtained in Currituck Sound, North Carolina, in 1870, and is recorded in Vol. V, p. 10, of the "American Naturalist."\*

---

\* In Dr. Brewer's "Catalogue of the Birds of New England" (from Proceedings of the Boston Society of Natural History, Vol. XVII, March 3, 1875) he excludes this species from our New England list, and also states that the specimen recorded by Mr. Lawrence as having been taken in North Carolina was

CAPTURE OF A SECOND SPECIMEN OF HELMINTHOPHAGA  
LAWRENCEI.

BY HAROLD HERRICK.

IN 1874 I had the pleasure of publishing in the "Proceedings of the Academy of Natural Sciences of Philadelphia" (p. 220, pl. xv) a description of a new species of *Helminthophaga* that I had just been fortunate enough to unearth. It has remained unique up to the present time, and although its friends have stoutly maintained its validity, the "hybrid" theorists have sorely tried their faith; therefore I am more than pleased to be able to set the matter permanently at rest by announcing the capture of a second specimen of *Helminthophaga lawrencei*. The specimen, oddly enough, was secured by Mr. Lawrence himself, who sends it to me with a letter of explanation, from which the following is an extract:—

"I obtained the specimen of *H. Lawrencei* last fall from a dealer, who called my attention to it as having a black throat, differing in that respect from any species he had ever before met with. He said it was sent to him last spring from Hoboken, N. J., with a miscellaneous lot of Warblers. I think the acquisition of a second specimen of this species should put at rest all doubt of its validity."

This specimen agrees precisely with the type, with this slight exception, that the type is an adult male, probably in the second or third year, while the bird under consideration is unquestionably a yearling male, and still has the immature yellowish tips to the coal-black feathers of the throat-patch. A slightly similar effect is seen in the yearling males of *Dendræca virens*. I cannot better describe it than by republishing the description of the type.

---

probably one of eight specimens which escaped from the grounds of a gentleman in Halifax in the fall of 1871 or 1872.

From Mr. Lawrence's record (*Am. Naturalist*, Vol V, p. 10) we find this Goose was captured on October 31, 1870, one or two years previous to the escaping of the Halifax birds.

In view of this fact may not Mr. Lawrence's specimen still remain as the first authentic instance of the occurrence of the Barnacle Goose in the United States; at all events, until we hear of a confined specimen having escaped previous to that date!—RUTHVEN DEANE.

“Upper parts and rump olive-green, a shade darker than in *pinus*. Wings bluish-gray, with two white bands, the upper not so clearly defined as in *pinus*. Tail bluish-gray, with the three outer tail-feathers with most of the web white, also a small white spot on the end of the fourth feather. Crown and under parts, from breast to vent, orange. A broad black patch extends from the bill through and behind the eye. Chin, throat, and fore-part of the breast black. A yellow stripe, commencing under the bill, extends back between the black eye- and breast-patches, and increases in width upon the shoulder. Length, 4.50; wing, 2.50; tail, 2.00. Measurements from the mounted bird.”

The measurements of the two birds are as nearly identical as is possible when one bird is mounted and the other a skin. Of its habitat, the plumage of the female, and its nesting peculiarities, we can only conjecture, but it seems not unreasonable to presume that its habitat is similar to that of its near congener, *H. pinus*, and that New Jersey may some day produce its nest and eggs, as it has already produced the only two known specimens of the bird.

The female, I believe, will be found to be not unlike that of *H. pinus*, and a close inspection of supposed specimens of the latter bird now in collections may develop some interesting facts.

In conclusion it may be well to add, what by inadvertence I omitted when the description was first published, namely, that for the correct delineation of the bird in the plate I am indebted to Mr. Robert Ridgway, of the Smithsonian Institution, to whom I take this opportunity of tendering my thanks.

---

#### NOTICE OF A FEW BIRDS OF RARE OR ACCIDENTAL OCCURRENCE IN NEW ENGLAND.

BY H. A. PURDIE

THE following data respecting the occurrence of the following fourteen species so far to the northeastward as New England are of special interest. I am indebted to Messrs. Frederic T. Jencks of Providence, R. I., Erwin I. Shores of Suffield, Conn., F. C. Browne of Framingham, Mass., and J. N. Clark of Saybrook, Conn., for facts relating to eleven of the birds here mentioned.

1. *Polioptila cærulea*. BLUE-GRAY GNATCATCHER. — Mr. Jencks writes: “Two were shot at Wauregan (Windham County), Conn., by

Mr. C. M. Carpenter, — a male in 1874 and a female in 1876. Three or four were seen by me at Providence, R. I., May 23, 1875."

A male was also shot, by Mr. Shores, at Silver Spring, near Providence, June 24, 1875, and several others have been seen by him at different times in Providence and vicinity.

2. *Helmitherus vermivorus*. WORM-EATING WARBLER. — Mr. Shores shot a male at Suffield (Hartford County), Conn., August 22, 1874. This is, I think, its most northerly record in the Atlantic States yet noted.

3. *Helminthophaga celata*. ORANGE-CROWNED WARBLER. — Mr. Jencks writes me that "a specimen was shot in Cranston, R. I., December 3, 1874." This is the fifth specimen reported for New England, and the second taken in the winter season.\*

4. *Dendroeca cærulea*. BLUE WARBLER. — A male was obtained by Mr. Shores at Suffield, June 12, 1875. This species and *Poliophtila cærulea*, though previously recorded as occurring in New England, have not been recently taken here.

5. *Myiodiocytes mitratus*. HOODED WARBLER. — A male was shot at Suffield, Conn., by Mr. Shores, July 8, 1875. This bird, though found regularly along the Sound shore of Connecticut, has not been noticed so far northward before in New England. This, as well as a few other species characteristic of the Carolinian Fauna, will probably be found to extend up the river-valleys of Connecticut, though not passing farther eastward.

6. *Pyrranga æstiva*. SUMMER REDBIRD. — Mr. Jencks informs me that a male was shot a few years since on Ten-Mile River, six or eight miles northeast of Providence. It has appeared before, but is sufficiently rare here to merit notice.

7. *Stelgidopteryx serripennis*. ROUGH-WINGED SWALLOW. — A female of this species was shot at Suffield, Conn., by Mr. Shores, June 6, 1874. At last this bird has been taken within our limits. It will be interesting to determine whether it proves to be in future a regular visitant to New England.

8. *Collurio ludovicianus* var. *excubitoroides*. WHITE-RUMPED SHRIKE. — A typical example of this variety was shot by Mr. Jencks in Cranston, R. I., September 2, 1873, and is now in his collection. Its previous record of having been found within our borders is somewhat doubtful. I believe it is hardly found regularly much east of Buffalo, N. Y. In this connection I would say that the *Collurio* taken in Massachusetts, recorded by me in the "American Naturalist" (Vol. VII, 1873, p. 115), was a typical "Loggerhead" Shrike (*C. ludovicianus*).

9. *Milvulus forficatus*. SWALLOW-TAILED FLYCATCHER. — Mr. Jencks informs me that a specimen of this species was shot by Mr. Carpenter, at

---

\* See this Bulletin, Vol. I, p. 94, for its previous New-England record.

Wauregan, Conn., about April 27, 1876. The bird first attracted Mr. Carpenter's attention by its opening and closing the tail while flying about a small sheet of water in quest of insects. The only other Eastern United States capture of this species is a male taken at Trenton, N. J., a few years ago, as recorded by Dr. C. C. Abbott.\* Of course its appearance here is entirely accidental.

10. *Porzana jamaicensis*. BLACK RAIL. — I have lately seen a skin of this species belonging to Mr. Browne, of Framingham. The bird was picked up dead, in August, 1869, by a relative of his, on Clark's Island, Plymouth Harbor, and was forwarded to him as something entirely new to our shores. This instance adds a new bird to the Fauna of Massachusetts.

Of this species Mr. Clark, of Saybrook, Conn., also writes me that a neighbor of his, while mowing at that place, July 10, 1876, swung his scythe over a nest of ten eggs on which the bird was sitting, unfortunately cutting off the bird's head and breaking all but four of the eggs. The only previous New England record of this species is that given by Dr. Brewer (Proc. Bost. Soc. Nat. Hist., Vol. XVII, p. 477).

11. *Rallus longirostris*. CLAPPER RAIL. — In the Natural History store of Brewster & Knowlton, Boston, I recently saw a mounted specimen of this species. The bird was captured by its flying on board a vessel in the harbor, May 4, 1875. Though recorded from Maine and Connecticut, it being in the latter probably quite a regular summer visitor (about the Sound), I believe its appearance before in Massachusetts has been questioned.

12. *Rallus elegans*. KING RAIL. — In the collection of Mr. George O. Welch, of Lynn, Mass., is a mounted specimen shot at Nahant, November 21, 1875. This is a second species new to Massachusetts, and has been but once or twice before recorded from any part of New England.

13. *Sterna fuliginosa*. SOOTY TERN. — Mr. Clark informs me that he has this species in his collection, mounted from a bird that last summer flew against the side of the steamboat-wharf depot at Saybrook, Conn. Stunned by the concussion, it fell and was picked up. It had been noticed for several days flying about the mouth of the river as something unusual.

14. *Pelecanus trachyrhynchus*. WHITE PELICAN. — At the Natural History store of Mr. A. J. Colburn, Boston, I saw, a few months since, a skin of this species, freshly made up from the flesh. The bird was shot at North Scituate, October 6, 1876, by Mr. George Pratt. It was a male, in fine plumage and good condition. Though not new to the State, I think its presence with us worthy of notice.

---

\* Amer. Nat., Vol. VI, p. 367.



## Recent Literature.

NOTICES OF FIVE RECENT ORNITHOLOGICAL PAPERS. — The first three numbers of the "Proceedings of the Zoological Society of London" for 1876 contain several important papers upon the anatomy and classification of several groups of birds, by Mr. A. H. Garrod, while among the numerous other ornithological articles of more or less special interest are papers by Mr. Howard Saunders, on the Skau Gulls and on the Terns; by Messrs. Sclater and Salvin, on the *Anatidae* of "Neotropical" America; an abstract of a memoir by Mr. W. K. Parker on *Ægithognathous* birds; and a paper by Mr. W. H. Hudson on the habits of some of the Rails of the Argentine Republic. Among the numerous new species of birds figured and described are quite a number from the Andean Region of South America.

Among Mr. Garrod's contributions is a short paper (l. c. pp. 275-277) on the anatomy of the Courlan (*Aramus scolopaceus*). He finds it to have, on the whole, decidedly closer affinities with the Cranes (*Grus*) than with any other group, especially in respect to its osteology, notwithstanding its many external resemblances to the Rails. Hence Mr. Garrod's researches confirm the views of recent systematists in respect to the affinities of this peculiar and interesting form. Mr. Garrod also writes (l. c. pp. 335-345, pls. xxvi-xxviii) concerning the anatomy of the Darter (*Plotus anhingæ*), a bird whose anatomy, aside from its skeleton, had previously received little attention. Mr. Garrod finds in its visceral anatomy several quite peculiar features, one of which is the protection of the pyloric orifice by "a mat of lengthy hair-like processes, much like cocoanut fibre, which nearly half fills the second stomach." These hair-like fibres are found to considerably resemble in structure true cutaneous hairs. In general terms, the Darter may be said to present many of the features characteristic of the Gannets, Pelicans, and their allies, in an exaggerated degree.

The most important and interesting of Mr. Garrod's contributions is a paper "On some Anatomical Characters which bear upon the Major Divisions of the Passerine Birds" (l. c. pp. 506-519, pls. xlviii-lviii). Mr. Garrod attaches great importance to the mode of insertion of one of the muscles of the wing (the *tensor patagii brevis*), to the character of the syrinx, the absence or presence of either the femoral or the sciatic artery, etc., to which points the researches here detailed are mainly directed. He concludes his paper with a tabular arrangement of the larger groups of the Passeres, expressive of his views of their affinities.

Mr. Howard Saunders's paper (l. c. pp. 317-332, pl. xxiv) on the Skau or Jager Gulls (*Stercorariina*) is devoted to the synonymy and range of the species, with incidental remarks on their progressive changes of plumage. Mr. Saunders recognizes six species, all of which he refers

to one genus, for which he adopts the name *Stercorarius* as being the only proper and tenable one. Two of these species belong to the Southern Hemisphere, the other four to the Northern, all of which latter occur in the boreal parts of North America, as well as in the Old World. The nomenclature adopted by Mr. Saunders for two of the Northern species differs from that commonly employed, Mr. Saunders maintaining, and apparently with good reason, that the Linnaean name *parasiticus* belongs not to the bird commonly so called, but to the bird usually called *capphus* or *buffoni*. Thus *Stercorarius eripidatus* Saunders is the *S. parasiticus* of Coues and most recent authors, while the *S. parasiticus* Saunders is the *S. buffoni* of Coues and others, which is again the *S. capphus* of Gray and other writers.

Mr. Saunders's paper on the Terns (l. c. pp. 638-672, pl. lxi) is similar in character to that on the Jager or Skau Gulls, treating mainly of nomenclature and distribution, being, like the other, preliminary to a monograph of the *Laridae*. The genera recognized are *Hydrochelidon*, *Sterna*, *Nania*, *Gygis*, and *Anous*. Of the forty-eight species recognized, thirty-eight are placed under *Sterna*. The most noteworthy change of names among the North American species is the substitution of the name *furiatilis* of Naumann for the hitherto almost universally accepted *hirundo* of Linnaeus for our Common Tern, which name he considers as originally embracing both the *hirundo* and the *macrura* of recent authors. The *Sterna portlandica* of Ridgway is referred to *S. macrura*, in accordance with Mr. Brewster's views, and the Least Tern is considered as specifically distinct from *S. superciliaris*, of which Dr. Coues deems it to be merely a variety. In most other cases Mr. Saunders's names as respects the North American species agree with those recently adopted by Dr. Coues in his "Birds of the Northwest."

Messrs. Sclater and Salvin's "Revision of the Neotropical Anatidæ" (l. c. pp. 358-412, pl. xxxiv) is a most valuable synopsis of the Ducks and Geese of Middle and Southern America, and embraces also a large proportion of the species of North America, including as it does all that reach Tropical America in their migrations. The paper includes notices of sixty-two species, of the greater part of which are given short descriptions, accompanied by pretty full lists of bibliographical references. Twenty-three of the species are considered as "Nearctic," leaving thirty-nine as properly "Neotropical." The paper closes with a very convenient tabular synopsis of the geographical distribution of the genera and species. — J. A. A.

VENNOR'S RAPACIOUS BIRDS OF CANADA. — With the title "Our Birds of Prey; or, The Eagles, Hawks, and Owls of Canada," by Henry G. Vennor, Messrs. Dawson Brothers have published an elegant royal octavo volume of one hundred and fifty-four pages, with thirty photographic illustrations. While these illustrations are probably as excellent as the photo-

graphic art can supply, they cannot be regarded as a very valuable addition to the work, certainly not in proportion to their cost. They do not supply those shades of tinting so essential to the student, and, being necessarily taken from mounted specimens, cannot remedy the inevitable shortcomings of their models. The text, which is largely compiled from the notes of other writers, gives a fairly digested summary of the individual history of each species.

Mr. Vennor includes two forms of Gyr-falcons, the *candicans* and the *labradora* of Audubon, but adds nothing of moment to our knowledge of the history of the former, and does not include, except inferentially, *Hierofalco islandicus* as among the birds of Canada. He gives, as a separate form, the dark Gyr-falcon, described by Audubon as *labradora* but he is mistaken in several of his statements in regard to this variety. It is probably not so very rare a bird as has been supposed, although it is little known in North American collections. The supposition that the two specimens in the Montreal Museum are the only ones known in all North America is incorrect. Mr. Boardman of St. Stephen possesses at least two very fine specimens, the Boston Museum has a very fine one, and there is at least one in the National Museum of Washington. Nor is Mr. Vennor the first to represent, in plate, this species (or variety?).

In the "Ornithological Miscellany," edited by Mr. George Dawson Rowley, and published by Trubner & Co., of London, Mr. Henry E. Dresser presented a very interesting memoir of this Hawk, accompanied with a very fine illustration. I am not aware that any copy of this work is in this country, and the writer can only refer to it from memory. From this it would appear that for several years past collections of skins received in Europe from Labrador have always contained skins of this bird. One of the museums of Germany was especially fortunate in securing a fine series of this bird, and Mr. Dresser, having learned the source from which it had been enriched, has himself since procured several very fine specimens. So far as is known it seems to be confined to Labrador, and its specific peculiarities, if it has any, are not publicly known. At present we know too little in regard to it to discuss the question whether it is to be regarded as a species or a race, or whether it may not be a melanistic form. It is much more distinct, in its external markings, from any of the three other forms, *gyrfalco*, *islandicus*, and *candicans* than they are from one another, and, so far as is known, there is much less variation in the markings of individuals. The writer has no doubt that the birds referred to (North American Birds, Vol. III, p. 311), under the supposition that they belonged to the Black Rough-legged Hawks, were really of this group.

In this connection it may be mentioned that Mr. Dresser refers the form of *Hierofalco* found on Anderson River, not to *H. candicans*, but to the more common Norway form of *H. gyrfalco*. — T. M. B

## General Notes.

A HUMMING-BIRD NEW TO THE FAUNA OF THE UNITED STATES. — I have again the pleasure of adding another bird new to our Fauna. A Humming-Bird (male), taken within the limits of Fort Brown, Texas, August 17, 1876, and forwarded to the Smithsonian Institution for identification, has just been determined to be *Amazilia cervineiventris*. It much resembles *Pyrrhophæna riefferi*, and has rusty (instead of white) leg puffs. — JAMES C. MERRILL, Fort Brown, Texas, December 4, 1876.

NOTE ON PODICEPS DOMINICUS.\* — This species was long since attributed (perhaps erroneously) to "California," by Dr. William Gambel; it was included by Baird among Birds of the Mexican Boundary, apparently on strength of its eggs found at Matamoras, and figured in his "Birds of North America" (ed. of 1860, not of 1858). It was also formally presented by me as North American (Birds of the Northwest, p. 736, where its habitat is given as north of the Rio Grande). — ELLIOTT COUES.

EASTWARD RANGE OF THE FERRUGINOUS BUZZARD (*Archibuteo ferrugineus*). — During the past summer (1876) I found this bird to be common on the prairies of Nebraska and Wyoming, where it might almost be considered as one of the characteristic species. In 1873 I observed it on the Pembina Mountains, in Eastern Dakota, near the Red River of the North; and in 1874 I found it nesting in Northern Montana, on one of the Two Forks of Milk River. In years previous I had only seen it in Arizona and Southern California. I can now record its range still farther eastward, — beyond the Mississippi, as I lately saw one in Illinois, a few miles from the river. The great size of the bird, its white tail, almost as conspicuous as that of the Bald Eagle, and white under parts, render it unmistakable at any ordinary distance. Its geographical distribution is apparently nearly coincident with that of the Lanier Falcon (*Falco polyargus* Cass.), a bird which I have also found very numerous in Nebraska, Wyoming, and open portions of Colorado. Both species are prairie Hawks, subsisting largely or chiefly upon the small rodent mammals which abound in such regions. — ELLIOTT COUES, Washington, D. C., October, 1876.

OCCURRENCE OF LECONTE'S BUNTING (*Coturniculus lecontei* Don.) IN IOWA. — One of my correspondents, Mr. E. W. Newton, of Franklin Grove, Ill., writes me that when on a recent collecting trip through Iowa, he had the good fortune to secure twenty-two specimens of this species in a small slough situated in Colo, Story County, near the centre of the State, one of which he kindly sent me for identification. The date of cap-

---

\* See this Bulletin, Vol. I, p. 83, November, 1876.

ture recorded on the label is October 10, 1876. Although he hunted carefully over equally desirable situations in other parts of the State, this was the only place where it was found. This forms its most eastern record, excepting the single specimen taken by Mr. E. W. Nelson at Riverdale, Ill.\* — H. B. BAILEY.

AUDUBON'S WARBLER IN MASSACHUSETTS. — While collecting in the neighborhood of Cambridge, Mass., November 15, 1876, I was fortunate enough to obtain a fine specimen of Audubon's Warbler (*Dendroica auduboni*). It was a male, and the yellow of the throat was very plainly marked. Dr. Coues, in his "Birds of the Northwest," gives Laramie Peak as about the eastern limit of this species. Its occurrence here must, of course, be regarded as entirely accidental. — A. M. FRAZAR.

OCCURRENCE OF THE SOOTY TERN IN MASSACHUSETTS. — In Mr. Allen's "Catalogue of the Birds of Massachusetts" we find the Sooty Tern (*Sterna fuliginosa*) given, on the authority of Mr. E. A. Samuels, as a rare summer visitor to Muskeget Island. But for some reason Dr. Brewer, in his recent "Catalogue of the Birds of New England," withdraws this species from the New England list, and challenges its right to be regarded as in any sense a New England bird. I have the pleasure of replacing this species by recording the capture of a fine adult male on the Merrimack River near Lawrence, Mass., on October 29, 1876. I examined the specimen at the store of Mr. Charles I. Goodale, taxidermist, who has finely preserved it, and it is now in the possession of Mr. A. W. Howland of Lawrence. — RUTHVEN DEANE.

THE BLACK GYR-FALCON (*Falco sacer* var. *labradora*) IN MASSACHUSETTS. — A fine specimen of this Falcon was shot on Breed's Island during the latter part of October, 1876. It proved to be a male, in nearly adult plumage, and is now in the collection of Mr. C. I. Goodale, through whose kindness I have had the pleasure of examining it. — C. B. CORY.

NOTES ON BIRDS NEW TO THE FAUNA OF MAINE, ETC. — Of the following five species, three are here for the first time recorded as birds of Maine, another as found for the first time so far in the interior, and another as found for the first time breeding on the New England coast.

1. **Ammodromus caudacutus** Swain. SHARP-TAILED FINCH. — I have found this species, now, I believe, for the first time recorded as a bird of Maine, a rare inhabitant of a certain part of the great marsh in Scarborough.

2. **Passerculus princeps** Maynard. IPSWICH SPARROW. — On the 9th of October, 1876, I met with one of these birds on a sandy point on the northwest shore of Lake Umbagog, in New Hampshire. I should hesitate

---

\* See Bulletin of the Nuttall Ornithological Club, Vol. I, p. 40.


to record the occurrence of this species in a locality so far removed from its known haunts, it not having been before observed so far in the interior, since, from the miss-fire of two cartridges in succession, I failed to capture my bird, were I, not perfectly acquainted with its almost unmistakable habits.

3. *Strix flammea* var. *pratinctola* Bonap. BARN OWL. — Mr. L. C. Daniels, of this city (Portland), has in his possession a specimen of this owl which he shot in Falmouth, June 10, 1866. It was killed while flying across an open field. It has not, I think, been before recorded as found in Maine.

4. *Tringa bairdii* Coues. BAIRD'S SANDPIPER. — My brother, Mr. Philip G. Brown, shot a young male of this species as it was flying along Scarborough Beach, on September 9, 1875. It was in company with another bird, apparently of the same species, which escaped. This is its first recorded appearance on the coast of Maine.

5. *Thalassidroma leachii* Bonap. LEACH'S PETREL. — This Petrel breeds in large numbers on several of the outer islands of Casco Bay, southeast of Portland. Although I have often been told by fishermen and sportsmen of the existence of colonies of these birds on certain of our islands, I never was able to verify their reports until the middle of last August, when I made three visits to two barren rocks known as the "Green Islands," once in company with Mr. E. N. Atwood of Cape Elizabeth. I found about forty nests, half of which at this late date were empty, the remainder containing squabs in different stages of development. — NATHAN CLIFFORD BROWN, *Portland, Me.*, November 12, 1876.

NORTHERN RANGE OF THE SHARP-TAILED FINCH (*Ammodromus caudacutus*). — My friend, Mr. William Stone of Cambridge, has recently presented me with five specimens of the Sharp-tailed Finch which he shot at Tignish, Prince Edward's Island, on August 2 and 3, 1876. The locality where they were taken, as he describes it to me, was exceptional, — a wide waste of marsh, dry, and at some distance from the sea, grown up to bushes, with a few scattered dead pine stubs, remnants of a former forest. Throughout this tract these birds were abundant, the males singing on all sides from the tops of the bushes. The individuals examined are all adults in very pale, worn breeding plumage. Dr. Coues, in his "Birds of New England" (Proc. Essex Inst., Vol. V, p. 282), gives *Ammodromus maritimus* as occurring at Rye Beach, New Hampshire, but this record, he informs me by letter, was a mistake, the bird which he found there being *A. caudacutus*. The finding of the Sharp-tailed Finch in numbers at Tignish, taken in connection with the fact of its recent detection at Scarborough, Me., by Mr. N. C. Brown [see above], renders it extremely probable that it may occur regularly, at suitable localities, all along the intermediate line of coast, — WILLIAM BREWSTER.

 THE continuance of the Bulletin beyond the present Volume is necessarily contingent upon the pecuniary support it receives from the public. We trust that our present patrons will not only all promptly renew their subscriptions, but will use their influence to extend its circulation. We hope it will soon be possible to permanently increase the number of pages from twenty-four to thirty-two in each issue. In order to do this a considerable increase to our subscription-list will be necessary, and we therefore appeal strongly to all friends of Ornithology to aid in extending its circulation. This being the only journal in this country devoted exclusively to this subject, it is hoped that our appeal for pecuniary support will meet with a prompt response.

In order to assist in defraying the expenses of publication, advertising sheets will be issued with future numbers, relating mainly to Natural History, and especially to Ornithology. The attention of dealers in specimens of Natural History and collectors' and taxidermists' materials, and of publishers of works relating to Natural History, is hence called to the Bulletin as a desirable advertising medium.

## A NEW MANUAL OF TAXIDERMY :

An illustrated book, board covers, giving complete instructions in Taxidermy, besides Hints on Collecting and Preserving Eggs and Insects, and other valuable receipts.

**From J. W. P. JENKS, Prof. of Natural History at Brown University.**

“Having perused your MS., I feel free to say that its suggestions are eminently practical, and cannot fail to render such aid to the beginner as he most needs, and indeed must have from some source, at the outset of his efforts to acquire the beautiful art of preserving and mounting specimens of Natural History.”

Every NATURALIST and BEGINNER should get it. Sent, post-paid, on receipt of 50 cents.

W. P. MANTON, 17 BEACON STREET,  
*Boston, Mass.*



# BULLETIN

OF THE

## NUTTALL ORNITHOLOGICAL CLUB.

---

Vol. II.

APRIL, 1877.

No. 2.

---

CORRECTIONS OF NOMENCLATURE IN THE GENUS SIURUS.

BY DR. ELLIOTT COUES, U. S. A.

THE genus itself, and all three of its species, require names different from those now generally used.

I. NAME OF THE GENUS. — Originally written SEIURUS by Swainson, who invented the term; also found under the forms *Sciurus*, by ignorance, inadvertence, or typographical blunder, and *Siurus*, the latter being correct. The word is compounded of the Greek *σειω*, "I wave or brandish," and *οὐρον*, "tail"; it is precisely equivalent to the Latin *mota-cilla*, French *hoche-queue*, English *wag-tail*. According to the rule that Greek *ει* becomes long *i*\* in Latin, the word should be spelled *Siurus*, as was first done, I think, in the *Ibis* for 1859, by Messrs. Selater and Salvin, and A. and E. Newton, so nearly simultaneously that I do not know to which of these scholars we owe the corrected orthography. *Seiurus* has been objected to on account of its identity in sound, though not in orthography, etymology, or signification, with *Sciurus*, "a squirrel," by German purists, who have proposed to substitute *Enicocichla* or *Henicocichla*; but this is inadmissible: *Siurus* and *Sciurus* being as different as *thee*, objective case of second personal pronoun, and *the*, definite article.† (Lat. *Sciurus* = Gr. *σκίουρος* = "shadow-tail.")

---

\* So, also, *Melopelia*, *Chamapelia*, *chrysoparia*, etc. (accent the penult), instead of *Melopleia*, *Chamapelia*, *chrysoparia*, etc.

† I am not of those rigid constructionists who require preservation of the original shape of a name, however faulty. While we cannot of course make actual substitution of one name for another without other than philological

2. NAME OF THE GOLDEN-CROWNED THRUSH. — Originally and usually written *aurocapillus*, which should give way to *auricapillus*. The word means simply "gold-hair," i. e., "golden-haired." The point is here: that the ablative of *aurum*, "gold," which is *auro*, is only to be used when the word with which it is compounded is an adjective or participial; otherwise *auri* is the correct form. If we were to employ the participial adjective *capillatus*, it would be correct to say *aurocapillatus*, i. e., literally and correctly, "haired with gold," *auro* being the *ablativus instrumenti*, that with or by means of which the bird is "haired." So we say rightly *aurocristatus*, *aurostriatus*, *auropunctatus*, crested, streaked, or speckled with gold, (color understood), but *auriceps*, *auricollis*, *Auriparus*, etc. So also, if we were to compound with the adjective *aureus*, "golden," we should say, e. g., *aureicauda*, not *aureocaudatus*. Either *auricapillus* or *auricapillus* is correct, but *aurocapillus* is not.

3. NAME OF THE SMALL-BILLED WATER-THRUSH. — It is to be noted that *Motacilla noveboracensis* of Gmelin, 1788, is precisely the same as *Motacilla navia* of Boddaert, 1783, both being based upon Planche Enluminée 752, fig. 1, which is the *Fauvette tachetée de la Louisiane* of Buffon, afterward the *New York Warbler* of Pennant and Latham. G. R. Gray seems to have observed this fact, but neither he nor any other author, according to my recollection, has acted upon the obvious requirement of the case, namely, that we must say *Siurus navius* (Bodd.), instead of *S. noveboracensis* (Gm.). Very curiously, Gmelin in another place made this species out to be a variety of the Cape May Warbler, *Perissoglossa tigrina*; for, Gmelin's *Motacilla tigrina* var.  $\beta$  (and so, also, Latham's *Sylvia tigrina* var.  $\beta$ ) is based exclusively upon the *Ficedula dominicensis fusca* of Brisson, Ornith., iii, 513, which is the Small-billed Water-Thrush. Vieillot, in 1807, noticed this curious circumstance, which authors have generally overlooked, and correctly allocated the synonymy. The name *navius* is unobjectionable, has priority, and must obtain.

4. NAME OF THE LARGE-BILLED WATER-THRUSH. — This is properly *Siurus motacilla* (Vieill.), Bp., for the *Turdus motacilla*, accurately described and recognizably figured by Vieillot in 1807, is unques-

---

reason, common sense certainly tells us to spell correctly if we can. If we are always to preserve the original forms of names, we must, for example, say *Scopolax* instead of *Scolopax* — it so stands in LIXN. Syst. Nat. i, 1766, p. 242.

tionably this species. Vieillot knew the other species, which he figured and described in the same work. Bonaparte called it *Seiurus motacilla* in 1850, though in 1824 he had called the other species *Turdus motacilla*, and Cabanis, in 1857, *Henicocichla motacilla*; but writers have usually adopted Audubon's term *ludovicianus*, proposed in 1832, notwithstanding that this author soon abandoned his species, under the wrong impression it was not different from *S. œvius*. "*Siurus motacilla*" is not a very choice name, meaning "wagtailed Wagtail," but it is no worse than *Mus musculus*, *Xanthocephalus icterocephalus*, *Columba palumbus*, *Regulus satrapa*, and a host of other names, the two terms of which mean the same thing; nor as bad as *Sialia sialis*, *Cupidonia cupido*, the sense and sound of which agree.

I append the synonymy of the species of this genus, the list of names here to be given being much more accurate, more extensive, and more nearly complete than any hitherto collated:

### 1. *Siurus auricapillus*.

*Motacilla aurocapilla*, LINN., Syst. Nat. i, 12th ed. 1766, 334, No. 29 (based on Brisson and Edwards, as below cited).

*Turdus aurocapillus*, LATH., Ind. Orn. i, 1790, 328, No. 6.

*Sylvia aurocapilla*, BONAP., Journ. Philada. Acad. iv, 1824, 35.

*Seiurus aurocapillus*, SWAINS., Philos. Mag. i, 1827, 369; Zool. Journ. iii. 1827, 171.

*Seiurus aurocapillus*, D'ORBIG., Ois. Cuba, 1839, 55.

*Siurus aurocapillus*, SCL. & SALV., Ibis, i, 1859, 9. — A. & E. NEWT., *ibid.* 142.

*Enicocichla aurocapilla*, "GRAY." (Reference not at hand as I write.)

*Enicocichla aurocapillus*, BREWER, Proc. Bost. Soc. N. H. vii, 1860, 306.

*Henicocichla aurocapillus*, CABAN., Mus. Hein. i, 1850, 16.

*Turdus auricapillus*, LICHT., "Preis-Verz. Mex. Vög. 1830, 2"; Journ. f. Orn. 1863, 57. (Orig. ref. not verified by me.)

*Accentor auricapillus*, RICH., Rep. Brit. Assoc. for 1836, 1837, 172.

*Seiurus auricapillus*, BONAP., Consp. Av. 1850, 306.

*Henicocichla auricapilla*, SCLAT., Proc. Zool. Soc. 1856, 293.

*Siurus auricapillus*, COUES, Birds Colorado Valley, 187-, (*MSS. ined.*).

*Turdus citreus*, ??? MÜLLER, Syst. Nat. Suppl. 1776, 141 (very problematical).

*Motacilla canadensis*, BODDAERT, Tabl. Pl. Enl. 1783, 24 (*in part*; the first ref., to P. E. 398, f. 2, and the ref. to Edw. Gl. 252, are to this sp., but the other refs. are to *Dendroica coronata*).

*Turdus minimus*, BARTR., Trav. Fla., 1st Am. ed. 1791, 290bis (not of Lafr., nor of authors).

- Turdus coronatus*, VIEILL., Ois. Am. Sept. ii, 1807, 8, pl. 64.  
*Anthus coronatus*, GERHARDT, Naumannia, iii, 1853, 38.  
*Ficedula pensilvanica aurocapilla*, BRISS., Orn. iii, 1760, 504, No. 57.  
*Figuier à teste d'or de pensilvanie*, BRISS., *op. loc. cit.*  
*Golden-crowned Thrush*, EDW., "Glean. 91, pl. 252." (Not verified by me.)  
*Grivelette de S. Domingue*, BUFF., "Hist. Nat. Ois. iii, 317." (Not verified by me.)  
*Petite Grive de St. Domingue*, of Planche Enlum. 398, f. 2 (see the citation of Boddaert, above).  
*Grive couronnée*, VIEILLOT, *op. loc. cit.*  
*Land Kick-up*, GOSSE, B. Jam. 1847, 152.  
*Golden-crowned Accentor*, *Golden-crowned Wagtail*, *Orange-crowned Accentor*, *Oven-bird*, of AUTHORS.

## 2. *Siurus nævius*.

- Motacilla nævia*, BODD., Tabl. 1783, 47; based on Pl. Enlum. 752, f. 1.  
*Siurus nævius*, COUES, Birds Colorado Valley, 187-, (MSS. *ined.*).  
*Motacilla noveboracensis*, GMEL., Syst. Nat. 13th ed. 1788, 958, No. 69  
 (based primarily on P. E. 752, f. 1 = *nævia* Bodd.).  
*Sylvia noveboracensis*, LATH., Ind. Orn. ii, 1790, 518, No. 33.  
*Turdus noveboracensis*, . . . . . ? — PEABODY, Rep. Orn. Mass. 1839, 306.  
*Turdus (Seiurus) noveboracensis*, NUTT., Man. Orn. orig. ed. i, 1832, 353.  
*Seiurus noveboracensis*, BONAP., Comp. and Geog. List. 1838, 21.  
*Siurus noveboracensis*, SCL. & SALV., Ibis, i, 1859, 10. — A. & E. NEWT.,  
 ibid. 142.  
*Seiurus noveboracensis*, HENSHAW, App. LL. Ann. Rep. Chf. Engr. U. S.  
 Army, for 1875, p. — (p. 59 of sep. paged pamph. List B. Arizona).  
*Enicocichla noveboracensis*, "GRAY." (Ref. not at hand; probably Gen.  
 of B.)  
*Henicocichla noveboracensis*, CABAN., "Schomb. Guiana, iii, 1848, 66";  
 Mus. Hein. i, 1850, 16.  
*Motacilla noveboracensis*, TURTON, Syst. Nat., English mal-version, i, 1806,  
 589.  
*Turdus (Seiurus) noveboracensis*, NUTT., Man. 2d ed. i, 1840, 402 (in part;  
 includes another species).  
*Seiurus noveboracensis*, . . . . . ? — PRATTEN, Trans. Illinois Agric. Soc. i,  
 1855, 601.  
*Seiurus noveboracensis*, PUTNAM, Proc. Essex Inst. i, 1856, 209.  
*Motacilla tigrina* var.  $\beta$ , GM., Syst. Nat. 13th ed. i, 1788, 985, No. 153  $\beta$   
 (= Briss. iii, 513, No. 62, pl. 28, f. 5).  
*Motacilla tigrina*, 2, TURTON, *op. loc. cit.*  
*Sylvia tigrina* var.  $\beta$ , LATH., Ind. Orn. ii, 1790, 537, No. 110  $\beta$  (= Gm.  
 No. 153  $\beta$ ).  
*Motacilla fluviatilis*, BARTR., Trav. Fla. 1st Am. ed. 1791, 291.  
*Turdus aquaticus*, WILS., Am. Orn. iii, 1811, 66, pl. 23, f. 5.

- Sciurus aquaticus*, SW. and RICH., Fn. Bor. Am. ii, 1831, 229, pl. 43.  
*Turdus aquatius*, BONAP., Journ. Phila. Acad. iv, 1824, 34.  
*Sylvia anthoides*, VIEILL., "Nouv. Dict. d'Hist. Nat. 1817, 208." (Not verified by me.)  
*Turdus motacilla*, BONAP., Journ. Phila. Acad. iv, 1824, 35 (not of Vieill.).  
*Sciurus tenuirostris*, SWAINS., Philos. Mag. i, 1827, 369.  
*Sciurus tenuirostris*, GAMB., Proc. Phila. Acad. i, 1843, 261.  
*Sciurus sulfurascens*, D'ORB., Ois. Cuba, 1839, 57, pl. 6.  
*Enicocichla sulphurascens*, "GRAY."  
*Sciurus sulphurascens*, BONAP., Consp. Av. i, 1850, 306.  
*Henicocichla sulphurascens*, GUNDL., Journ. für Orn. 1855, 471.  
*Anthus l'herminieri*, LESSON, "Rev. Zoologique 1839, 101." (Not verified by me.)  
*Sciurus gossii*, BONAP., Consp. Av. i, 1850, 306.  
*Fauvette tacheté de la Louisiane*, BUFF., "Hist. Nat. Ois. v, 161" ; Pl. Enlum. No. 752, f. 1 (is the basis of Bodd.'s and Gm.'s names).  
*Figuiier brun de S. Domingue*, BRISS., Orn. iii, 1760, 513, No. 62, pl. 28, f. 5 (obviously this sp. ; sole basis of *Mot. tigrina* var.  $\beta$ , Gm.).  
*Ficedula dominicensis fusca*, BRISS., *op. loc. cit.*  
*New York Warbler*, LATH., Syn. ii, pt. ii, 1783, 436, No. 29 (= *Mot. noveboracensis* Gm.).  
*Spotted Yellow Warbler*, var. A, LATH., Syn. ii, pt. ii, 1783, 483, var. A (= *Sylvia tigrina* var.  $\beta$ , Lath.).  
*Fauvette brune*, V., O. A. S., l. s. c.  
*Bessy Kick-up, River Pink*, GOSSE, B. Jam. 1847, 151 (basis of *S. gossii* Bp.).  
*Grive de rouisseaux, ou Hochequeue*, LE MOINE, Ois. Canad. 1861, 173.  
*Water Thrush, New York Water Thrush, Aquatic Wagtail, Aquatic Wood-Wagtail, Aquatic Accentor*, of AUTHORS.

### 3. *Siurus motacilla*.

- Turdus motacilla*, VIEILL., Ois. Am. Sept. ii, 1807, 9, pl. 65 (not of Bp., 1824).  
*Sciurus motacilla*, BONAP., Consp. Av. i, 1850, 306.  
*Henicocichla motacilla*, CAB., Journ. für Orn. 1857, 240.  
*Siurus motacilla*, COUES, Birds Colorado Valley, 187-, (*MSS. ined.*).  
*Turdus ludovicianus*, AUD., Orn. Biog. i, 1832, 99, pl. 19 (afterward merged in *S. noveboracensis*).  
*Sciurus ludovicianus*, BONAP., Comp. Geogr. List. 1838, 21.  
*Siurus ludovicianus*, SCLAT., P. Z. S. 1859, 363.  
*Sciurus ludovicianus*, TRIPPE, Proc. Bost. Soc. xv, 1873, 234.  
*Henicocichla ludoviciana*, SCLAT., Cat. Am. B. 1860, 25.  
*Henicocichla major*, CABAN., Mus. Hein. i, 1850, 16.  
*Enicocichla major*, BREWER, Proc. Bost. Soc. Nat. Hist. vii, 1860, 306.  
*Grive hochequeue*, VIEILL., l. s. c.  
*Louisiana or Large-billed Water Thrush*, AUTHORS.

NOTES ON THE BREEDING OF THE BLACK TERN (*HYDRO-  
CHELIDON LARIFORMIS*) IN MINNESOTA.

BY T. S. ROBERTS.

THE Black Tern is the most abundant representative of its family in this State, making its appearance in the vicinity of Minneapolis about the middle of May. Stragglers remain until the first week in September, but the majority leave during the latter part of August. For a short time after their arrival they are to be seen flying leisurely around the larger lakes; but as the nesting-season approaches they select some prairie slough or marshy lake, and there spend the greater part of their time until the young are able to fly. Late in May or early in June the nest is built and the eggs are laid, or the eggs are deposited without any nest, as the case may be. Dr. Coues mentions (*Birds of the Northwest*, 1874) meeting with a colony breeding along the Red River, and states that there were no nests whatever, the eggs being placed on beds of decaying reeds. Such is their habit under some circumstances, but only two instances of the kind have come under my notice as yet. Once, I found three eggs laid directly on the mud on an abandoned, broken-down muskrat house in the midst of a large slough. The same day I found another set of two eggs on a bed formed by the bending over of the tops of some tall dead grass. They were thus raised more than a foot above the water, which was of considerable depth. There was no indication of a nest, the eggs being held in place by resting among the coarse grass. A very interesting and valuable note on this subject occurs in a short article by Dr. P. L. Hatch, published in the *Bulletin of the Minnesota Academy of Natural Sciences* for 1876. It is an extract from a letter written by Mr. E. W. Nelson of Chicago, and although the observations were not made in this State, I will introduce them here: "I have seen the eggs of *Sterna plumbea* deposited on masses of floating weeds in several instances, but only for the *third* brood, the bird having previously built two nests and deposited the eggs in both, which had been removed by myself to ascertain how many they would lay. The result was almost invariably as follows: first nest, three eggs; second nest, two eggs; and the third, one egg. In

several instances I found the nests floating in two and a half to three feet of water without the least sign of floating rushes in the vicinity ; in fact, there were no rushes or anything else except fine swamp grass growing anywhere near, and of this the nests were built."

As already stated, they build in this section (vicinity of Minneapolis) in the latter part of May or early in June, usually placing the nest in a prairie slough or marsh bordering an open pond. The material used in the construction is short bits of grass and reeds disposed in such a manner that a neat, but loose structure is formed. Occasionally greater skill is displayed, longer material being used, which is slightly interwoven, so that the nest may even be removed alone without injuring it. These frail structures are sometimes found upon floating masses of decayed débris, and when so situated it is necessary, with but few exceptions, to detach a portion of this underlying bed in order to remove the nest intact. But they are oftener placed upon the tops of small mounds of partially decayed vegetable matter. These mounds, undoubtedly made by the Terns as foundations for their nests, are seven or eight inches in diameter, and rise one or two inches above the surface of the water. They are placed over beds of live moss, and are partly supported by the water and partly by the moss below. It takes but a slight motion of the water to rock them, and they would undoubtedly often go adrift were they not generally protected by the grass growing around them. To obtain the nest in good condition the hand may be inserted beneath the pile and the whole lifted up.

The average external diameter of the nest of this Tern is about five inches ; internal diameter, three inches ; while the depth varies from a slight depression to three fourths of an inch or more. The eggs are either two or three in number, perhaps oftener three than two. Their ground-color varies from deep brown to greenish white. The markings consist of blotches, dots, etc., of various shades of brown. On some specimens there are a few, and on others numerous, obscure pale spots in the shell. Frequently the markings are nearly equally distributed over the entire surface of the egg, but usually are aggregated to form a wreath around the larger end. So far as my observations have extended, all the eggs taken from one nest have about the same ground-color and character of marking. The average measurement of fifteen eggs before me is 1.35 inches in length by .98 inches in width.

During the day the parent birds sit on the nest very little, leaving the incubation of the eggs greatly to the heat from the sun and the warmth arising from the damp decaying vegetable matter upon which they rest, for the nests are almost always moist inside. When the site where a colony is breeding is approached nearer than the parent birds deem safe, they make a great clamor, and dart repeatedly at the head of the intruder, occasionally venturing within a foot or two. If the nest of a pair be removed, and the birds left to themselves, they show considerable distress at their loss. Hovering over the spot from which the nest has been taken, they utter incessant cries and frequently alight to look in vain for their lost treasure. All the Terns in the neighborhood join in the cries of the bereaved pair, and the lamentation becomes general.

I once had the fortune to meet with a young Tern of this species which had evidently entered this world but a few hours before. It was a curious-looking little creature, and could swim very well. The following description may convey some idea of its appearance: body covered with a soft, fluffy down; beneath, pale sooty; above, obscure yellow, washed with grayish, and tinged with rufous on the posterior parts of the body. Scattered over the upper parts were irregular spots of black. The under surfaces of the wings, lores, and feathers next the base of the upper mandible were white. The bill was black, with a white spot at the end of the upper mandible. Legs very dark flesh-color, with a reddish tinge.

I am of the opinion that these miniature Terns leave the nest very soon after emerging from the egg. The one just described was found swimming about several feet from the nest, while just at the time one of his brothers was working his way into the world by neatly cutting the shell into halves with the point of his bill.

As soon as the young Terns are able to fly they are conducted to some suitable situation around a pond or lake, where they can sit while the parent birds supply them with food. I once counted thirty-seven sitting thus at one time on four or five panels of fence, which extended from the shore a short distance into a lake.



## TWO UNDESCRIBED NESTS OF CALIFORNIA BIRDS.

BY WILLIAM BREWSTER.

THE following nests, previously unknown to science, were collected for me by Mr. Charles A. Allen, of Nicasio, Marin County, California : —

1. CALIFORNIAN PURPLE FINCH (*Carpodacus purpureus* var. *californicus*, Baird). Two nests of this variety of the Purple Finch taken at Nicasio, Marin County, California, are before me. The first, with a set of five eggs, was collected May 10, 1876. It is a somewhat smaller structure than the nest of the Eastern bird, and is much more closely compacted. It measures externally 5.75 inches in diameter, by 2.75 in depth. Internally 2.00 inches in diameter by 1.50 in depth. The outer framework is composed of rather fine weed-stalks and coarse grasses firmly interwoven, while the inner nest is fitted smoothly and warmly with a peculiar fibrous hemp-like material of a rich bay color. This nest was found in a garden in Nicasio. It was placed in the fork of two limbs at the height of about eight feet above the ground. The eggs differ very materially from those of *Carpodacus purpureus* var. *purpureus*, and much more closely resemble eggs of the House Finch (*C. frontalis*). Their ground-color is *white* with a scarcely perceptible shade of bluish, about as much, in fact, as obtains in average eggs of the Indigo Bird (*Cyanospiza cyanea*). A very few lines and dots of black or dark brown about the larger ends constitute the only markings. They are in shape a blunted oval, and measure .73 of an inch in length by .55 in breadth. The other nest contained young, and as it was not secured until after they had left it, is in rather poor condition. It, however, agrees very closely with the one just described, and is lined with the same peculiar material. The parent bird — a male — sent with these nests is quite typical of the variety which it represents.

2. RUFOUS-CROWNED SPARROW (*Peucaea ruficeps*, Baird). — A nest of this species collected by Mr. Allen on Black Mountain, near Nicasio, July 10, 1875, presents the following features : It is outwardly composed of coarse grass and weed-stalks, and lined somewhat scantily with horse-hair. It is very loosely put together, and the original shape is so nearly destroyed that measurements are almost impracticable. An approximation would, however, be nearly as follows : External diameter, 4 inches ; internal, 2.25 inches. External depth, 2 inches ; internal, 1.25 inches. It contained three pure white eggs, which measure .89 of an inch in length by .65 in breadth. The locality was an open heathy tract on the mountain-side, and the nest was placed on the ground under a bush. Mr.

Allen, having only his rifle with him at the time, was unable to secure the female, but as she sat closely and was distinctly seen by him, there seems little reason to doubt the correctness of the identification, especially as in position of nest, color and size of eggs, etc., we find nothing incompatible with the corresponding breeding characteristics of the other and better-known species of this genus.

Mr. Allen has since informed me, by letter, that a nest satisfactorily determined as belonging to this species, and which agrees closely with the one just described, was discovered by Captain Charles Bendire in Oregon (1).

---

## A CONTRIBUTION TO THE BIOGRAPHY OF WILSON'S PHALAROPE.

BY E. W. NELSON.

ALTHOUGH this species (*Steganopus wilsoni*, Coues) is more or less common in portions of the country frequently visited by Ornithologists, it is remarkable that its life-history should be so little known. The account of nearly every author who has mentioned the species contains more or less error, and none give anything like a complete history of it. To remedy this to some extent is the object of the present paper, since I have had abundant opportunity for observing the bird in the field.

But first I wish to make a few quotations from and remarks upon the principal accounts of the species. Ord, in his edition of "Wilson's Ornithology (Vol. III, p. 205), states as follows: "Our figure of this species [*Phalaropus lobatus*, Ord] bears all the marks of haste; it is inaccurately drawn, and imperfectly colored; notwithstanding, by a diligent study of it, I have been enabled to ascertain that it is the *Coot-footed Tringa* [Phalarope] of Edwards, pls. 46 and 143, to which bird Linnæus gave the specific denomination of *lobatus*." Thus far Ord is undoubtedly correct, as is evident by a comparison of the plates in question. As Dr. Coues has already stated (Birds of the Northwest, p. 467), *Tringa lobata*, Linn. is *Lobipes hyperboreus*, (L.) Cuv., and I perfectly agree with Ord in referring Wilson's plate to the same species; but farther on Ord describes an undoubted specimen of *Steganopus wilsoni*, taken near Philadelphia, as being identical with Wilson's plate of *lobatus*, which is certainly a bad case of mal-identification. From references I have been enabled to make, I

think it extremely doubtful that Wilson ever saw a specimen of *S. wilsoni*.

Audubon's account of the sexes of this bird is quite erroneous. Concerning a pair taken near Great Egg Harbor, in June, 1829, he states that, "on examining the birds when we returned, I saw that the female had been sitting";\* and on the opposite page, "I observed scarcely any difference in the coloring of the sexes, the female being merely larger than the male"; and he again states: "The female, which is somewhat larger, is in color precisely similar to the male." The few specimens seen by Audubon during the breeding-season were apparently all females, and, taking it for granted that the males were equally bright, he so stated. In his plate of this species he figures a "female" young of the year and an adult "male," which is, in reality, a female in breeding plumage. Audubon's statement regarding the likeness of the sexes in the breeding plumage has been accepted as true by subsequent authors, even when they have had the opportunity to settle the matter for themselves in the field.

Nuttall adds considerable to the known range of the species, but makes his statements curiously conflicting, as the following quotations show: "Taking the interior of the continent for its abode, it is seen not uncommon on the borders of lakes, in the vicinity of the City of Mexico. In these situations, choosing the shelter of some grassy tuft, it forms an artless nest, in which it deposits two or three pyriform eggs, between yellowish-gray and cream-color, interspersed with small roundish spots and a few larger blotches of umber-brown somewhat crowded towards the obtuse end." He also states that "it is unknown in summer beyond the 55th parallel, passing the period of reproduction on the plains of the Saskatchewan, being also a stranger to the coasts of Hudson's Bay"; and again, that "*in the United States it can only be considered as a straggler.*" †

Dr. Coues, in his "Birds of the Northwest," arranges the synonymy of the species in a very satisfactory manner, but makes essentially the same statement as Audubon regarding the sexual plumages, and adds nothing of importance to the life-history of the species. To Mr. A. L. Kumlien‡ is due the credit of being the

---

\* Birds of Amer., Vol. V, pp. 229, 230, pl. 341.

† Man. Orn., Vol. II, pp. 245, 246.

‡ Field and Forest, July, 1876.

first to announce the true relations of the sexes of this species. His statements that "the male attends to the duties of incubation almost entirely alone," and that "not only is the female much more brilliant in plumage, but also considerably larger," are certainly true, but that the females "pursue" the males during the pairing-season seems to me to be rather doubtful, unless, as might be the case, Mr. Kumlien has mistaken for this their habit of flying restlessly about the marsh in small parties of three or four individuals, when the males are usually in advance. At these times the nearest approach I have observed to pursuit is in a habit they have of suddenly darting off for a short distance at right angles to their general course, but this appears to be in mere sport, for nearly the same relative positions are kept by the birds, and this erratic course is rarely pursued beyond a few rods.

In fact, throughout the pairing-season I have always found the Phalaropes very undemonstrative toward each other, the choice of mates being conducted in a quiet, unobtrusive way, quite unlike the usual manner among birds. Neither have I ever seen the males "drop as if shot, within two feet of me, and feign the most distressing pains," when the nest is discovered; nor even when the newly hatched young have been captured do they evince any such emotion, and at no time have I ever seen any more anxiety shown by the male than by the female. Mr. Kumlien describes the nest as being built in a tussock of grass, "much in the same manner as the *Agelaius phoeniceus*," which is certainly a considerable variation from the situations chosen by the birds in Northern Illinois, as a comparison of the above statement with my description of the situation of the nest will show.

My experience with the species has been to prove that during the breeding-season, at least, they are averse to any large body of water, and I have never found the young away from the midst of the grassy marshes until fully fledged. The last author before quoted, however, states that "the young are conducted to the shore soon after they are hatched, and if suddenly surprised take to the water and swim and dive with the greatest ease."

In Northern Illinois, where the following observations were made, Wilson's Phalarope is the most common summer resident, occurring about grassy marshes and low prairies, and is not exceeded in numbers by even the ever-present Spotted Sandpiper. As is the case with several other species of birds, Lake Michigan appears to form

a limit to its common occurrence in the eastern portion of its range. On the west it extends to the Rocky Mountains, and between these limits it has been recorded during the breeding-season from the Saskatchewan to the Arkansas (Coues) and to the city of Mexico (Nuttall). It is more closely confined to its favorite haunts than most water-birds, and this may, in a measure, account for the little hitherto known regarding its habits. During the first two weeks of May, the exact date varying with the season, this beautiful bird first makes its appearance in Northeastern Illinois. Its arrival is heralded by a few females, which arrive first, and are found singly about the marshes. At this time the females have a peculiar harsh note, which I have heard but a few times, and only from solitary individuals before the arrival of the main body.

A few days later small flocks, embracing both sexes, may be found along the borders of grassy pools, or lying at midday on the sunny side of some warm knoll in the marsh. As the breeding-season approaches they become more restless, flying from place to place, and finally separate into small parties of two or three pairs. About the middle of May their love-making commences, and is at first indicated by the increasing solicitude they show for each other's welfare. The appearance of a person in their vicinity at this time is the signal for all the birds near to come circling about, though generally not within easy gunshot. By a careful approach one may now and then find a small party swimming about in some secluded pool. The charming grace of movement exhibited at such times, combined with their tasteful elegance of attire, form one of the most pleasing sights one could witness, as they swim buoyantly from side to side of the pool, gracefully nodding their heads; now pausing an instant to arrange a feather, or to daintily gather some fragment of food, and now floating idly about, wafted by the slight breeze which at intervals ripples the surface of the water. A more common, but scarcely less pleasing sight is presented when, unconscious of observation, they walk sedately along the border of the water, never departing from their usual easy grace of movement. Their food is generally found in such places, where the receding water furnishes a bountiful supply. The only demonstrations I have observed during the pairing-time consist of a kind of solemn bowing of the head and body; but sometimes, with the head lowered and thrust forward, they will run back and forth in front of the object of their regard; or again a pair may often be seen to

salute each other by alternately bowing or lowering their heads ; but their courtship is characterized by a lack of the rivalry and vehemence usually exhibited by birds. A male is often accompanied by two females at first, but as soon as his choice is made the rejected bird joins her fortunes with some more impressible swain.

The nesting-site is usually in some thin tuft of grass on a level spot, but often in an open place concealed by only a few straggling blades of small *carices*. The male scratches a shallow depression in the soft earth, which is usually lined with a thin layer of fragments of old grass blades, upon which the eggs, numbering from three to four, are deposited about the last of May or first of June. Owing to the low situations in which the nests are placed, the first set of eggs is often destroyed by a heavy fall of rain, causing the water to rise so as to submerge the nest. In this case the second set, numbering two or three, are often deposited in a depression scratched in the ground, as at first, but with no sign of any lining. Accidents of this kind cause the second set of eggs to be sometimes deposited as late as the last of June.

The young usually appear about the third week of June, and are able to fly in about three weeks. Generally a number of pairs nest upon the same marsh. In some instances as many as fifty may be counted within the radius of a mile ; but, notwithstanding this, their nests are extremely difficult to discover, the material and the color of the eggs correspond so closely to the appearance of the surrounding surface. If they are disturbed while building, the nest is usually abandoned. Incubation is attended to by the male alone.\* The female, however, keeps near, and is quick to give the alarm upon the approach of danger. The females are frequently found at this time in small parties of six or eight ; and should their breeding-ground be approached, exhibit great anxiety, coming from every part of the marsh to meet the intruder, and, hovering over his head, utter a weak nasal note, which can be heard to

---

\* [As above stated by Mr. Nelson, Mr. Kumlien was the first to call attention to this fact, as regards the present species, as well as to the fact of the female being larger and brighter-colored than the male. European authors have recorded the same sexual peculiarities of plumage in the Red and the Northern Phalaropes (*Phalaropus fulicarius* and *Lobipes hyperboreus*), and also, in respect to the former, that the male alone undertakes the duties of incubation. In these species the male is said to show much greater devotion to the young, when exposed to danger, than does the female. — J. A. ALLEN.]

only a short distance. This note, which is possessed by both sexes, is nearly always made while the birds are in the air, and its production requires apparently considerable effort; the head and neck being inclined downward, and then suddenly raised as the note is uttered, the flight being at the same time momentarily checked. The movements of the birds usually render it an easy matter to decide whether or not they have nests in the immediate vicinity. After the first alarm, those having nests at a distance disperse, while the others take their course in the form of an ellipse, sometimes several hundred yards in length, with the object of their suspicion in the centre; and, with long strokes of their wings, much like the flight of a Killdeer, they move back and forth. As their nests are approached the length of their flight is gradually lessened, until at last they are joined by the males, when the whole party hover low over the intruder's head, uttering their peculiar note of alarm. At this time they have an ingenious mode of misleading the novice, by flying off to a short distance and hovering anxiously over a particular spot in the marsh, as though there were concealed the objects of their solicitation. Should they be followed, however, and a search be there made, the manœuvre is repeated in another place still farther from the real location of the nest. But should this ruse prove unavailing, they return and seem to become fairly desperate, flying about one's head almost within reach, manifesting great distress. If possible, still greater agitation is shown when they have unfledged young, — they even betraying their charge into the hands of the enemy by their too obvious solicitude, they then hovering directly over the young, and uttering their notes of distress. The young have a fine, wiry peep, inaudible beyond a few feet. They are very pretty little creatures, covered with yellowish-buff-colored down, with black spots on the upper surface of the body. Even when first hatched they are quite lively and difficult to capture.

About the middle of July the females suddenly disappear, and a little later the males and the young also leave, with the exception of a few stragglers, which occasionally remain until the last of August. The main portion rarely remain as late as the 10th, and are usually gone by the 5th. The males commence their fall moult before they leave; but I have never taken a specimen in which the winter plumage was very evident.

## A DEFENCE OF HIS CATALOGUE OF THE BIRDS OF NEW ENGLAND.

BY T. M. BREWER.

MESSRS. EDITORS:—There were two objects set prominently in view in my list, and distinctly stated. One was to furnish a list that shall be reliable *so far as it goes*. The other was to present a separate list of those birds attributed to New England, but in regard to which, up to May, 1875, I could \* “find no evidence that would warrant me in retaining them.” These statements seem sufficiently intelligible. The one suggests the incompleteness of the list and my expectation of additional facts. The other explains the challenged list as one given, after many years of careful investigations, as my own conclusions, for which I alone am responsible. It is my indisputable right, having made my own investigations, to form and to express my own conclusions.

In confining myself to what is reliable I necessarily had to omit all generalizations where the data were open to conflicting constructions. Thus in referring to seven species I confined myself to the single prominent feature in their New England life, their residence here in summer. The record shows (North American Birds, *passim*) that I was also well aware of their more or less limited presence in winter. To my mind their occasional presence does not necessarily prove them to be, properly speaking, *resident*, a term only applicable to cases where the same individuals are both generally and constantly present. It should not be applied, except with careful qualifications, to species where this presence is limited to a small proportion, or where it may be altogether doubtful.

---

\* My friend Mr. Deane, in recording the capture of *Sterna fuliginosa* near Lawrence, Mass., speaks of my having for some unknown reason *withdrawn* this species from the New England list and of its being now *replaced*. I object to this phraseology as calculated to give an erroneous impression. If the bird had been rightfully in the list, it was not in my power to *withdraw* it. If there is no evidence in favor of this right, it cannot be *replaced*. It was first mentioned by Mr. Samuels as breeding in Muskegat. Every one familiar with that island knows that there is not even a probability that it has ever done so. The whole statement was obviously incorrect. So well satisfied was Mr. Samuels himself of the incorrectness of his information that in his “Ornithology of New England” he omits this species. This Tern is now generally regarded as a cosmopolitan, intertropical species, rarely occurring north or south of the two tropical lines, and is not known to have occurred on Long Island, the coast of New Jersey, or anywhere north of the Chesapeake prior to the present record.



To my mind it is simply an absurdity to speak of a species as *resident* when not one individual of the entire species resides in any part of New England more than a fraction of the year. The word "race" is still a good English word, the meaning of which is so obvious that there is no occasion for misunderstanding it. According to Worcester it is "a series of descendants from one stock." In this sense, and in this only, our Summer and our Winter Robins are of different races, though specifically the same.

*Corvus americanus*, considered as a bird of *all* New England, is almost exclusively a summer resident. The few that winter are the exception, not the rule; are restricted to a very small part of New England; and are probably merely winter visitants from beyond our borders, and therefore not residents. What your correspondent quotes from my language in reference to *Picus villosus* had reference to all the United States, and not exclusively to New England, though in a more restricted sense it is also applicable. I cheerfully admit that in this case it would have been more correct, on my part, to have given it qualified as partially a resident.

It is not safe to assume, because a limited number of individuals of the other four species named are occasionally taken here in the winter, that they are necessarily resident. Without attempting to generalize, on data to my mind insufficient, I confined myself to that feature in their New England life in regard to which there would not be two opinions, leaving in abeyance all that admits of controversy. These birds are probably only winter visitants, and in no proper sense resident, or only very exceptionally resident.

Your correspondent takes up nearly a third of his second article with various *opinions* as to the occurrence in New England of the five species that formed the subject of his interrogation in his first article. But when I ask for bread he gives me only a stone. I ask for *facts*, and he gives me only *opinions*. He does not cite a single reference that I had not already fully considered. In one instance, while he goes back several years to cite opinions then expressed, but long since given up, he omits to quote the views now entertained by the same party, and in entire variance with what he does cite. In reference to *Quiscalus major*, he quotes Dr. Coues's opinion given in 1868. Twice since then Dr. Coues has publicly given his opinion *against* the occurrence of this species north of the Carolinas; first, in his admirable biography (*Ibis*, IV. 1870) of this bird, where he speaks of it as "*restricted to a narrow belt along the coast of the ocean and gulf from North Carolina to Mexico, and as rarely ever occurring north of the Carolinas*"; secondly, in a work with which, judging from his quotations, your correspondent seems to be sufficiently familiar (*Birds of the Northwest*, p. 204), where he speaks of it as "*not authentic in New England*." Why rake up an opinion given nine years ago and long since disclaimed? Why omit his real opinion now? Dr. Linsley was a correspondent of mine, and from his own account of this species I was satis-

fied that his opinions as to its occurrence were based wholly on hearsay and unreliable testimony, and subsequent claims, when tested, were invariably of the same vague, inconsistent, and contradictory character. Positive proof, such as the preservation of the alleged species or reliable witnesses, was never forthcoming. The occurrence of this peculiarly semi-tropical and local species in New England, when totally unknown north of the Chesapeake, was in itself so improbable that in the absence of any proof I could only discredit such claims. In this opinion I am fully sustained by your correspondent's strongest witness, Dr. Coues.

The same is eminently true of *Corvus ossifragus*. Your correspondent can cite only opinions. Even Mr. Brewster's record of its occurrence, though he is an expert as little likely to be mistaken as any one, does not even now, to my view, bring this species into the list of those whose occurrence with us has been indisputably proven, though it may make its future capture more probable.

In regard to *Egialitis wilsonius*, we have the *opinion* of Mr. Linsley, which rested upon no evidence; of Dr. Coues, given inferentially and with a "perhaps"; of Dr. Wood, on Long Island (!); of Mr. Allen; and again of Dr. Coues, the latter again speaking qualifiedly ("probably"). What I have said of this species still stands uncontroverted by any *facts*, and the opinions cited are in full accordance with my own given in my list. *Nettion crecca*, as I state, "is a bird liable to occur in New England," but the only instance cited was founded in error on hasty, and, as I satisfied myself at the time, incorrect conclusions. The specimen had been taken in North Carolina and not in Massachusetts. *Sula fiber*, from Mr. Linsley's own account of the specimen, which was not preserved, proved to be an immature *Sula bassana*. Mr. Putnam wrote me that he could give me no authority for his reference.

Your correspondent is skeptical in regard to *Ægiothus canescens*, *Myiodi-ocetes minutus*, *Anser gambeli*, *Bernicla hutchinsi*, *Lagopus albus*. In regard to the last-named I feel somewhat doubtful myself. The first rests on the high authority of Mr. G. A. Boardman; the second, waiving the specimen I took myself in 1836, and which was identified by Mr. Audubon, rests on the excellent authority of a good ornithologist, Dr. Charles Pickering, confirmed by no less authority than that of Mr. Nuttall himself. *Anser gambeli*, between 1836 and 1846, was much more common than it apparently is now, but even now there is no lack of evidence of its presence, though it may have escaped your correspondent's notice. A fine specimen in immature plumage has been recently taken in Gloucester, and is now in the collection of Mr. William Jeffries of Boston. In the winter of 1836-1837, Hutchins's Goose was abundant in our market from this neighborhood, as was also the Pied Duck, the Harlequin, and others now rarely seen. Several specimens were procured by me, preserved in alcohol, and sent to London for Mr. Audubon's anatomical investigations.

In my list are six or seven species given without defining the extent of their distribution,—some of them, though found to my certain knowledge all over New England, and beyond its borders, are only found in favorable localities; others probably have a more restricted range. In regard to all these, my views as to the extent of their range are fully given elsewhere, and, as your correspondent shows, are sufficiently known to him. Yet in spite of this knowledge he did not scruple to attribute to me views which he now shows he knew I did not entertain. This is especially noticeable in the case of *Vireo noveboracensis*. Here, as it seems, he knew that it is my recorded opinion that this bird rarely, if ever, goes north of Massachusetts, yet he professes to understand me as signifying *all* New England, when I had not said so, and when I had elsewhere—unrestricted as to space—stated that I did not so believe!

And where are his *facts* demonstrating that *Helminthophaga chrysoptera* is not a *rare* bird in New England? We have again only an *opinion* that a bird must not be called rare if it regularly breeds here in numbers. The numbers must be very small in this case, and the finding of the fourth nest during ten or twelve years' search by hosts of collectors, is to be spoken of in the future tense! A bird that has only been found in a very restricted area, perhaps a thousandth part of New England, and so uncommon that only two or three of its nests have ever been taken, must not be spoken of as rare!

In the case of *Coturniculus passerinus* your correspondent is excusable for misunderstanding my real meaning, as it is somewhat blindly expressed. What I intended to convey was, that while it is chiefly confined to Southern New England, it is, as a general rule, rare throughout a very extended region into which it sparsely spreads itself. Wherever found it is a species of very irregular and unequal distribution. It wanders into Northern New England, and occurs even as far to the northeast as St. Stephen, N. B. In all this extended area the localities in which it can be said to be at all common are restricted in area and few in number. Your correspondent refers to its being exceptionally common in Nantucket. All this while he well knew that the fact of its abundance on that congenial island was well known to me. (See *North American Birds*, Vol. I, p. 554, lines 20, 21 and 22.)

More than forty years ago I ventured to publish a supplementary list of the birds of Massachusetts (*Boston Jour. Nat. Hist.*, I, 435). In this list I placed inferentially and with a ? *Polioptila carulea*. From that day to the time of the publication of my catalogue I have vainly sought for any confirmation of my supposition. Your correspondent is the first to come to my support and to confirm my conjecture, but, prior to May, 1875, there is no "record" whatever confirmatory of its claim to be counted as a bird of New England. Yet because, nearly two years after the preparation of my paper, your correspondent hears of its having been taken in

Connecticut, he speaks of "its record of occurrence having been as good as any of those just cited"; that is, a *subsequent* occurrence can establish a *prior record*!

The same indefensible claim is made in behalf of *Dendroica cærulea*. This was given by Mr. Putnam as a bird of Essex County, on the supposed authority of Mr. Brickett of Portland. Mr. Brickett, when appealed to, wrote me that he had been misunderstood, that he only referred to *D. cærulescens*. So *D. cærulea* fell to the ground, and was left with absolutely no record. Its record is now wholly *ex post facto*. The fact remains indisputable that there was no authentic record of its appearance in New England at the time I so stated.

Having exhausted the all too insufficient limits to which I am restricted, I am compelled to omit nearly all that I have written in reference to *Micropalama himantopus*. I will only state that in characterizing it as "migratory, Mass.," I should have added "N. H.," in which it has been taken twelve miles from our boundary line. Though invited to do so, your correspondent is unable to give any data to show that it is migratory along the entire New England coast. It has not been found in any part of that coast from St. Andrew's to Kittery, or from Buzzard's Bay to East River, and the sweeping statement of your correspondent still remains an entirely unsupported assumption.

Here all controversy, on my part, with your correspondent ends. Whatever reference I may hereafter make to any facts or opinions bearing upon any of our New England birds, will be without any reference to a controversy that has been forced upon me, but in which I cannot do full justice to myself without becoming an infliction upon your readers.\*

---

### Recent Literature.

BURROUGHS'S "WAKE-ROBIN."—Hurd and Houghton have reprinted Mr. John Burroughs's charming little volume "Wake-Robin," wherein the wild wood-life of the birds, from Washington to the Adirondaeks, is picturesquely sketched. Mr. Burroughs has a keen eye and a loving heart towards the birds, and it is encouraging to know that this volume of his ornithological essays finds a continued sale. The present edition differs from the original (although it is labelled "revised") only in the addition of a chapter on the Bluebird, the addition of a copious index, and in the

---

\* By some oversight, which I can neither explain nor excuse, *Dendroica blackburnie* is omitted in my catalogue. It should have been given as breeding at least as far south as Massachusetts. The latest instance was noticed by Mr. Geo. O. Welch of Lynn last summer.

insertion of some wood-cut illustrations from Baird, Brewer, and Ridgway's large work, which, with the exception of the frontispiece, by Miss Brydges, almost uniformly mar, rather than beautify the volume. The very first cut is of an inconsolable Olive-sided Flycatcher, which is written down "Hermit Thrush"! But this is the fault of the publishers, who also betray their ignorance in the bad spelling of the preface, and not of the author, who did not see the proof-sheets. It is to be hoped that Mr. Burroughs will collect his later essays on birds into a second volume, which would meet with a very hearty welcome. — E. I.

MINOT'S "BIRDS OF NEW ENGLAND."\* — It would not be generous, or even just, to criticise this work as a scientific treatise or as a mature production. We prefer to side with the youthful author, who is evidently a lover of birds, keenly alive to the delights they are capable of affording, and enthusiastic in the pursuit of his favorite study, who has in an incredibly brief period trained himself to become a really good observer, and who shows that he possesses qualities which go to make a first-rate ornithologist. In this volume he not only imparts to others the knowledge of birds he has acquired, but also endeavors to awaken the same pleasurable emotions he has experienced in the acquisition: the former design is carried out with fidelity, precision, and detail, while the freshness, *naïveté*, and no little good taste which the literary execution of the work displays will go far toward meeting the latter indication; for the color of personality — if it be the genuine thing, as it is in this case unquestionably — always lends a charm to natural-history narrative. The work, moreover, shows traces of kindly interested supervision during its preparation, and the contributions to its pages are not the least valuable of its contents. There is very little technicality, chiefly taken from Baird and another writer; the descriptions, however, are tersely original. The instructions for collecting eggs differ from those ordinarily given mainly on the score of humanity, showing what may be accomplished without destroying the parents; but we waver here, saying frankly that as between a bird's life and the identification of an egg we are merciless. Next after the biographies of the birds, which are conveniently divided into sections relating respectively to the nest and eggs, the general habits, and the song or other notes, and which embody no little information not already the property of ornithologists, — on the night-habits of some species, for instance, — the most prominent and most original features of the work are the artificial "keys," in one of which the birds themselves are analyzed somewhat after a method lately introduced, the eggs of

---

\* The Land-Birds and Game-Birds of New England, with Descriptions of the Birds, their Nests and Eggs, their Habits and Notes. By H. D. Minot. Salem, Mass.: Naturalist's Agency. Boston: Estes and Lauriat. "1877" [i. e. Dec., 1876]. 1 vol. 8vo. pp. xvi, 456, figg. xylog. 29 (on 1 pl.) + 22 (in text).

Massachusetts birds being similarly handled in the other. Those who are familiar with these "short cuts" know that it is a stand-off between convenience and fallibility; but the reviewer is the last person who should find fault with them. To appreciate Mr. Minot's work as a whole, we may say that its defects are in no way the author's fault, and that they are of the obtrusive and superficial rather than of the grave or serious kind, much easier to pass over than to dwell upon ungraciously; and that its merits entitle it to full recognition by ornithologists, while they commend it very highly to the student and amateur. The mechanical execution of the volume sustains the high reputation the Salem press deserves for good work. — E. C.

---

---

### General Notes.

WESTERN RANGE OF *CONURUS CAROLINENSIS*. — Mr. E. L. Berthoud, of Golden, Col., writes under date of December 2, 1876: "I saw the Carolina Parrot, at this place (lat.  $39^{\circ} 45'$ ; long.  $105^{\circ} 8'$ ) and at Denver, on the S. Platte, in 1860-61, and on the Little Thompson River, Col., in 1862. It was abundant in Kansas in 1865-67, since which year I have seen but few, on Smoky Hill and Republican Forks. I have also seen it near old Fort Lyon, on the Arkansas River." I am not aware that the species has hitherto been reported as occurring so far west as Colorado. — ELLIOTT COUES, *Washington, D. C.*

FECUNDITY OF THE CAROLINA WREN (*Thryothorus ludovicianus*). — About April 25 I found "our pair of Wrens" very busy, the male being followed by five nearly full-fledged young, and the female actively engaged in constructing (under the rafters of our stable) another nest, in which she soon deposited five beautiful eggs, and commenced sitting, cheered by the loud and happy notes of the male, who had by this time got rid of his noisy brood. In due time five more young Wrens made their appearance, and never did birds work harder than did their parents to supply their insatiable appetites. Spiders, bugs, and larvæ of every description, were brought in quick succession, and, as a consequence, a rapid growth was the result, and the brood was out by the fore part of July, following the male and "quivering" their wings in supplication for food. The female immediately set herself at work on another nest, this time under the eaves of a porch. Large quantities of dry leaves and coarse grass and weeds were carried up, and a compact oval structure was made, with a round cavity in the top, partly roofed over. On the 19th of July I found five eggs in the nest, and the female again sitting. Are three broods in a season commonly reared by this species? — CHARLES DURY, *Avondale, Hamilton County, Ohio.*

THE LOUISIANA HERON IN INDIANA. — My friend Mr. F. T. Jencks, of Providence, R. I., writes me that on the 26th of June, 1876, while passing through a large marsh between Plymouth and Hanna, on the line of the Pittsburg and Fort Wayne Railroad, in Northern Indiana, he saw a fine adult specimen of *Demigretta ludoviciana* spring up close beside the railroad-track and fly off in full view. As Mr. Jencks is well acquainted with the species in question, I have no doubt of the correctness of his identification. — E. W. NELSON, *Chicago, Ill.*

NOTE ON THE CINNAMON TEAL (*Querquedula cyanoptera*). — A small lake, which feeds one of the headwaters of the North Platte, in North Park, Colorado, was found to be a breeding-place of large numbers of wild Geese (*Branta canadensis*) and other water-fowl, among which Wigeons (*Mareca americana*), Shovellers (*Spatula clypeata*), and the species the name of which heads this paragraph, were the most numerous. I was on the spot too late in the season to take eggs, but newly-fledged birds of each of the three species of Ducks mentioned, as well as old birds in moult, were secured or observed. The Cinnamon Teal here seemed to replace the common Blue-winged, none of which were ascertained to occur. The spot was on the Atlantic side of the main water-shed, though practically on the divide, as it was only a day's march from the edge of Middle Park, the waters of which area flow into the Colorado of the West and so into the Gulf of California. The Teal associated as usual with various other kinds of Ducks, and no peculiarity of habits was noted. Two young birds were captured alive, in a natural excavation of an embankment, in which they had apparently crawled to hide, as the hole showed no traces of a nest. — ELLIOTT COUES, *Washington, D. C.*

ÆGIOTHUS EXILIPES IN EUROPE. — It may be interesting to the readers of the Bulletin to learn that one of the two Redpoles procured upon the Petchora River in Northeastern Russia, in 1875, by Mr. Henry Seebohm and myself, and on a former occasion, at Archangel, by Mr. E. R. Alston and myself, has turned out to be identical with *Ægithus exilipes*, Coues. We are thus able to extend the distribution of that species in European Russia as far as Archangel (40° E. longitude from Greenwich). This species was also procured by M. Severtzoff, in Turkestan. In our papers in the "Ibis," January, 1873, p. 64, and "Ibis," January, 1876, we have erroneously named this bird *Æ. rufescens*, which name is only applicable to the form found in Great Britain, and migrating southward in winter. The other species found in North Russia is true *Linota linaria*, Linn. *Ægithus exilipes*, Coues, will thus probably be found to be almost circumpolar in its distribution, as it is reasonable to suppose that it is the common species occurring throughout Northern Siberia in summer. References to notices of this species in Europe will be found in the "Ibis" as above quoted, in the "Zoölogist" for January, 1877, and in an Appendix Mr. Seebohm and myself are preparing to our "Notes of the Birds of the

Lower Petchora"; and it will be doubtless still more fully treated of in a forthcoming part of Mr. H. E. Dresser's "Birds of Europe." — J. A. HARVIE BROWN, *Cor. Mem., Dunipace House, Larbert, Scotland.*

A NOTE ON CUPIDONIA CUPIDO VAR. PALLIDICINCTUS, RIDGWAY. — In the latter part of January, 1877, I found in Fulton Market about thirty specimens of this form; they were generally unfit for preserving, but I got two in good condition. On examination they agreed accurately with Mr. Ridgway's description (*N. Am. Birds*, Vol. III, p. 446). I ascertained that they came from Pierce City, Southwestern Missouri. I have been unable to make much inquiry for others since. I lately learned from a large dealer that they had been quite abundant in market, all coming from Southern Missouri. The marketmen objected to buying them on account of their small size. I found their average weight to be one and three-fourths pounds, some weighing but one and a half pounds. All I talked with said they had not noticed them before this winter. — GEORGE N. LAWRENCE, *New York.*

CAPTURE OF THE EGYPTIAN GOOSE ON LONG ISLAND. — On the 3d of January, 1877, I received a remarkably fine specimen of a species of Goose entirely unknown to me. The bird was killed in a pond of fresh water near Carnarsie, Long Island, and has every appearance of being a wild bird. The plumage is in fine condition, and the feet are free from warts. On exhibiting it to our well-known ornithologist, Mr. G. N. Lawrence of New York, he expressed great surprise, and promised to investigate the matter. I have since received from him the following communication: —

"The Goose shown me yesterday is the Egyptian Goose (*Chenalopez aegyptiacus*, Linn.). It inhabits all of Africa, and numerous specimens have been killed in Great Britain. Its acquisition is worthy of being noted, and whether a straggler or an escaped specimen may be ascertained in the future."

The specimen will be placed in the Museum of the Long Island Historical Society of Brooklyn. — JOHN AKHURST, *Brooklyn, N. Y.*

MACCOWN'S LONGSPUR IN ILLINOIS. — While looking over a box of Snow-Buntings and Shore Larks in the market, January 15, 1877, I found a specimen of *Plectrophanes maccowni*, shot at Champaign, Illinois. January 17, another box containing Lapland Longspurs was sent from the same place, and among them was a second specimen of *P. maccowni*, which is now in the collection of C. N. Holden, Jr., Chicago. January 19 I obtained a third specimen from the same source, which has been sent to Mr. E. W. Nelson, of this city. They were all males, showing plainly the chestnut coloring on the bend of the wing and the peculiar white markings of the tail. This is, I think, the first record of the occurrence of this bird in Illinois, if not east of Kansas. — HENRY K. COALE, *Chicago, Illinois.*



# BULLETIN

OF THE

## NUTTALL ORNITHOLOGICAL CLUB.

Vol. II.

JULY, 1877.

No. 3.

### DESCRIPTION OF A NEW SPECIES OF HUMMING-BIRD FROM CALIFORNIA.

BY H. W. HENSHAW.

#### **Selasphorus alleni**, NOBIS.


SP. CHAR. — *Adult Male*. Two outer tail-feathers very narrow, linear, the outer nearly acicular; second only slightly larger; third abruptly larger. Upper parts bright golden-green, dullest on the crown. Under tail-coverts, sides, and belly rufous, paler along the median line. Jugulum white. Wings purplish-dusky. Tail-feathers cinnamon, tipped and edged with purplish-brown. Throat and ruff bright coppery-red, with brassy-green reflections. *Female* similar to the female of *S. rufus*, but smaller; tail-feathers narrower, especially the outer.

*Habitat*. Coast district of California; northward?

#### DIFFERENTIAL DIAGNOSES.

*S. RUFUS*. — *Male*. Tail-feathers successively graduated in size, the outer the smallest; fourth from the outer abruptly attenuated, and with a deep notch at the base of attenuation on the inner web. Crown (only) metallic-green. Upper parts clear cinnamon, sometimes with flakes and patches of metallic-green on the back.

*S. ALLENI*. — *Male*. Smaller. Two outer tail-feathers very narrow, linear, the outer nearly acicular;



\* These figures were kindly drawn by Mr. Ridgway, and afford a very exact idea of the differences in the form of the tails of the two birds.

third abruptly larger ; the fourth intermediate in length between the third and fifth. Entire upper parts to the tail golden-green.

SOME time since my attention was attracted by a specimen of the above-described bird, in the collection of Mr. William Brewster, which had been received from Mr. C. A. Allen, of Nicasio, California, the locality where it was taken. Experience in the interior, especially in Arizona, had afforded opportunity for the examination of hundreds of the *Selasphorus rufus*, and this specimen, an adult male, differed so decidedly from any I had ever seen that I was led to the belief that there might be on the west coast a variety peculiar to that region. The examination of specimens, however, showing what I considered intermediate phases of coloration from this same locality, led me to conclude that the form was not sufficiently distinct to receive a name, an impression which I now think was wholly erroneous. The reception of new material and a reconsideration of the question has convinced me that the form in question is quite distinct specifically from *S. rufus*, from which it is separable by perfectly good and stable characters of external structure. As might be expected in a family where the females of totally different genera are often distinguishable only with difficulty, it is in the males that the differences are chiefly to be noted. The narrow outer tail-feathers are in the case of *S. alleni* sufficiently peculiar, however, to enable one to distinguish the females and even the young birds.

The adult males are at once separable, not only by the bright green back, the green extending partially over the upper tail-coverts and contrasting sharply with the rufous of the tail, but even more readily by the very peculiar characters of the tail, as above indicated. These are perfectly constant in all the specimens I have seen, and have proved to be so also in all of the many individuals which have passed under Mr. Allen's notice.

In reference to the coloration of the *S. alleni*, I can state that the amount of variation in the adult males is very small. The back is always of a pure metallic green. Mr. Allen, whose authority is unquestionable, and who has had ample opportunity for ascertaining, writes me that the variation is extremely slight, and that a series of thirty males then before him showed no differences. On the other hand, I can assert with equal positiveness that the *S. rufus* never assumes this complete green phase of coloration. Of specimens

from the interior, I have seen at least two hundred adult males, and they all possessed the bright rufous back, only an occasional individual showing a faint trace of the green. In California, however, where the *S. rufus* occurs in its typical condition, that is, with an unmixed rufous back, specimens are not uncommonly found which exhibit a strong approach to the coloration of *S. alleni*. In these the rufous is seen to be mixed with green in varying proportions. Beyond a certain point they appear never to pass, and all such specimens have the peculiar tail characters of *S. rufus* as strongly marked as in the most typical examples of the species. We find the same difference between specimens of *S. rufus* taken in California and on the west coast generally, and those from the interior, that usually obtains in species possessed of a similar range, — that is, the coloration is appreciably darker. The locality of specimens, whether from the coast or the interior, may by this means be readily told.

In Mexico only the *S. rufus* appears to occur, and Mr. A. Boucard informed Mr. Ridgway that, having examined hundreds of the *Selasphorus* from there, he had never seen an example of the Green-backed form, which, however, he was familiar with from California. One of the strongest proofs of the specific distinctness of the two is seen in the fact that the two birds, in their typical condition, occur at the *same locality* and at the *same season*. Both breed about Nicasio (situated about twenty-five miles north of San Francisco), though Mr. Allen finds the *S. alleni* very much more abundant than the other. Thus, during the present season (1877), he has shot about forty-five of the former bird, and has only succeeded in taking four of the rufous-backed species. Last year's experience was a similar one.

This gentleman, whose investigation of these two birds has extended over several years, and who, it is but proper to state, has always believed them to be distinct, has found a constant difference in size, the *S. alleni* being the smaller. Such I find to be the case in the several specimens I have measured, and the discrepancy between them is quite considerable when the diminutive size of the birds of this family is taken into account.

*Habits.* I am in possession of but few notes bearing upon the habits of this Hummer. Mr. Allen remarks incidentally in a letter that the Green-backs are much the livelier and more active of the two, keeping constantly in the open, and always perching upon the most prominent dead twigs they can find. Their extreme shyness,

as contrasted with the unsuspecting nature of the Rufous-backed, is quite remarkable. They seem to possess a larger share than usual of the courage and pugnacity which is so constantly displayed in birds of this family. Not only do they always come off the victors when chance encounters take place between them and the Rufous-backs, but Mr. Allen has seen a pair attack and put to rout a Red-tailed Hawk; while, as he remarks, "Sparrow-Hawks have no chance at all with them." He has often seen the little fellows in hot chase after these latter birds, and their only care seemed to be to get out of the way as soon as possible of foes so determined.

The Rufous-backed Hummer, on the contrary, frequents the thickets, and is always unsuspecting and readily approached. The different localities they affect may indicate a difference in the flowers from which they obtain their food.

*Habitat.* The habitat of *Selasphorus alleni* seems to be confined to the coast district of California, though subsequent investigation may show that it extends its range to the northward. It appears to be strictly limited to the western slope of the Sierras, and may indeed wander but a short distance from the coast. It is thus very local, while the *S. rufus* has a very widely extended habitat. Few species, indeed, of the family range over as many degrees of latitude as this, which appears equally at home in the valleys and elevated table-lands of sub-tropical Mexico and in the less inviting regions of Southern Alaska, while it occupies in summer most of the intermediate ground. Specimens are in the Smithsonian collection from Sitka, which forms its most northern recorded locality. In the interior it appears to be of less general occurrence than in the region west of the Sierras, though this apparent absence is doubtless due in part to our lack of knowledge of the Avifauna, especially of the northern interior. The eastern slope of the Sierras is apparently occupied by it throughout their whole length. Mr. Ridgway found it in autumn in the East Humboldt Mountains, and Dr. Coues in Montana, and it may be said to occur over most of the Rocky Mountain region, either as a summer resident or as a fall migrant. Dr. Coues is the only one who has found it breeding in the interior, but I think it probable that not only the mountains about Fort Whipple, which formed his field of observations, but the mountain-fastnesses throughout the Territory of Arizona, and also in New Mexico, and perhaps in Southern Colorado, may furnish the species a summer home. Certain is it that in August this Hum-

ming-Bird is found almost simultaneously over all the more inviting parts of this lower country, and in certain sections where flowers abound, as in the mountains of Eastern Arizona, their numbers far exceed all other species put together.

Though apparently so hardy, and though pushing its migrations in spring in search of its summer home to a point attained by no other species of its family, it yet in winter retires far south, and appears to find the climate, even of California, too rigorous for its endurance. Possibly, however, the winter flora is not suited to its habits, and this may govern its choice of a winter residence, rather than inability to live in a climate which appears well adapted to its less fastidious rival, the Anna Humming-Bird (*Calypte annæ*). However this may be, it is certain that the great proportion, perhaps all, of the two species of *Selasphorus* leave our territory entirely in fall, and if any remain they are but a very limited number, which take refuge in the warm interior valleys of California.

Below is appended a table of measurements of the two species. The specimens of *S. rufa* are purposely selected from different localities, and the measurements give a good idea of the amount of variation:—

Sex.	Date.	Locality.	Collector.	Wing.	Tail.	Bill.
<b>S. rufus,</b> ♂ ad.	March 29, 1877	Nicasio, Cal.	C. A. Allen	1.59	1.23	.61
" ♂ ad.	March 29, 1877	" "	" "	1.62	1.33	.63
" ♂ ad.	— — 1876	San Francisco, Cal.	F. Gruber	1.67	1.28	.65
" ♂ ad.	— — 1876	" " "	" "	1.60	1.27	.65
" ♂ ad.	May 29, 1855	Columbia River,	J. K. Townsend	1.52	1.32	.63
" ♂ ad.	— — 1855	Mirador, E. Mex.	Dr. Sartorius	1.52	1.27	.66
" ♂ ad.	April 21, 1851	Stellacoom, W. T.	Dr. Suckley	1.57	1.27	.63
" ♂ ad.	August 25, 1873	Apache, Ariz.	H. W. Henshaw	1.58	1.32	.63
" ♀ ad.	July 24, 1873	El Moro, N. Mex.	" "	1.75	1.22	.72
" ♀ ad.	August 12, 1874	Ft Garland, Col.	C. E. Aiken	1.77	1.25	.68
" ♀ ad.	August 17, 1875	Tejon Mts., Cal.	H. W. Henshaw	1.66	1.17	.63
" ♀ juv.	August 18, 1875	" " "	" "	1.73	1.21	.67
<b>S. alleni,</b> ♂ ad.	March 24, 1875	Nicasio, Cal.	C. A. Allen	1.47	1.21	.64
" ♂ ad.	March 31, 1877	" "	" "	1.48	1.17	.63
" ♂ ad.	March 31, 1877	" "	" "	1.50	1.17	.67
" ♂ ad.	March 31, 1877	San Francisco, Cal.	" "	1.52	1.17	.63
" ♀ ad.	May 18, 1876	Nicasio, Cal.	" "	1.63	1.15	.68
" ♀ ad.	March 17, 1877	" "	" "	1.61	1.16	.67
<b>S. rufus,</b> Average of eight males				1.58	1.29	.64
" " " four females				1.72	1.21	.68
<b>S. alleni,</b> " " four males				1.49	1.18	.64
" " " two females				1.62	1.15	.67

The number given of *S. alleni* is less than would have been desirable, and I regret that I have not at hand the large number of

measurements taken by Mr. Allen. As he informs me, however, that the variation in this respect is very small, it is probable the average would be little changed by their addition.

In bestowing the specific name, I have paid but a deserved compliment to the zeal and enthusiasm of Mr. C. A. Allen, but for whose efforts in obtaining the specimens necessary for comparison, and careful field-notes, the species might have remained for a long time still unrecognized.

---

---

## THE BIRDS OF GUADALUPE ISLAND, DISCUSSED WITH REFERENCE TO THE PRESENT GENESIS OF SPECIES.

BY ROBERT RIDGWAY.

THE importance of insular faunæ in their bearing on the subject of the derivation of species, has been recognized by eminent writers; and certain islands, remote from the mainland, among which may be mentioned the Galapagos, have received marked attention. An excellent memoir upon the birds of the latter group has lately been published by Mr. Osbert Salvin,\* who discusses the relationship of certain peculiar forms to their continental allies, and even goes so far as to point out among the latter the present representatives of the parent stocks from which the genera now peculiar to the Galapagos have been derived. That Mr. Salvin is probably correct in his conclusions, is very strongly indicated by certain facts developed from a study of the fauna of Guadalupe, where in the case of almost every species, precisely the same local modifications are observable; the process of change in the latter case, however, has been either more recent or more gradual, since close affinity to continental representatives is clearly seen in every species, the extreme differentiation met with in the Galapagos types not having yet been reached.†

---

\* On the Avifauna of the Galapagos Archipelago. By Osbert Salvin, M. A., F. R. S., etc. Transactions of the Zoological Society of London, Vol. IX, Part IX, pp. 447-510, pls. lxxxiv-lxxxix. May, 1876.

† These modifications seem to be, (1) enlargement, or elongation, of the bill and feet, (2) abbreviation of the wings and tail, and (3) darkening of the colors

The island of Guadalupe is situated off the coast of Lower California, between latitude  $28^{\circ} 45'$  and  $29^{\circ} 10'$  north, and about two hundred and twenty miles southwest from San Diego. Until very recently nothing was known regarding the ornithology of this island, but in the spring of 1875 a collection of birds was made by Dr. Edward Palmer, and forwarded to the National Museum at Washington. This collection, embracing eight species and seventy-two specimens of land-birds, was placed in the hands of the writer for identification, and reported upon in the Bulletin of the U. S. Geological and Geographical Survey of the Territories, Vol. II, No. 2, April 1, 1876 (pp. 183-195).\*

As above stated, the land-birds contained in the collection from Guadalupe embrace only eight species, so that the fauna of the island is by no means fully represented; indeed, the collector observed a Humming-Bird, two kinds of Owls, and a Hawk, of which no specimens were obtained. This is to be regretted, since most, if not all, of these would doubtless have proved new. It is altogether likely, too, that other species escaped notice, and thus remain to be discovered; a rich field is therefore left to the future explorer. The affinities of the birds of Guadalupe are, so far as known, almost entirely with those of Western North America, there being no peculiar types, each species having a more or less closely related representative on the continent. The species thus far known are the following, their continental representatives being given opposite:—

*Guadalupe Species.**Mainland Representatives.***Sylviidæ.**1. *Regulus obscurus.**Regulus calendula.*

of the plumage. The following remarks by Mr. Salvin are those which refer more particularly to these peculiarities:—

“In the formation of its bill, it [*Geospiza*] hardly differs at all from some species of *Guiraca*, as *G. conereta* and its allies. The legs and feet, however, are much longer and stronger than in any species of *Guiraca*; and the tail in proportion to the wings is very short” (l. c., p. 478).

“From *Controstrum* [the continental representative genus] *Certhidea* [of the Galapagos] differs in having much shorter wings and tail” (l. c., p. 476). [*Italics our own.*]

\* Ornithology of Guadeloupe [*leg* Guadalupe] Island. Based on Notes and Collections made by Dr. Edward Palmer.

**Troglodytidae.**

- |                                      |                               |
|--------------------------------------|-------------------------------|
| 2. <i>Thryomanes brevicauda</i> .    | <i>Thryomanes bewicki</i> .   |
| 3. <i>Salpinctes guadalupensis</i> . | <i>Salpinctes obsoletus</i> . |

**Fringillidae.**

- |                                |                               |
|--------------------------------|-------------------------------|
| 4. <i>Carpodacus amplus</i> .  | <i>Carpodacus frontalis</i> . |
| 5. <i>Junco insularis</i> .    | <i>Junco annectens</i> .      |
| 6. <i>Pipilo consobrinus</i> . | <i>Pipilo maculatus</i> .     |

**Picidae.**

- |                                 |                             |
|---------------------------------|-----------------------------|
| 7. <i>Colaptes rufipileus</i> . | <i>Colaptes mexicanus</i> . |
|---------------------------------|-----------------------------|

**Falconidae.**

- |                               |                            |
|-------------------------------|----------------------------|
| 8. <i>Polyborus lutosus</i> . | <i>Polyborus tharus</i> .* |
|-------------------------------|----------------------------|

The more prominent characteristics of these Guadalupe birds, as compared with the mainland forms, are (1) increased size of the bill and feet, (2) shorter wings and tail, and (3) darker colors; these variations are by no means uniform, however, in the several species, the differentiation being in some slight, while in others it amounts to almost generic distinctness; but, what is very remarkable, and of great interest in connection with the subject in hand is, that no matter how great may be the modification of form and proportions, *the specific characters, so far as coloration is concerned, are in every case strictly preserved!* † Thus, *Carpodacus amplus* has the wing and tail barely longer than *C. frontalis*, but the bill is three times as large, and the feet twice as stout, *while the colors and markings are substantially identical*; *Junco insularis* has precisely the same plumage as *J. annectens* (except that the shades of color are just perceptibly darker), but the bill is so elongated, and the wings and tail so much abbreviated, that in form it is much more like the species of *Ammodromus* than those of typical *Junco*! *Thryomanes brevicauda* does not differ more than just appreciably in colors and marking from *T. bewicki*, but the difference in form is so great as to render it necessary either to institute a new genus, or, as the only alternative, to draw up a generic diagnosis entirely

\* Singularly enough, the *Polyborus* from Guadalupe resembles much more closely the South American species than that from Mexico (*P. cheriway*), though it is very distinct from either.

The only water-bird collected on the island was a specimen of *Colymbus pacificus*, in perfect breeding plumage, *found dead on the shore*.

† With the sole exception of *Polyborus lutosus*.



different from that heretofore considered to embrace the most important characters. In the case of *Carpodacus*, there has been reached in the Guadalupe race almost, if not quite, that degree of differentiation which distinguishes the Galapagoan genus *Geospiza* from continental *Guiraca*, the modifications being moreover of precisely the same character. *Junco insularis* may likewise be compared with the species of *Cactornis*, in which the bill has become so extremely produced as to have almost lost its fringilline character.

The three species above-mentioned exhibit the local modifications to the greatest extent, but the rule may be traced through the whole series; and to show the exact extent of these modifications of form we present the following table of measurements of each of the Guadalupe species compared with those of the mainland representative form, the measurements representing the *maximum and minimum of a large series* of each:—

SPECIES.	MAXIMUM.					MINIMUM.				
	Wing.	Tail.	Bill from Nostril.	Tarsus.	Middle Toe.	Wing.	Tail.	Bill from Nostril.	Tarsus.	Middle Toe.
<i>Regulus obscurus</i> * .....	2.20	1.95	.25	.80	.40	2.00	1.70	.22	.80	.38
" <i>calendula</i> .....	2.45	2.03	.22	.76	.35	2.20	1.60	.20	.65	.35
<i>Thryomanes brevicauda</i> ..	1.90	1.80	.50	.75	.50	1.85	1.80	.45	.70	.50
" <i>bewicki</i> .....	2.38	2.70	.45	.80	.50	2.00	2.00	.38	.63	.45
<i>Salpinctes guadalupensis</i>	2.75	2.30	.60	.90	.55	2.50	2.00	.55	.80	.50
" <i>obsoletus</i> .....	3.00	2.42	.50	.90	.58	2.65	2.20	.50	.73	.55
<i>Carpodacus amplus</i> .....	3.35	2.90	.45	.85	.65	3.10	2.60	.40	.75	.58
" <i>frontalis</i> .....	3.20	2.90	.35	.70	.50	2.85	2.40	.30	.62	.50
<i>Junco insularis</i> .....	2.85	2.60	.38	.85	.60	2.50	2.30	.35	.80	.55
" <i>annectens</i> .....	3.43	3.40	.30	.80	.60	3.00	2.80	.28	.80	.55
<i>Pipilo consobrinus</i> .....	3.25	2.80	.40	1.08	.75	2.90	3.25	.35	1.00	.60
" <i>megalonyx</i> .....	3.60	4.60	.40	1.05	.75	3.30	4.00	.35	1.05	.70
<i>Colaptes rufipileus</i> .....	6.25	5.30	1.60	1.10	.92	5.90	4.75	1.35	.....	.....
" <i>mexicanus</i> .....	7.00	6.00	1.25	1.20	.85	6.65	5.50	1.15	.....	.....
<i>Polyborus lutosus</i> .....	16.40	11.65	1.35	3.75	2.10	15.00	10.50	1.25	3.50	1.80
" <i>cheriway</i> .....	16.50	10.00	1.48	3.75	2.10	14.60	9.00	1.20	3.20	1.90

\* The Guadalupe species are in Italics.

A close perusal of the above figures leads to the discovery of some exceptions to the rule of variation in proportions. Thus, *Regulus obscurus*, while conforming in other respects, does not have the tail constantly shorter than *R. calendula*, although it averages shorter. There is also no appreciable difference in the absolute length of the tarsus in the two forms of *Thryomanes* and *Salpinctes*, though the comparative difference in favor of *T. brevicauda* and *S. obsoletus*, when contrasted with the length of the wings and tail, is very marked.

These exceptions we are unable to explain, but even unaccounted for they do not detract from the high importance of the variations we have noted. It will also be observed that there is no essential difference in dimensions or proportions between *Polyborus lutosus* and *P. cheriway*, the modifications being almost entirely in the plumage, which in the former species is so distinct, at all ages, from that of the other in corresponding stages that it may be regarded as one of the most completely differentiated birds of the whole series.

Not only in proportions, but also in colors, do the modifications presented by these Guadalupe birds correlate with characteristics of the Galapagoan forms. A conspicuous character of the latter is their sombre plumage of black or fuliginous-brown; now, excepting only *Polyborus lutosus*, precisely the difference in plumage of the Guadalupe birds from their Continental allies consists in their darker colors. *Carpodacus amplus*, although a bird of at least double the bulk of *C. frontalis*, is so nearly identical in plumage that positively the only difference consists in the slightly darker shade of all the colors; *Junco insularis* is darker than *J. amnecetus*, but is otherwise similar in plumage; *Regulus obscurus* is much darker than *R. calendula*; *Thryomanes brevicauda* and *Salpinctes guadalupensis* are likewise darker in colors than *T. bewicki* and *S. obsoletus*, while *Colaptes rufipileus* differs from *C. mexicanus* in having one half more black on the under side of the tail, besides being darker generally. In *Pipilo consobrinus*, however, the black portions of the plumage are hardly so intense black as in the mainland forms of *P. maculatus*, but the female is almost if not quite as black as the male,\* while in the others she is more or less conspicuously different, being some shade of brown or gray instead of black. As remarked before, the only real exception to the rule is *Polyborus lutosus*, but this has a quite different distribution of colors from the two Continental species; it may be observed, however, that while the black markings are replaced by dark brown, the lighter markings are pale clay-color instead of white; and further, that there is far less difference between the young and adult stages.

Not the least interesting fact concerning these Guadalupe birds

---

\* The similarity of the sexes in birds having a black plumage is remarkably prevalent among the West India birds, as Professor Baird has somewhere noted.

is their anomalous geographical relation to their mainland representatives, the latter being the Rocky Mountain or Middle Province races *instead of those from the intervening coast districts!* Thus, *Carpodacus amplus* agrees with *C. frontalis* in the restriction of the red in the male to sharply defined and limited areas, the coast form, *C. frontalis rhodocolpus*, having the red "spread," as it were, over the greater portion of the plumage; *Junco insularis* is a perfect repetition of *J. annectens* so far as plumage is concerned (except that the shades of color are somewhat darker), and does not at all resemble *J. oregonus* of the coast. *Thryomanes brevicauda* is colored more like *T. bewicki leucogaster* (the Upper Rio Grande form) than *T. bewicki spilurus* (the coast form); while *Salpinctes guadalupensis* differs in the same way from *S. obsoletus*, which, moreover, *is not represented at all in the coast district*, except perhaps rarely and locally in Southern and Lower California.

The peninsula above mentioned also presents in many respects closer affinities to the middle region than to the coast district, especially in the fauna found at Cape Saint Lucas; but on the western side many of the true Californian forms replace those of the Middle Province, *Carpodacus rhodocolpus* being a case in point, this species thus entering as a separating wedge between *C. frontalis* and *C. amplus!* Now very similar anomalous cases occur among Galapagos birds, an entirely parallel instance being afforded by *Dendroica aureola*, of which Mr. Salvin (l. c., p. 474) remarks: "The bird from the Galapagos [meaning the above-named form] is the same as that from Jamaica,\* whereas on the intervening continent two other (so-called) species occur — namely, *D. aestiva* as a winter migrant, and *D. vieilloti* as a resident — but never, as far as we know, *D. petechia.*" Another quite similar case is afforded by *Myiarchus magnirostris* of the Galapagos, since Mr. Salvin says that "its nearest allies are perhaps the island races of the Antilles rather than those of the continent; and in this respect the affinities of *Dendroica aureola* are, to some extent, repeated; but in the present case the specific characters of *M. magnirostris* are well defined" (l. c., p. 492).

In the paper above referred to, are incorporated notes by the collector, Dr. Habel, on the habits of the species; and in these refer-

---

\* We do not agree with Mr. Salvin in considering the forms of this species from the West Indies and the Galapagos *absolutely* identical, but recognize in them well-marked races, differing from each other about as much as the Guadalupe birds do from those of the mainland.

ence is made to the exceeding tameness of certain Galapagos birds. A similar confidence in man was likewise found to be a characteristic of some of the Guadalupe species, as an instance of which we quote the following from Dr. Palmer's notes regarding *Junco insularis* [see page 189 of our paper, cited at the head of this article]: "These are the most abundant birds of the island, and are so tame that they may be killed with a stick or captured in a butterfly-net. While I was looking for insects under stones and logs, these birds would sometimes join in the search, and hop almost into my hands. They gathered chiefly ants and their eggs. At times they even enter the houses, picking up anything edible they can find. Numbers boarded the schooner as we neared the island, and made themselves perfectly at home, roaming over every part of the vessel in search of food." It seems, however, that not all the birds of the island were thus unsuspecting and familiar, since Dr. Palmer remarks that it is difficult to secure *Thryomanes brevicauda*, on account of its shyness.

In conclusion, a few words regarding the derivation of these insular forms may not be out of place. As to those of the Galapagos, Mr. Salvin expresses the following opinion: "Considering their purely volcanic nature, it cannot reasonably be doubted that these islands have always been islands since they emerged from the sea. Such is Mr. Darwin's view; and it is fully indorsed by Dr. Hooker and others. The birds that are now found, being related to American birds, must have emigrated thence and become modified by the different circumstances with which they became surrounded. The oldest immigrants seem to be indicated by their generic difference from their continental allies, the more modern comers by their merely specific distinctness, and the most recent by their identity with birds now found on the adjoining continent. On this view the islands were first taken possession of by individuals of the parent stock of *Certhidea* and *Conirostrum*, *Geospiza* and *Guiraca*, *Camarhynchus* and *Neorhynchus*. Then came perhaps the ancestors of *Buteo* [*galapagensis*]\*; after these followed *Mimus*, *Pyrocephalus*, and *Myiarchus*, *Strix* and *Asio*, *Zenæda*, *Larus* and *Spheniscus*. Then those of *Dendroæca*, *Progne*, *Butorides*, *Nycticorax*, and *Porzana*, and, finally, *Dolichonyx oryzivorus*, *Ardea herodias*, and the Ducks, Flamingo, Gannets, Plovers, and Sandpipers, though of these last a constant stream

---

\* The nearest ally of the Galapagoan *Buteo* is *B. poliosomus* of the Patagonian region.

of immigrants may have been maintained from the earliest times. It must be remembered, however, that no precise order of immigration can be laid down, even approximately; for one term in the proposition is an absolutely unknown quantity. We know nothing of the rate of change that has taken place in any one species. Outward circumstances may have acted on one species, so as to leave it little changed in a given time, whilst in the same time another species may have assumed distinctive generic characters. Viewing the very peculiar physical characters of these islands when contrasted with the neighboring American shores, it would seem reasonable that the rate of change demanded of an immigrant species would be high; consequently the origin of the islands need not be dated back to a more distant period than seems indicated by their volcanic origin."

Considered in connection with the subject discussed above, the birds of Guadalupe are of extreme interest, since they apparently represent a transition stage through which those of the Galapagos once undoubtedly passed. Nothing, unfortunately, is known to the writer as to the geological structure of Guadalupe; the character of the modifications presented in its birds, however, point strongly to its volcanic origin, and render it extremely probable that the upheaval took place at a more recent date than that of the Galapagos. The earliest immigrants to this island were probably the ancestors of *Polyborus lutosus*, which has become completely differentiated in plumage but not perceptibly altered in the details of structure,\* and those of *Carpodacus amplus*, whose modifications of external struc-

---

\* The case of this species presents a very curious problem. Its origin from *P. cheriway*, the only species now inhabiting Middle America, and even northern South America, can scarcely be doubted; but the modifications which the Guadalupe species has undergone tend toward the distinguishing characters of the South American form (*P. tharus*). The two continental representatives of this genus have undoubtedly had a common origin, the differences between them coming under the scope of ordinary geographical laws of variation in this family, as at present understood. The differentiation of the Guadalupe form is of a most remarkable kind, however, being apparently a partial reversion to the features of the Southern form; but some of the characters which distinguish the latter from its Northern analogues are even greatly exaggerated in this Northern insular form! In this instance, then, the differentiation has been a kind of retrocession, with no change in details of structure, while in all the other forms of the island the differentiation has been of the opposite kind, affecting the proportions more than the colors.

ture have almost, if not quite, reached generic distinctness, while the colors have remained essentially unaltered. The same may be said of *Junco insularis*, *Thryomanes brevicauda*, and *Colaptes rufipileus*, while the remaining species, *Regulus obscurus*, *Salpinctes gadalupensis*, and *Pipilo consobrinus* are either more recent arrivals or species in which the process of change has been comparatively slow.

---

---

AN UNDESCRIBED HYBRID BETWEEN TWO NORTH  
AMERICAN GROUSE.

BY WILLIAM BREWSTER.

IN the preparation of the following paper I have hesitated not a little as to the propriety of giving a name to the bird about to be described. That it is a hybrid between the Pinnated Grouse (*Cupidonia cupido*) and the Southern race of the Sharp-tailed Grouse (*Pediocetes phasianellus* var. *columbianus*) is unquestionable, and, further, I consider it almost equally certain that offspring resulting from such unnatural connections are of regular, perhaps even not uncommon, occurrence wherever the two just mentioned species are found together. Indeed, I am aware at the time of writing, of three other similar specimens in private cabinets, and I have heard of additional ones. Although I have examined but one besides my own, I understand that they are all in every way nearly identical, and the fact of their having been procured from different localities must go far towards proving that their occurrence is by no means exceptional or unique. Granting this to be a fact, it seems reasonable that so distinctly specialized a form should bear a distinguishing name, for though certainly the result of a *mésalliance*, and combining in itself characters peculiar to two different species, it is yet unlike either.

But I do not claim originality for a system that has been long established among European authorities. In respect to the name to be adopted I shall follow the practice of Mr. Robert Collett. This gentleman, in writing upon the "Rakkellhane," a hybrid between *Tetrao urogallus* and *T. tetrix*, says,\* it "is a *compound* and

---

\* Remarks on the Ornithology of Northern Norway by Robert Collett. From the Forhandl. Vidensk. Selsk. Christiania, 1872. (Page 50 of the reprint.)

not a simple species, and should, therefore, as such, have a compound and not a simple name." The propriety of this must, I think, impress every one, but in endeavoring to carry out his plan in the present instance I have experienced a serious difficulty.

In naming hybrid forms Mr. Collett makes use of the *generic* title of the *male* parent alone, the "*compound*" part being made up from the *specific* appellations of *both* parents. Thus he calls the offspring of the male Ptarmigan (*Lagopus albus*), paired with the female Black Grouse (*Tetrao tetricus*), *Lagopus tetrici-albus*, the numerous recorded facts at his disposal enabling him to decide upon the respective specific relation of both parents with almost absolute certainty. But in the present case the entire absence of any facts bearing upon this part of the problem reduces me to the somewhat dangerous limits of mere conjecture, or, what is little better, the relative preponderance of specific resemblance exhibited by the specimens before me. Not to weary the reader by a too exhaustive preliminary discussion of detail, I will restrict myself to the simple statement that after careful examination I believe the hybrid Grouse about to be described the offspring of a connection between *Cupidonia cupido*, *male*, and *Pediocetes phasianellus* var. *columbianus*, *female*, and I accordingly bestow upon it, provisionally, the compound name *Cupidonia cupidini-columbiana*.

**DISTINCTIVE CHARACTERS.** — Adult male, from a specimen in my collection, obtained in Iowa. Size and general proportions of *Pediocetes phasianellus* var. *columbianus*. Tail of sixteen feathers exclusive of *two central projecting ones*. Tarsi feathered as in *Cupidonia*. *Neck-tufts* 1.50 inches long. Upper tail-coverts *coextensive* with the rectrices. Above similar to *Cupidonia cupido*; wing-coverts (but not the scapulars) white-spotted, as in *Pediocetes*. Breast and sides barred transversely, as in *C. cupido*; abdomen *white*, sparsely covered with obtuse V-shaped spots of brown. Head, neck, and throat-markings precisely as in *C. cupido*. Neck-tufts dark brown; the longer ones not so stiff as those of *C. cupido*, the shorter dull yellow. Tail generally similar in shape and color to that of *C. cupido*, but with a central pair of elongated feathers "with parallel edges and truncated ends," which project .52 of an inch beyond the next pair. These projecting feathers are tipped with light brown like the other rectrices; subterminally for the space of about an inch they are solidly black, — anteriorly, with ragged rusty-yellow bars. The outer webs of the outer pair of rectrices are irregularly white. The measurements, taken from the dried specimen, are as follows: Wing, 8.57; tail, 3.25, — two central feathers, .52 longer; bill, depth, .40, length from nostril, .50; tarsus, 2.03; middle toe, 2.75.

It will appear from the above description that this bird combines in nearly equal proportions the characters of *Pediocetes* and *Cupidonia*. In the general pattern of coloration of the plumage it most resembles *C. cupido*, but the abdomen is spotted like the breast of *Pediocetes*, and the wing-coverts are marked precisely as in that species. It has the neck-tufts of *Cupidonia* and the projecting tail-feathers of *Pediocetes*, both of these characters, however, slightly modified. A remarkable feature appears in the extension of the upper tail-coverts nearly to the tips of the rectrices.

---

### Recent Literature.

NELSON'S "BIRDS OF NORTHEASTERN ILLINOIS."\*—Under the above title Mr. E. W. Nelson gives us the results of three years' investigation in Cook and Lake Counties in the northeastern corner of Illinois, "a belt about twenty-five miles wide, bordering Lake Michigan in Illinois," including the field considered. As he remarks, the locality seems to form "a kind of four-corners where the avian faunæ of four regions intergrade"; hence we find a somewhat novel juxtaposition of species. On or near the lake occur many birds formerly considered as more or less exclusively maritime. Notably among those of this class found in summer are *Ammodramus caudacutus* and *Egialitis melodus*; during the migrations, *Streptopelia interpres*, *Tringa maritima*, *T. canuta*, *Calidris arcuaria*, and *Micropalama himantopus*; in winter, *Histrionicus torquatus*, *Harelda glacialis*, *Somateria mollissima*, *S. spectabilis*, *Larus glaucus*, and *L. leucopterus*. As might be expected, the species properly belonging to the Carolinian fauna which reach this point are, with a few exceptions, of either uncommon or rare occurrence, and they here seem to touch the extreme northern limit of their range in that longitude. But most interesting are the records of northern birds breeding so far south, especially Limicoline and Natatorial species. Thus Mr. Nelson has found nesting in greater or less abundance, *Tringa minutilla*, *Totanus melanoleucus*, *T. flavipes*, *T. solitarius*, *Marca americana*, *Fulix affinis*, *F. collaris*, *Erismatura rubida*, *Mergus serrator*, and some others.

It is not, however, from the simple enumeration of species, that this list derives its chief value and interest, but from the unusually complete and satisfactory character of the biographical annotations, which embrace good descriptions of the habits of many birds previously but little known.

Thus Mr. Nelson describes the songs of *Turdus aliceæ* and *Oporornis*

---

\* Birds of Northeastern Illinois. By E. W. Nelson. Bulletin of the Essex Institute, Vol. VIII, 1876, Nos. 9-12, pp. 90-155, April, 1877.



*agilis*, the eggs of *Totanus melanoleucus*, and tells us of Night Herons (*Nycticorax grisea nivea*) breeding in the open marshes of Fox River, placing their nests among the wild rice. Emphatically there is no lumber about this paper. It gives, in clear, concise language, the results of extended, carefully and intelligently conducted observations in a region almost wholly unworked, and from its geographical situation and topographical character and surroundings, most rich in results. As an important and valuable faunal contribution to our knowledge of North American Ornithology, Mr. Nelson's list cannot fail to take first rank. — W. B.

SALVIN ON THE PROCELLARIIDÆ. — In the fourth part of the "Ornithological Miscellany," edited by Mr. G. D. Rowley, Professor Osbert Salvin has given the first of a valuable series of papers, in which he seeks to throw all possible light upon this very obscure family. This paper is in two parts. The first is devoted to an examination of the unpublished "Banks' drawings," and the manuscripts of Dr. Solander, so far as they relate to the Petrels. These drawings are sixteen in number, and are presumed to have been drawn by Sydney Parkinson, one of the artists in the employ of Sir Joseph Banks, in the "Endeavor," under Captain Cook. The manuscript notes of Dr. Solander are in the British Museum. As Bonaparte and Gray have introduced Dr. Solander's names into our ornithological nomenclature, even where unaccompanied by descriptions and unpublished, Mr. Salvin has done the world good service in testing the vitality of these names.

*Procellaria oceanica* of Solander stands as *Oceanites oceanicus* of Kuhl. It is better known as *Thalassidroma wilsoni*. *Procellaria aquorea* of Solander, (= *P. marina* of Latham, and confounded by Kuhl with *P. fregata*, a distinct species, in which he was followed by Gray, and for a time by Coues), stands as *Pelagodroma marina*. *Procellaria fregata* of Solander stands as *Fregata grallaria*. *Procellaria turtur* (Sol.) = *Prion turtur*. *P. velox* (Sol.) "must continue doubtful." *P. gigantea* (Sol.) stands as *Ossifraga gigantea*. *P. fuliginosa* Mr. Salvin traces with some difficulty to *Majaquicus equinoctialis*, but without doubt. *P. sandulata*, a "long-lost" species, now reappears in the *Estrelata armingoniana* of Giglioli and Salvadori (Ibis, 1869), and to their name Mr. Salvin gives the preference, following the Golden Rule in questions of nomenclature.

*P. lugens* of Solander cannot be placed. Kuhl and Gray made it the same with *P. grisea* = *Estrelata kidleri* Coues = *Æ. brevirostris* Lesson, — the latter being the proper name, and the same as *P. tristis* Forst. and *P. amaurosoma* Coues; but, according to Mr. Salvin, incorrectly. *P. lugens* must thus be left in abeyance. *Nectris fuliginosa* of Solander Mr. Salvin is convinced = *Puffinus griseus* Finsch. *Nectris munda* Solander may apply to *P. garia* Forst. (= *P. opisthomelus* Coues), but this is regarded as doubtful. *Diomedea antarctica* (Sol.) is probably = *D. fuliginosa*, and *Diomedea profuga* possibly = *D. chlororhyncha*.

Mr. Salvin's second paper is a careful examination of the new species of Petrels obtained by Dr. H. H. Giglioli during the voyage of the Italian corvette "Magenta" round the world, and described in the "Ibis" in 1869. Mr. Salvin examined the type specimens, and accompanies his review with fine illustrations drawn by the well-known bird-painter, Keulemans. These species, according to Mr. Salvin, should all stand as good and novel species. Their names are as follows: *Estrelata magenta*, *E. armingoniana*, *E. trinitatis*, *E. deflippiana*, and *Puffinus elegans*. — T. M. B.

CATALOGUE OF THE BIRDS OF THE ISLANDS OF MALTA AND GOZO. — The Boston Public Library has recently received a copy of Mr. Grant's catalogue of Birds found on the islands of Malta and Gozo. Naturally, a list of the birds occurring in these islands, situated midway between the northern coast of Africa and the southern shores of Europe, divides into three principal groups: migratory birds, properly European; visitants, from Africa; and resident species. Besides these, not always distinguishable, are purely accidental or chance callers. Among the most noticeable of this small group it is interesting to note such purely American species as *Tringoides macularius*, *Actiturus bartramius*, and *Larus atricilla*. *Tryngites rufescens*, American, but not uncommon in Europe, has also been found there. We also notice the occurrence in these islands of birds that are somewhat cosmopolitan as well as North American, such as *Rissa tridactyla*, *Larus marinus*, *Plectrophanes nivalis*, *Tringa canutus*, *T. maritima*, *T. subarquata*, *T. maculata*, *Streptopus interpres*, *Anous stolidus*, and several species of Ducks, common to both continents. — T. M. B.

RIDGWAY'S "STUDIES OF THE AMERICAN FALCONIDÆ." — During the last two years Mr. Robert Ridgway has given us, in a series of special papers, some of the results of his protracted investigation of the American *Falconidæ*. In June, 1875, appeared his "Outlines of a Natural Arrangement of the *Falconidæ*,"\* based on an extended examination of the osteology of the leading types of raptorial birds. In this paper he indorses most fully the classification of the Birds of Prey proposed in 1867 by Professor Huxley, which unites the Old-World Vultures (*Vulturidæ* of most authors) with the *Falconidæ*, separates the Secretary Bird as an independent family (*Serpentariidæ*), and makes the Vultures of the New World a family (*Carthartidæ*) distinct from the other Diurnal Birds of Prey. Mr. Ridgway recognizes among the *Falconidæ* only two subfamilies, — namely, *Falconinæ* and *Buteoninæ*, — thereby differing very widely from most previous writers, some of whose classifications he rather sharply criticises. The subfamily *Falconinæ* he arranges in four minor divisions or "groups," under the names *Falcones*, *Polybori*, *Micrastures*, and *Herpcto-*

\* Bull. U. S. Geol. and Geogr. Survey of the Terr., Vol. 1, No. 4, pp. 225 - 231, pls. xi - xviii, June 10, 1875.

theres, and similarly divides the subfamily *Buteoninae* into three "groups," which he calls *Pandiones*, *Pernes*, and *Buteones*. While Mr. Ridgway endeavors to make use of all available characters as the basis of his divisions, his classification is based primarily upon the character of the coracoid apparatus, particularly as to whether the scapular process of the coracoid is produced forward so as to meet the clavicle, or is separated from it by a wide interval; and also upon the character of certain bones of the skull, especially the supra-maxillary, the nasal, and the superciliary process of the lachrymal, of which parts are given numerous outline illustrations. Whether Mr. Ridgway has quite hit the "natural arrangement" here aimed at, and whether he has not placed undue value upon the modifications of the scapular process of the coracoid, are matters respecting which there is evidently room for honest differences of opinion.

This paper, which closes with a synopsis and diagnoses of the genera and subgenera of the *Polybori*, is followed a few months later by his "Monograph of the *Polybori*."\* This is strictly an American group, of very peculiar appearance and habits. They are chiefly tropical in their distribution, inhabiting all parts of South America, where they most abound, and do not extend northward beyond sub-tropical limits, a single species only (the Caracara, *Polyborus cheriway*) reaching the southern border of the United States. About ten species only are recognized by Mr. Ridgway, which he refers to four genera, namely, *Polyborus*, *Phalcobanus*, *Milvago*, and *Ibycter*. The more important generic characters are illustrated by outline figures, and the various phases of plumage of the species are quite fully described.

A little later, under the general title "Studies of the American *Falconidae*," † Mr. Ridgway continued his account of the American *Falconidae*, describing in detail sixteen genera and some forty species, besides a number of varieties. Thirteen species are referred to the single genus *Nisus* (= *Accipiter* of many authors), while twelve other genera have each but a single species. A few only of the species described in this paper are North American. These are, "*Nisus cooperi*," "*N. fuscus*," *Onychotes gruberi*, *Antenor uncinatus* var. *harrisi*, and *Elanoides forficatus*, only three of which range very far into the United States.

Mr. Ridgway also, in the mean time, published a paper "On *Nisus cooperi* (Bon.) and *N. gundlachi* (Lawr.)," ‡ in which he maintains the entire specific distinctness of "*N. gundlachi*," which had been previously by several authors, including Mr. Ridgway himself, referred to "*N. cooperi*."

This paper is immediately followed by another, "On the Buteonine

\* Studies of the American *Falconidae*: Monograph of the *Polybori*. Bull. U. S. Geol. and Geogr. Survey of the Terr., Vol. I, No. 6, pp. 451-473, pls. xxii-xxvii, February 8, 1876.

† Bull. U. S. Geol. and Geogr. Survey of the Terr., Vol. II, No. 2, pp. 91-182, pls. xxxi, xxxii, April 1, 1876.

‡ Proc. Acad. Nat. Sci. Phila., 1875, pp. 78-88.

Subgenus *Crawircx* (Gould),\* a group represented most numerously in South America. Two species only (*Buteo swainsoni* and *B. pennsylvanicus*), of the six referred to this subgenus, range far into North America. This group is distinguished from the ordinary *Buteones* by having only three (instead of four) of the outer primaries emarginated on their inner webs. While agreeing in this feature, they vary considerably in respect to the relative length of the wing, the size and length of the tarsus, and in other details of structure; and it is perhaps an open question whether our author does not place too much importance upon what he regards as the distinctive feature of the group, namely, the number of emarginated primaries. The various phases of plumage presented by the different species is described in considerable detail, with very full citations of synonymy. The account of *Buteo swainsoni* is particularly full, nearly one half the paper (about fifteen pages) being devoted to this species alone. Among its prominent synonyms are *Buteo bairdii* of Hoy, *B. insignatus* and *B. oxypterus* of Cassin, and *B. fuliginosus* of Selater.

A little later appeared his "Monograph of the Genus *Micrastur*,"† of which seven species are recognized, and of which are given detailed descriptions. The group is mainly restricted to Central and Northern South America, being pre-eminently tropical in distribution. While the species are said to present "no appreciable sexual variation," they exhibit "two well-marked growth-stages," the young birds generally greatly differing from the adult. In addition to this, several of the species are dimorphic, being subject to erythrism, analogous to the dimorphism met with in *Syrnium aluco* of Europe, in *Scops asio*, and in the several species of *Glaucidium*, among the Owls. Shortly after this appeared another paper on this genus, by Mr. Ridgway, entitled "Second Thoughts on the Genus *Micrastur*."‡ This is mainly a summary of his "Monograph." The "Second Thoughts" relate to his earlier revision of this genus, published in 1873, § — in which only five, instead of seven, species were admitted, — rather than to the "Monograph," to which there is in this paper no allusion.

During the elaboration of these important papers on the American *Falconidae*, Mr. Ridgway has had access to the material contained in all the larger public and private ornithological collections of the United States, and has been kindly favored with the use of specimens from abroad. He has in this way been able to base his investigations upon the examination of a larger amount of material than has, in most cases at least, fallen un-

---

\* Proc. Acad. Nat. Sci. Phila., 1875, pp. 89-119.

† Studies of the American *Falconidae*: Monograph of the Genus *Micrastur*. Proc. Acad. Nat. Sci. Phila., 1875, pp. 470-502, figs. 1-9.

‡ Ibis, January, 1876, pp. 1-5.

§ Revision of the Falconine Genera, *Micrastur*, *Geranoospiza*, and *Rupornis*, and the Strigine Genus, *Glaucidium*. Proc. Bost. Soc. Nat. Hist., Vol. XVI, pp. 73-81, December, 1873.

der the inspection of any other investigator in the same field. In these papers we have the results of much hard and patient study, and while experts in the same field may differ from him in respect to minor points of classification and nomenclature, no one can question the fact of his having immensely advanced our knowledge of this large and difficult group. We trust that he will soon be able to give us further instalments of his work on this family, and that eventually we may have the whole reissued connectedly in a well-rounded monograph of the American *Falconidae*. — J. A. A.

RECENT ORNITHOLOGICAL ARTICLES IN AMERICAN JOURNALS.\* —

Among the varied contents of the last and current volumes of the "American Naturalist," † we note the following original articles relating to birds. The number for January, 1876 (Vol. X) contains a note on the "Proper Specific Name of the Song Sparrow" (pp. 17, 18), by David Scott, who claims for this species the name *fasciata*, Gmelin, 1788, in place of *melodia*, Wilson, 1810, the probable tenability of which had been previously suggested by other writers. Also an article on the "Availability of Certain Bartramian Names in Ornithology" (pp. 21-29), by J. A. Allen; a note on the extinction of the Great Auk at the Funk Islands (p. 48), by the same; on the occurrence of Bewick's Wren in the Atlantic States, and the breeding of *ibis ordi* at Camp Harney, Oregon (p. 48), by Dr. Elliott Coues; on the Early Nesting of the Anna Humming-Bird (pp. 48-50), by Dr. J. G. Cooper; and on the occurrence of the European Tree Sparrow (*Pyrgita montana*) at St. Louis, Mo. (pp. 50, 51), — where it has been introduced with the *P. domestica*, — by Dr. J. C. Merrill. The February number (same volume) contains notes on the "Summer Birds of the White Mountain Region" (pp. 75-80), by H. D. Minot; an article on "Californian Garden Birds" (pp. 90-97), by Dr. J. G. Cooper; a reply to Mr. Allen's article in the preceding number on Bartramian names (pp. 98-102), by Dr. Elliott Coues; also, by the same, notes on the "Breeding Range of the Snowbird," *Junco hyemalis* (p. 114); on "Protective Resemblance in the Yellow-Bird," *Chrysomitris tristis* (p. 115), by Henry Gill-

---

[\* In addition to notices of papers published in the transactions of different scientific societies, or as special works, it is intended to note in the Bulletin all original ornithological articles appearing in various periodicals (beginning with the year 1876), thereby making the Bulletin a complete record of North American ornithological literature. Owing to limited space, it will be impracticable to give generally more than the titles of articles. The record here begun will be continued in future numbers. — Eds.]

† The American Naturalist. Edited by Dr. A. S. Packard, Jr. Boston: H. O. Houghton & Co.; New York: Hurd and Houghton. Vol. X and Vol. XI, January to June. (The departments of Ornithology and Mammalogy are now in charge of Dr. Elliott Coues.)

man. In the March number, Dr. W. Wood writes of the "Game Falcons of New England: the Goshawk" (pp. 132-135); J. A. Allen appears with a further note on Bartramian names (pp. 176, 177), and R. E. C. Stearns speaks of the abundance of Pelicans in San Francisco Bay, Cal. (p. 177). In the April number are notes on the breeding of the Red Crossbill (*Loxia curvirostra*) at Riverdale, N. Y. (p. 237), by E. A. Bicknell; on Bewick's Wren in New Jersey (p. 236), by Dr. C. C. Abbott; on "Habits of Western Birds,"—*Buteo swainsoni*, *Icterus bullocki*, *Stellula caliope*, *Corvus americanus*, and *C. corax*,—(p. 238), by Dr. W. J. Hoffman; on "Unusual Nesting-Sites of the Night-Hawk and Towhee Bunting (p. 239), by Dr. Elliott Coues; and on "Small Birds [*Chrysomitris tristis* and *Dendroica coronata*] caught by the Burdock" (p. 239), by A. K. Fisher. The May number contains a call for information respecting the distribution of the Labrador Duck, *Camptolomus labradorius* (p. 303), by Dr. Coues. In the June number is a record of the capture of the European Woodcock in Virginia (p. 372), and a note on change of habits in the Bank Swallow (p. 373), by Dr. Coues; also remarks on the food of the Chapparral Cock, *Geococcyx californianus* (p. 373), by V. T. Chambers. In the July number J. Clarence Hersey notes the occurrence of the Little White Egret (*Ardea candidissima*) in Colorado (p. 430). The August number contains an account of a brief sojourn of large numbers of White Egrets at Trenton, N. J. (pp. 469-473), by Dr. C. C. Abbott; Robert Ridgway here states that the Bank Swallows (p. 493) referred to in the June number are the *Cotyle serripennis*, instead of *C. riparia*, as first stated. In the September number appears an historical article on "Progress of Ornithology in the United States during the Last Century" (pp. 536-550), by J. A. Allen, and a note on Bluebirds feeding on the berries of the Virginia Creeper (p. 556), by Henry Gillman. The October number contains a note on the egg of *Chionis* (p. 628), by Dr. J. H. Kidder. An article in the December number, by Dr. Elliott Coues, on the "Destruction of Birds by Telegraph Wire" (pp. 734-736), concludes the ornithological articles of Volume X.

In Volume XI we find, in the January number, "Notes on some Oregon Birds" (p. 44),—seven species,—by George R. Bacon, and a note on the Whistling Duck, *Bucephala americana* (p. 44), by J. F. LeBaron. In the February number, notes on some of the birds of the Fanning Islands (pp. 68-72), by Dr. T. H. Streets; in the April number, record of the occurrence of the Raven and the Sooty Tern (*Sterna fuliginosa*) at Williamstown, Mass. (p. 243), by Sauborn Tenney; in the May number, "Glimpses of Mind in Birds" (pp. 276-286), by Dr. C. C. Abbott; some observations on the winter birds of Arkansas (p. 307), by H. S. Reynolds; and a note on a carnivorous propensity of the Red-headed Woodpecker, *Melanerpes erythrocephalus* (p. 308), by Charles Aldrich. In the June number Judge J. D. Caton writes of the "Wild Turkey and its Domestication" (pp. 321-330).

f "Field and Forest" \* continues to devote a considerable portion of its space to ornithology. Volume II (beginning July, 1876) contains the following: "On the Habits of *Steganopus wilsoni*" (pp. 11, 12), by A. L. Kumlien (already noted in this journal, Vol. I, p. 71); "Notes on Forster's Tern" (pp. 29-31), by Pierre Louis Jouy; "Sexual, Individual, and Geographical Variation in the Genus *Leucosticte*" (pp. 37-43), by Robert Ridgway; "Drumming of the Ruffed Grouse" (pp. 57-60), by David Scott; "Crows Seeking Water" (p. 65), by N. B. Webster; "Sexual Variation in the Genus *Leucosticte*" (pp. 76-79), by J. A. Allen; the "Tendency of Birds to vary in their Habits" (pp. 107-114), by David Scott; a Congress of Birds (p. 122), by L. F. Ward; "Unusual Accidents to Birds" (p. 106), by A. J. Kumlien; "Ornithological Notes from Texas" (pp. 154-156), by Ludovic Kumlien; "Catalogue of the Birds of the District of Columbia" (pp. 154-156, 178-181), by Pierre Louis Jouy, — a nominal list of 240 species; Remarks on the Birds of the District of Columbia (pp. 191-194), by Drs. E. Cones and D. W. Prentiss; "Mrs. Maxwell's Colorado Museum, — Collection of Birds" (pp. 194-199), by Robert Ridgway. — J. A. A.

Since January, 1876, several ornithological lists, of more or less importance, have appeared from time to time in "Forest and Stream." † While our space will not permit of a lengthy notice of these lists, a simple enumeration of them will be useful for reference. In Volume VI, pp. 99, 132, 163, 214, 266, 318, 354, 402, and Volume VII, pp. 147, 164, 276, Mr. Adolphe B. Covert, of Ann Arbor, Mich., has given an interesting List of the Birds of Lower Michigan, including two hundred and twelve species. In this list are recorded specimens of *Dendroica kirtlandi*, *Scelopax rusticola* (one specimen obtained by Dr. Wm. E. Lewitt, May 9, 1870), and *Camptolarnus labradorius* (one specimen taken at Delphi Mills, Mich., April 17, 1872). In Volume VI, p. 148, Mr. G. Aug. Smith gives a List of the Birds of Fort Wayne, Ind., including fifty-two species. In Volume VI, pp. 233, 284, 337, 402, and Volume VII, pp. 36, 52, 84, 180, 230, Mr. H. G. Fowler, of Auburn, N. Y., gives a Partial List of the Birds of Central New York, from observation made in the counties of Cayuga, Seneca, and Wayne. In this list Mr. Fowler records the capture of *Querquedula cyanoptera* on Seneca River. This, we believe, is the only record for this species east of the Rocky Mountains. He also notes the occurrence of *Sterna fuliginosa* on Owasco Lake, a specimen having been taken there

---

\* Field and Forest. A Monthly Journal devoted to the Natural Sciences. Charles R. Dodge, Editor. Washington, D. C. Vol. II, Nos. 1-12, July, 1876, to June, 1877.

† Forest and Stream, a Weekly Journal devoted to Field and Aquatic Sports, Practical Natural History, Fish Culture, the Protection of Game, etc., etc. New York: Forest and Stream Publishing Company.

on September 20, 1876. The list comprises one hundred and sixty-eight species. In Volume VI, p. 300, Mr. H. Ernst, of Cleveland, Ohio, gives a List of the Warblers of the Western Reserve, in which he notes thirty-one species. In Volume VII, pp. 389 and 404, Colonel A. S. Brackett, U. S. A., gives a List of Birds of Southeastern Wyoming, including eighty-four species. In Volume VIII, pp. 33, 49, 96, 113, a List of the Birds of Webster, N. H., and adjoining towns, is given by Mr. Chas. F. Goodhue. This list comprises one hundred and thirty-three species. In Volume VIII, pp. 176, 192, 224, 241, 261, is a List of the Birds of the Coteau des Prairies of Eastern Dakota, by Chas. E. McChesney, M. D., U. S. A., of Fort Sisseton, comprising one hundred and three species. — R. D.

CALIFORNIAN ORNITHOLOGY. — Dr. J. G. Cooper has recently published a paper\* of fourteen pages, entitled "New Facts relating to Californian Ornithology. — No. 1." This is supplemental to the same author's "Ornithology of California," and "includes only observations not previously published, and such opinions as differ from those of later authors." About fifty species are noticed, the notes respecting them relating mainly to their seasonal distribution and habits, but include descriptions of the nests and eggs of several species, with occasional remarks upon disputed points of nomenclature. He claims the name *nanus* for the species of Thrush usually known as *Turdus pallasi* (as has also Dr. T. M. Brewer), but inclines to the opinion that the name *guttatus* of Pallas (1811) will finally prove to be the only tenable name for the species. He also claims, on the ground of priority, that *sandwichensis* should take the place of *savanna* for the species of Sparrow, commonly known as *Passerculus savanna*. The paper is replete with interesting matter, and forms a valuable contribution to our knowledge of Californian Ornithology. — J. A. A.

MCCAULEY'S NOTES ON TEXAN ORNITHOLOGY.†—Lieutenant C. A. H. McCauley has just given us the results of six weeks' observations made in May and June, 1876, on the ornithology of the country about the source of the Red River of Texas, embracing a portion of the region known as the Staked Plain. The paper includes notices of about one hundred species, with quite copious notes respecting the habits of a considerable proportion of them, with, in some cases, descriptions of their nests and eggs. This is almost the first special paper treating of the ornithology of Western

\* Proc. Cal. Acad. Sci., 1876. Only the author's separates have yet been seen by the writer.

† Notes on the Ornithology of the Region about the Source of the Red River of Texas, from Observations made during the Exploration conducted by Lieutenant E. H. Ruffner, Corps of Engineers, U. S. A. By C. A. H. McCauley, Lieutenant Third United States Artillery. Annotated by Dr. Elliott Coues, U. S. A. Bull. U. S. Geol. and Geogr. Survey, Vol. III, No. 3, pp. 655-695, May 15, 1877.



Texas, and forms a most welcome contribution to our knowledge of the ornithology of that region. The list shows an unusual mingling of eastern and western species, and, as would be naturally expected, a considerable proportion of strictly southern forms. On the barren Staked Plain few birds were met with, and these mainly about water-holes; along the timber-skirted streams, however, bird life was abundant. — J. A. A.

---

---

### General Notes.

NEST AND EGGS OF TOWNSEND'S FLYCATCHER. — In July, 1876, while rambling with my brother over the mountains of Summit County, Colorado, it was my good fortune to find, at an altitude of about ten thousand feet, the nest of Townsend's Flycatcher (*Myiadestes townsendi*), and as no description of its eggs has as yet appeared, perhaps the following may not be uninteresting: The nest was very loosely, and, externally, shabbily built of long dry grasses, straggling two feet or more below it. It was placed in the upper bank of a miner's ditch (running from the Bear River, above Breckenridge, to the Gold Run and Buffalo Flat diggings), and was partly concealed by overhanging roots; yet it was rendered so conspicuous by the loose swaying material of which it was composed, as well as by that which had become attached to the overhanging roots during its construction, as to attract the eye of an experienced collector when yet some rods away. On nearing the nest the bird immediately took flight, and alighted on the topmost branch of the nearest pine. Resting uneasily here for half a minute, it then, in short, uncertain flights, worked its way down the mountain side and out of sight. Withdrawing to a convenient cover, we had only to wait a few moments for the bird to return, perch herself on a branch a few feet from the nest, peer anxiously into it, and then quickly resume her task of incubation. Moving cautiously along the bank above the ditch, we tried to capture the bird by placing a hat over the nest, but, miscalculating its location by a few inches, the bird eluded the stroke and made good her escape, as she did also on our second attempt to capture her. Again retreating to cover, we waited for half an hour for the bird to return, when suddenly we espied it flying from branch to branch, displaying by its restless motions more anxiety and suspicion than before, yet constantly working nearer its home, which it soon reached and settled quietly again to business. After the last unsuccessful attempt to catch the bird, a stick was placed on the bank directly over the nest, to mark its exact locality, and this time, moving with less haste and more caution, we gained the desired position, lay down on the bank, and taking a hat in each hand quickly covered the opening and secured the unfortunate bird, and also the opportunity of giving to ornithologists an authentic

account of the number, size, and coloration of the eggs. The nest contained four eggs, very closely resembling those of the Shrikes. The ground-color is dull white or bluish, thickly blotched or freckled with reddish-brown. The measurements of the three specimens preserved are 1.01 by .66, .94 by .68, and .88 by .66. Incubation had been going on for about ten days, and unfortunately one egg was destroyed in cleaning.—WILBUR F. LAMB, *Holyoke, Mass.*

PERSISTENCY AT NEST-BUILDING IN A HOUSE-WREN.—A House-Wren (*Troglodytes ædon*) has this season manifested a strong predilection for the nozzle of a pump for a nesting-site. The pump being in daily use, the nozzle, much to our surprise, was repeatedly found to be obstructed with sticks. An investigation of the novel incident led to the discovery of the cause, it being found that a House-Wren was industriously at work carrying materials into the pump for the construction of its nest. The bird was finally left one morning to carry on his work, when, at the end of two hours, it was found that he had filled the pump so full that water could not be obtained until a part of the sticks had been removed. The nest, through the necessary use of the pump, was three times destroyed before the persevering little fellow abandoned his work.—ABBOTT W. FRAZAR, *Watertown, Mass.*

A NEW BIRD TO MASSACHUSETTS.—Mr. Charles W. Townsend, of Boston, shot, July 28, 1876, a male specimen of *Plectrophanes ornatus*. It was taken in Magnolia, near Gloucester, Mass., in a field near the seashore, and has been by Mr. Townsend presented to the New England collection of the Boston Natural History Society. It is an adult male, in worn plumage.—T. M. BREWER, *Boston, Mass.*

A NEW FORM OF SURNIA TO NEW ENGLAND.—Two fine specimens of the Hawk Owl have recently been taken in Houlton, Maine, and have been mounted by Mr. Welch in his usual superior style. They are both males, and while one is in the plumage usually known as *Surnia hudsonia*, the other is in that distinguished by the separate name of *Surnia ulula*, and supposed to be exclusively Palearctic.—T. M. BREWER, *Boston, Mass.*

CAPTURE OF THE PHILADELPHIA VIREO IN NEW HAMPSHIRE.—A specimen of this bird (*Vireo philadelphicus*) was shot in Hollis, New Hampshire, May 26, 1876, by Mr. A. F. Eaton. It was feeding in company with two other birds of the same kind, in some low oak-bushes.—W. H. FOX, *Concord, Mass.*

OCCURRENCE OF PASSERCULUS PRINCEPS IN NEW YORK.—One of my correspondents, Mr. Frank E. Johnson, of Gravesend, Long Island, writes me that when out collecting, on December 20, 1876, on Coney Island, in New York harbor, he shot three specimens of a Sparrow new to him,

which were shown to Mr. George N. Lawrence, and pronounced to be the Ipswich Sparrow (*Passerculus princeps*). They were shot on the salt meadows of the island, and were in company with Savannah Sparrows (*Passerculus savanna*) and Swamp Sparrows (*Melospiza palustris*). This is the most southern record of this species.\* — H. B. BAILEY, *Newton, Mass.*

THE PIGEON-HAWK (*Falco columbarius*) AT SEA. — While returning from a trip to Labrador, last summer, I observed small Hawks, undoubtedly of this species, at a considerable distance from land, on two occasions.

The first occasion was on the 5th of September. We were crossing the Gulf of St. Lawrence, and were in sight of the coast of Newfoundland, which was about twelve or fifteen miles distant. As many as four Hawks were seen, which came so near that we were able to recognize them as Pigeon-Hawks. They seemed to be perfectly at home, flying over the water, and showed no fear of the vessel, several times alighting on the rigging.

The first that appeared had a Leach's Petrel, dead, in his talons. He alighted with this, on the fore-crossrees, and proceeded to eat it. The sailors were unwilling that we should fire into the rigging, so a young man went up the fore-rigging, and nearly caught the Hawk, which flew off, leaving his prey behind him. Three other Hawks came off to the vessel during the day, and were all shot, but all, unfortunately, fell into the water and could not be secured. The day was bright, clear, and warm, with a light wind from the north, so that we made very little progress. The Hawks appeared to come from the direction of Newfoundland.

The second occasion was during our run from Cape Sable to Boston, about fifty miles from the nearest land. It was the 10th of September, a bright day, with a strong northwest wind. A small Hawk, probably a Pigeon-Hawk, passed the vessel, flying to windward. Dr. Cones, in his "Notes on the Ornithology of Labrador," mentions that a Hawk of this species came on board their vessel during their return voyage, in a very exhausted condition. This bird, however, was very shy, and was immediately frightened away from the vessel. He also mentions seeing several Sparrow-Hawks while in the Gulf of St. Lawrence, off Cape Breton Island, which "circled quite closely around the vessel, showing but little fear." — JOHN MURDOCH, *Cambridge, Mass.*

CAPTURE OF A SECOND SPECIMEN OF HELMINTHOPHAGA LEUCOBRONCHIALIS. — In the first number of the Bulletin for the year 1876, Mr. Wm. Brewster described a new species of *Helminthophaga* (*H. leucobronchialis*), which he obtained in Newtonville, Mass., on May 18, 1870. He says in his article, "Whether it must be placed in the same category with the

---

\* For other records of occurrence of this species see this Bulletin, Vol. I, pp. 18, 52, and Vol. II, p. 27.

unique *Euspiza townsendi*, *Regulus curvirostris*, etc., or, like *Dendroica kirtlandi*, will turn up occasionally in the future at different points, or still again, as in the case of *Centronyx bairdii*, will be found in large numbers, time alone can decide." It is with pleasure, therefore, that I can announce the capture of a second specimen of this species, so new to Ornithology, and particularly also because it was taken in a locality so far distant from where the first one was obtained. The specimen under consideration was shot by Mr. Christopher D. Wood, on the afternoon of May 12, 1877, in an apple orchard near Clifton, Delaware County, Pa. It proved to be a male, and answered to the description given by Mr. Brewster. It is, without doubt, a veritable specimen of *H. leucobronchialis*, and goes to prove the species a good one. It was first called to my attention by Mr. Wood himself, who told me that he had shot a specimen of *H. leucobronchialis* near Clifton. He afterwards showed me the bird, which he had been comparing with the plate of the former specimen, and found it to be identical. Whence these rarities come, whether they are abundant in certain sections, and the characters of the females, are matters not yet known; yet it is more than likely that at no very distant day both the present species, as well as *Helminthophaga lawrencei*, may prove to be nearly if not quite as abundant as the other species of the same genus. — SPENCER TROTTER, Philadelphia, Pa.

THE MOTTLED OWL AS A FISHERMAN. — On November 29, 1876, I took from a Mottled Owl's hole (*Scops asio*) the hinder half of a Woodcock (*Philohela minor*). Within two weeks after I took two Owls from the same hole, and on the 19th of January last I had the good fortune to take another. After extracting the Owl I put in my hand to see what else there was of interest, and found sixteen Horned Pouts (*Amiurus atrarius*), four of which were alive. When it occurred to me that all the ponds in the vicinity were under at least two feet of snow and ice, I could scarcely conjecture where the Horned Pouts could have been captured. After visiting all the ponds, I found they had most probably been captured in one fully a mile away, where some boys had been cutting holes through the ice to catch pickerel bait. The Owl probably stationed himself by the edge of the hole and seized the fish as they came to the surface. What a busy time he must have had flying thirty-two miles after sixteen Horned Pouts! I may also state in this connection that I once found the ground under a Great Horned Owl's nest (*Bubo virginianus*) literally strewn with fish-bones. — A. M. FRAZAR, Watertown, Mass.

BREEDING OF LEACH'S PETREL ON THE COAST OF MAINE. — In the January number of the Bulletin (Vol. II, 1877) Mr. N. C. Brown refers to the Leach's Petrel (*Thalassidroma leucorhoa*, Linn.) "as found for the first time breeding on the New England coast," and mentions meeting with its

nests on the Green Islands in Casco Bay. That Mr. Brown was not the first person to find it breeding even on the Green Islands would not be a fact of sufficient moment to call for correction did not his statement suggest the quite important error implied: that it is not known, and has not been known, to breed elsewhere on the coast of Maine. That this Petrel breeds along the greater part of the coast of Maine has been known as a fact for many years. Whether Casco Bay is its most western point remains to be ascertained.

In June, 1850, I made several weeks' explorations in the neighborhood of Eastport, and found this bird breeding in all the Grand Menan group — which geographically, if not politically, are part of the Maine coast — on the island of Eastport itself, and on a small island between Eastport and Machias. An account was published (*Bost. Jour. Nat. Hist.*, Vol. VI, p. 297). On the following year, in company with Dr. H. R. Storer, I continued these explorations, and ascertained that this species breeds abundantly on every suitable island as far west as Mt. Desert. Several years afterwards, in the summer of 1855, and again in 1856, in company with Dr. Dixon, of Damariscotta, we traced their breeding, in considerable numbers, as far west as Round Pond harbor, in Bristol, and in the Damariscove Islands, in the ocean, not far from the mouth of the Kennebec. In 1873, on Peakes Island, I saw specimens of the eggs and birds taken by Messrs. Franklin Benner and Spencer Baird Biddle in Casco Bay the same summer. On the strength of these observations, made by others as well as myself, in my Catalogue of the Birds of New England (*Proc. Bost. Soc. Nat. Hist.*, Vol. XVII, p. 450), I spoke of this Petrel as a summer resident on the coast of Maine. I have by me, in MS., the letters of both these gentlemen in regard to their observations. I subjoin a brief extract from the notes of Mr. Benner:—

“The first visit was made to Junk-of-Pork Island, about three miles northeast of Peakes, in Casco Bay, on July 16, 1873. The island has an area of half an acre at low water, and in the centre is an almost perpendicular piece of rock about forty feet in diameter and nearly twenty-five feet high. A dozen or more burrows of this Petrel, each with their single egg, were found in the earth that had accumulated on the top of this rock. The eggs were about half incubated. In two nests young were found only a day or two old. One of the parent birds was found in each burrow, and in one instance both.

“On the 22d of the same month I visited White Bull Island, located twelve miles farther to the eastward, and comprising a much larger extent of surface than the first. Here were also found the nests of the Petrel among many of the Terns. Young birds were found in many of them, and some eggs.”

He speaks of having found them “abundant,” and probably breeding in several other “of the many barren islands in the neighborhood.”— T. M. BREWER, *Boston, Mass.*

NEST AND EGGS OF THE ALASKAN WREN. — In a small collection of birds' skins, nests, and eggs recently acquired by the Museum of Comparative Zoölogy, collected at the Pribylow Islands, Alaska, is the nest and two eggs of the Alaskan Wren (*Troglodytes parvulus* var. *alascensis*), which are believed to be the first ever seen by naturalists. The nest is quite large and very compactly built, being composed externally of fine moss of a bright green color, interwoven with fine roots, and lined heavily with hair and feathers. Conspicuous among the latter are the rosy-tipped feathers of the *Leucosticte griseinucha*. The hairs are rather coarse and white, three to four or five inches in length, and appear to be hairs of the Polar Bear. The nest was obtained in June, 1876, on St. George Island, by Mr. W. J. McIntyre, to whom it was brought by a native. It is said to have been placed deep down in the crevices of large rocks, and to have originally contained twelve eggs, all but two of which were broken before they came into Mr. McIntyre's possession. These measure, respectively, .68 by .51 and .60 by .50. Their general color is dull white, with a very few minute dots of reddish, so few and small as to be easily overlooked. The nest is represented to be very hard to find, being placed so deeply among the rocks, this being the only one Mr. McIntyre could obtain during two years' residence at the Islands, although he had a standing offer for them of about ten dollars in gold each. — J. A. ALLEN, *Cambridge, Mass.*

JUNCO OREGONUS IN ILLINOIS. — October 14, 1875, I saw a flock of some dozen birds in a willow-tree, and killed one with a sling. The rest flew off, and were not seen again. The specimen was sent to Mr. E. W. Nelson, who identified it as *Junco oregonus*, the first one of this species captured in this State, its extreme eastern range as heretofore known being Kansas. — H. K. COALE, *Chicago, Ill.*

LEPTOPTILA ALBIFRONS, A PIGEON NEW TO THE UNITED STATES FAUNA. — Mr. George B. Sennett, a diligent and zealous ornithologist, who has been making collections and observations in Southern Texas, writes as follows from Hidalgo, Tex., under date of May 2, 1877 : —

"I have a dove which I do not identify, and accordingly send you a description of a specimen killed April 18. This is a male. I have secured four specimens, and hope to find the nest, as I am satisfied they breed here. Their cooing is low and short, ending with a falling inflection, and is easily recognized by its peculiarity. . . . Length, 12.50; extent, 19.50; wing, 6.35; tail, 4.50; tarsus, 1.37, middle toe and claw the same; bill, .62, black. Iris yellow. Orbital space small, faintly red and blue. Tail square, of twelve feathers. Upper parts greenish-olive, the metallic coloring purple with bronzy-green reflections, and restricted to back of neck. Crown drab, shading to nearly white on the forehead. Chin white. Foreneck creamy-slate. Belly white. Sides ashy. Wings brown, slaty below, and whole underwing-coverts are bright chestnut, which color ex-

tends even on the sides. Middle tail-feathers like the back ; others brown above, and tipped with white in increasing amount, till the outer ones are white for half an inch ; tail below black, with the white tips, as just said. Under tail-coverts pure white. In general habits, the bird is quiet and not readily alarmed ; it associates with the White-winged Doves (*Melopelia leucoptera*), and prefers tall trees to undergrowth."

I sent my correspondent's letter to Mr. Ridgway, who kindly compared the description with specimens of *Leptotila albifrons* in the National Museum, and made this identification. — ELLIOTT COUES, *Washington, D. C.*

MELOPELIA LEUCOPTERA IN COLORADO. — Mr. E. L. Berthoud, writing from Golden, Col., March 7, 1877, informs me of the occurrence of this species near timber line on the head of Cub Creek, Jefferson County. He saw a dozen or more of the birds — rare in this region — in July, 1869. This verifies my surmise (*Birds of the Northwest*, p. 386) of the actual occurrence of the species beyond hitherto recorded limits. — ELLIOTT COUES, *Washington, D. C.*

THE RUFF AND THE PURPLE GALLINULE IN OHIO. — Dr. Theodore Jasper, of this city, obtained, November 10, 1872, at the Licking Reservoir, thirty miles east of Columbus, a Wader which remained unidentified till recently. I was of the opinion that it would prove to be either *Philomachus pugnax*, or a nondescript. On communicating my views to Mr. H. W. Henshaw, of Washington, he kindly offered to compare the specimen with others in the National Museum. He writes that the bird, which was a male (probably young), is positively identical with specimens of that species in the collection of the National Museum.

I have also just received from my friend, Dr. Howard E. Jones, a fine skin of the Purple Gallinule (*Porphyrio martinica*), killed by him at Circleville, Ohio, May 10, 1877. This bird is now recorded for the first time, on unimpeachable authority, as a visitor to this State. Dr. Jones tells me that it has been seen before in the vicinity of Circleville. In my Catalogue of the Birds of Ohio (*Ohio Agric. Rep.*, 1860), it was inserted on what I afterward discovered to be insufficient authority, and for that reason it was omitted from a subsequent list (*Food of Birds*, etc. 1875). I have several times been favored with reports, and once or twice with skins, presumed to be of this species, which proved, however, to be those of the Florida Gallinule, which is not a rare summer resident throughout the State. — J. M. WHEATON, *Columbus, O.*

NOTES ON NYCTALE ACADICA. — Although not generally common in any locality, the Acadian or Saw-whet Owl has been of quite frequent occurrence in Chicago and immediate vicinity during the past three years. A female of this species in my collection was caught alive while sleeping on one of the lower branches of a pine-tree, June 23, 1874. In July of the same year three adult specimens were shot by a boy, who saved only

the wings, as he did not understand preserving skins. November 4 an adult female was shot in a small grove of pine-trees near the south city limits. It measured,  $8.12 \times 18.60$ ; wing, 5.50. March 26, Mr. C. C. Whitacre caught an adult male at the same place. It was kept alive for several days, when it died from being caught between the bars of its cage. He afterwards shot two adult females in the same grove (which has since gone by the name of the Acadian Grove), and also found one dead in his yard. March 28 Mr. J. B. Osborne shot an adult female, and June 15 a young female, both of which are now in my possession. The latter measured,  $7.37 \times 19.25$ ; wing, 5.62. Disc dark brown; forehead, wings, and tail beginning to show white markings, as in the adult. July 10 a second young specimen was brought to me alive. Although just captured, it showed no fear on being handled. In the shade the iris was hardly visible, while in the sun the pupil contracted so much as to appear as only a small black spot. The bird always sat, when perched, with two toes before and two behind, puffing out its feathers at times so that it looked nearly as round as a ball. The white markings were more clearly defined than in the other, extending farther back on the forehead, and entirely round the outer edge of the disc. This specimen is in the collection of Dr. J. W. Velie, who also found one dead on the lake shore at Hyde Park, Ill. July 16 a third juvenile was shot in a poplar-tree opposite my residence. It was still more advanced toward adult plumage than either of the others, especially about the head and wings. J. Strickland (taxidermist) has a young specimen which was caught here about a year ago. December 20 Mr. G. F. Clingman shot an adult female in Acadian Grove, and March 4, 1876, a second specimen was shot in the same place. These, with one male and two females which are also in collections here, make a total of twenty, including two adult males, fourteen adult females, and four in immature plumage. — H. K. COALE, *Chicago, Ill.*

PROBABLE BREEDING OF THE ACADIAN OWL (*Nyctale acadica*) IN MASSACHUSETTS. — The capture of this species in the adult state is by no means of rare occurrence in Massachusetts, but its presence is generally detected in the winter months. Of its breeding so far south in New England I think there has hitherto been no instance recorded. We are now able, however, to note the capture of three specimens in the plumage of the so-called "*albifrons*." The first was taken in Newton, Mass., on June 28, 1876; and the second at Hingham, Mass., on July 5, 1876; the third was captured in one of the cells in the Penitentiary on Deer Island, Boston Harbor, on the 8th of the same month, by an inmate of the prison. These localities being some ten or fifteen miles apart, it would seem hardly probable that these three Owls belonged to the same brood. On April 4, 1877, a specimen in adult plumage was captured in the Penitentiary on Deer Island, where the above-mentioned immature specimen was taken. I am indebted to Mr. Wm. J. Knowlton, of Boston, for the above facts, and from him I obtained one of the young specimens. — RUTHVEN DEANE, *Cambridge, Mass.*



# BULLETIN

OF THE

## NUTTALL ORNITHOLOGICAL CLUB.

---

Vol. II.

OCTOBER, 1877.

No. 4.

---

### NOTES ON *MOLOTHRUS ÆNEUS*, WAGL.

BY J. C. MERRILL, ASSISTANT SURGEON, U. S. A.

THE occurrence of this species north of Mexico was noted in the Bulletin of November, 1876 (Vol. I, p. 88). It is now more than a year since it was first observed, and during that time I have had ample opportunity to study its habits, a short account of which may be of interest. This Cowbird is found in Mexico, Guatemala, and Veragua, as well as in Southern Texas; how far it penetrates into the latter State I am unable to say. My first specimens were taken at Hidalgo, on the Rio Grande, seventy miles northwest of Fort Brown, where, however, they are not so abundant as lower down the river. Here they are common throughout the year, a small proportion going south in winter. Those that remain gather in large flocks with the Long-tailed Grackles, common Cowbirds, and Brewer's, Red-winged, and Yellow-headed Blackbirds; they become very tame, and the abundance of food about the picket-lines attracts them for miles around. *M. æneus* is readily distinguishable in these mixed gatherings from the other species by its blood-red iris and its peculiar top-heavy appearance, caused by its habit of puffing out the feathers of the head and neck. This habit is most marked during the breeding-season and in the male, but is seen throughout the year.

About the middle of April the common Cowbird, Brewer's, and Yellow-headed Blackbirds leave for the North; the Long-tailed Grackles have formed their colonies in favorite clumps of mesquite trees; the Redwings that remain to breed have selected sites for their nests; the dwarf Cowbirds (*Molothrus pecoris* var. *obscurus*)

arrive from the South, and *M. œneus* gather in flocks by themselves, and wait for their victims to build. The males have now a variety of notes, somewhat resembling those of the common Cowbird (*Molothrus pecoris*), but more harsh. During the day they scatter over the surrounding country in little companies of one or two females and half a dozen males, returning at nightfall to the vicinity of the picket lines. While the females are feeding or resting in the shade of a bush, the males are eagerly paying their addresses by puffing out their feathers, as above noted, strutting up and down, and nodding and bowing in a very odd manner. Every now and then one of the males rises in the air, and, poising himself two or three feet above the female, flutters for a minute or two, following her if she moves away, and then descends to resume his puffing and bowing. This habit of fluttering in the air was what first attracted my attention to the species. In other respects their habits seem to be like those of the eastern Cowbird (*M. pecoris*).

My first egg of *M. œneus* was taken on May 14, 1876, in a Cardinal's nest. A few days before this a soldier brought me a similar egg, saying he found it in a Scissor-tail's (*Milvulus*) nest; not recognizing it at the time, I paid little attention to him, and did not keep the egg. I soon found several others, and have taken in all twenty-two specimens the past season. All but two of these were found in nests of the Bullock's, Hooded, and small Orchard Orioles (*Icterus spurius* var. *affinis*). It is a curious fact that although Yellow-breasted Chats and Red-winged Blackbirds breed abundantly in places most frequented by these Cowbirds, I have but once found the latter's egg in a Chat's nest, and never in a Red-wing's, though I have looked in very many of them. Perhaps they feel that the line should be drawn somewhere, and select their cousins the Blackbirds as coming within it; the Dwarf Cowbirds are not troubled by this scruple, however. Several of these parasitic eggs were found under interesting conditions. On six occasions I have found an egg of both Cowbirds in the same nest; in four of these there were eggs of the rightful owner,\* who was sitting; in the other two the Cowbird's eggs were alone in the nests, which

---

\* It would be interesting to know what would have become of the three species in one nest, and had the latter been near the fort, where I could have visited them daily, I should not have taken the eggs. It is probable, however, that *M. œneus* would have disposed of the young Dwarf Cowbird as easily as of the young Orioles.

were deserted : but I have known the Hooded Oriole to sit on an egg of *M. aeneus*, which was on the point of hatching when found ; how its own disappeared I cannot say. Once two eggs of *aeneus* were found in a nest of the small Orchard Oriole (var. *affinis*). Twice I have seen a broken egg of *aeneus* under nests of Bullock's Oriole on which the owner was sitting.

Early in June a nest of the Hooded Oriole was found with four eggs and one of *M. aeneus*, all of which I removed, leaving the nest. Happening to pass by it a few days later, I looked in, and to my surprise found two eggs of *aeneus*, which were taken ; these were so unlike that they were probably laid by different birds. Still another egg, and the last, was laid in the same nest within ten days. But the most remarkable instance was a nest of the small Orchard Oriole, found June 20, containing three eggs of *aeneus*, while just beneath it was a whole egg of this parasite, also a broken one of this and of the Dwarf Cowbird (var. *obscurus*). Two of the eggs in the nest were rotten ; the third, strange to say, contained a living embryo. As the nest was certainly deserted, I can only account for this by supposing that the two rotten ones were laid about the first week of June, when there was considerable rain, and that the other was deposited soon after, since which time the weather had been clear and very hot. On one occasion I found a female *aeneus* hanging with a stout thread around its neck to a nest of the Bullock's Oriole. The nest contained one young one of this Cowbird, and it is probable that its parent, after depositing the egg, was entangled in the thread on hurriedly leaving the nest, and there died ; it had apparently been dead about two weeks. This case supports the view that the eggs or young of the owner are thrown out by the young parasite, and not removed by its parent, though I could find no trace of them beneath the nest.

Twenty-two eggs of *M. aeneus* average  $.90 \times .70$ , the extremes being  $.95 \times .75$  and  $.82 \times .65$ . The color is a *greenish-white, unspotted, soon fading to a dull opaque white*. There is more than the usual variation in shape. Some are almost perfectly elliptical, others are nearly round ; some are quite pointed at the smaller end, while others still are there abruptly truncate.

The young, soon after leaving the nest, have the plumage uniform dull black ; cheeks and sides of head bare ; iris brown.

*Fort Brown, Texas.*

## ON SEVENTY-FIVE DOUBTFUL WEST-COAST BIRDS.

BY J. G. COOPER, M. D.

IN Ornithological works written previous to 1860 we find many species of birds mentioned as from "California" and the "Northwest Coast," which have not been confirmed as from that region, and are therefore not referred to in more recent works. It is an interesting question whether they were always wrongly allocated there, or whether they may not, in many cases, have actually occurred as stragglers, and deserve, therefore, a place in our Fauna. As an aid in deciding these questions, I have compiled a list of all I can find, with such remarks as seem required respecting the chances of their occurrence, based on my own experience as a collector there for more than twenty years, and the observations of others on their usual range of distribution.

In 1852 Professor Baird published in Stansbury's Report of his Exploration of the Great Salt Lake, p. 327, a list of such species as were then known, including all given by authors as from west of the Mississippi, but not figured by Audubon. The one hundred and fifty-three nominal species included ninety-one from the Pacific slope, of which only twenty have not been since confirmed as belonging to our Fauna. Most of these were referred to again in his "Birds of North America." Mr. Cassin undertook to figure and describe "all" of these in his "Illustrations," but was undoubtedly saved from the repetition of many errors by the extensive collections soon after made by the Pacific Railroad Expeditions, although he has introduced several not since found in the United States.

In Volume XII, Part II, p. 288, of the Pacific Railroad Reports ("Natural History of Washington Territory") Dr. Suckley and myself, in 1859-60, printed a hastily prepared list of birds, not confirmed by us as from the "Northwest Coast," most of which, however, do not require to be excluded at present, only twenty out of one hundred and twenty-three coming into this list. I have carefully reviewed every accessible authority, and included only such as are distinct species and not represented within the regions named by geographical races or near analogues, which might

reasonably be mistaken for them. The older authors in confounding the races were quite excusable in giving them the same specific names. Those now to be noticed come under three divisions as to geographical distribution, viz.: first, those of tropical and South America; second, those of Asia; third, those of the Eastern United States. All circumpolar species, even if not recently found on this coast, are omitted as very likely to be found. The probability of the occurrence of the others varies chiefly with the nearness of their usual range and their powers of flight, so that we may frequently consider their occurrence as next to impossible, as with the Mexican Quails and Jays which have been so often credited to "California."

What we know of the Asiatic stragglers to Alaska shows that they are usually strong flyers, and I may here mention two of the Raptores found only in California so far, and each but once, which have a very Asiatic aspect, viz. *Buteo cooperi* and *Onychotes gruberi*.

The first doubtful West-coast birds were given by Gmelin as from "Unalaska, Nootka," etc.; the next came from "Oregon"; and not until 1830 did species begin to be credited to "California." At that time the peninsula was included in this name, and yet very few species, if any, were wrongly located on that account, most of those not found in "Upper California" being now unknown on the peninsula, though often found on the West Mexican coast. The cool sea-breezes of the coast act as an obstacle to the northward migration of many tropical species occurring on the Atlantic coast in summer much farther north, while the Colorado Desert, over one hundred miles wide, debars many more from the hotter interior.

A few have been called "Californian" which occur about the Gulf, but not within sixty miles of our line, though of course the water-birds may follow so far up the Colorado River. Many were no doubt carelessly labelled by collectors, or designedly misrepresented as from regions then almost inaccessible, to increase their value.

As to "Oregon" of authors before 1853, it must also be remarked, that they included under the name the whole country drained by the Columbia River; and even now the State includes part of the "Middle Province" of Baird, in which are found birds not known from the coast. Captain Bendire is now for the first

time collecting thoroughly the birds of Eastern Oregon; and his published notes, as well as those of Allen and Henshaw for Utah, of Ridgway for Nevada, and Coues, Henshaw, and others for Arizona, show very clearly what species distinguish the Middle and Western Provinces. Although about fifty species of Land Birds are given in the first volume of "Ornithology of California," which belong properly to the Middle Province or to Lower California, their habitats are so carefully described there that it is not necessary to include the species here, except in a very few striking cases.

As shown by Professor Baird from the Xantus collection made at Cape Saint Lucas, the Middle Province birds become common on the coast and peninsula south of latitude thirty-five degrees, to the exclusion of most of the characteristic Californian species, while very few of those of tropical Mexico occur on the peninsula, so that the chances are largely against the occurrence of the latter within our limits. Assistance derived from original observations and investigations by scientific friends, often unpublished before, is acknowledged by giving their names as authority.

1. **Turdus flavirostris**, *Swains.*, 1827, = *T. rufopalliatu*s, *Lafres.*, 1840, "Monterey, Cal." An abundant West Mexican bird, which may straggle northward with flocks of *T. migratorius*, which it closely resembles in habits.

2. **Harporhynchus rufus** var. **longirostris** (*Lafres.*), 1838. "California and Mexico." The occurrence of this Eastern Mexican form is not improbable, and it may have been the bird seen by me at Clear Lake, Cal., as recorded in History of N. A. Birds, III, 500.

3. **Ægithina leucoptera**, *Vieill.*, 1807, "North America," = ? *Motacilla leucoptera*, *Vig.*, 1839, "Western North America," Baird, List, 1852 (not of Quoy and Gaimard, which is a Palearctic bird). If = *Sylvia leucoptera*, *Wilson*, Index, it is *Dendroica carulescens*, not known far west of the Mississippi (*Coues*). [*Ægithina leucoptera*, *Vieill.*, according to Gray (Hand-List), is from India, while *M. leucoptera*, *Vigors*, is from Persia (*Lawrence*).]

4. **Sialia sialis** (*Lin.*). "Columbia River," *Aud. Syn.*, 1839 (error?). Not mentioned from there by Townsend nor Nuttall, who were then the chief authorities. Still it very probably will occur west of the Rocky Mountains. Some specimens of *S. mexicana* are stated to approach very near it, from which Audubon's statement may have arisen.

5. **Parus carolinensis**, *Aud.* "Oregon," Nuttall, 1840, by error for *P. atricapillus* var. *occidentalis*, which is very near it.

6. **Geothlypis velatus** (Vieill.) = *Trichas delafeldi*, Aud., 1839, Nutt., 1840, Heermann, 1858. "Oregon and California." It seems strange that a bird so common as *G. trichas* is in the East should be confounded with such a different South American species. Townsend's specimen was doubtless from South America, but Heermann's were only *G. trichas*. Lately recorded from Mazatlan, Mexico, and may therefore reach the Lower Colorado.

7. **Siurus auricapillus** (Linn.). "Oregon," Nuttall, 1840, and recently recorded from Idaho, as well as Alaska and Mazatlan. "California," Bonap., Notes Delat., 1853, but may have been obtained in Nicaragua. Still it very probably migrates occasionally through California.

8. **Dendrocæa montana** (Wils.). "California," Aud., 1839 - 41, Nutt., 1840. There is no further information respecting this very rare and scarcely known bird than is given by the above-named authors.

9. **Dendrocæa striata** (Forst.). "Oregon," Aud. Syn., 1839. Not given by Townsend nor Nuttall, though it should pass west of the Rocky Mountains in going from Alaska to the tropics. Yet it is not recorded as from the Rocky Mountains or Mexico.

10. **Dendrocæa cærulea** (Wils.) = *azurea*, Steph. "Oregon," Townsend's List, 1839, Aud., 1839, Nutt., 1840. Recently found west of the Rocky Mountains, so that it may not have been confounded by Townsend, etc., with *Polioptila*, as I suspected. It has not been detected near the coast.

11. **Setophaga ruticilla** (Linn.). "California," Bonap., Notes Delat., 1853, but was perhaps from Nicaragua. Still, as it is common throughout the Rocky Mountains in summer, it may occur in California, though not known from Western Mexico.

12. **Collocalia? unalascenkensis** (Gmel.). According to Cassin (Illustr., 1855, 251), Gmelin's bird belongs to the genus *Collocalia*, which is peculiar to the Pacific (tropical) islands, Japan, etc. If really found on the Aleutian Islands, some may be expected to straggle to our coast in winter, with other Asiatic species found there by Dall.

13. **Hypocolius ampelinus**, Bonap., 1850 (Consp.). "California." This species is now known to be from Sennaar, Upper Egypt, and is undoubtedly to be removed from the list of West-coast birds.

14. **Lanius lahtora**, Sykes, = *L. elegans*, Swains., 1831, Nutt., 1840. Supposed to have been received from Northwest America, but it inhabits Siberia. Stragglers may, however, cross Behring's Straits, like several other birds. "*L. elegans*" of other American authors is *Collurio ludovicianus* (L.) var. *robustus*, Baird, 1873.

15. **Vireo agilis**, Licht., 1823, = *V. virescens*, Cass., not of Vieill., which is *V. bartrami*, Swains., 1831; not of Aud., 1839, nor Nuttall, 1840. This species, confounded by former authors with *V. gilvus* var. *swainsoni*, has not been found north of Mexico, nor perhaps in North America, unless Douglas really found it at the Columbia River, as supposed by Swainson.

16. **Coccothraustes ferreirostris**, *Vigors*, 1839, = *papa*, *Kittl.*, 1830. "California," *Baird*, 1852.\* [This is now recognized as the *papa* of *Kittlitz* from the Bonin Islands (*Allen*).]

17. **Pyrrhula inornata**, *Vigors*, 1839. "Northwest coast of America," *Baird*, *List*, 1852. Probably Asiatic; certainly not North American.

18. **Ramphopsis flammigerus**, *Jard.*, about 1830, "Columbia River." [Now known to be South American, — *Columbia*, *Sclater* and *Salvin*; *New Grenada*, *Gray* (*Allen*).]

19. **Chrysopoga typica**, *Bonap.*, 1850. "California." [A Mexican and Central American species, not yet confirmed as from California (*Allen*).]

20. "**Emberiza**" **atricapilla**, *Gmel.*, 1788, *Aud.*, 1839–41. These authors confound this Sandwich Island bird with the *Zonotrichia coronata*, *Pall.*, a common California bird described in 1831.

21. "**Fringilla maculata**," *Aud.*, 1839 (*Townsend's List*), was a confusing of *Hedymeles melanocephalus*, *Swains.*, with some foreign species, perhaps *Pipilo maculata*, *Swains.* It occurs only in *Audubon's plate*. †

22. **Pyrgisoma biarcuatus** (*Lafres.*), 1855, "California," is a Central American bird.

23. **Saltator rufiventris**, *Vigors*, 1839, "West coast of North America," *Baird*, *List*, 1852. [A Bolivian species (*Allen*).]

24. **Icterus baltimore** (*Linna.*), "Columbia River." *Aud. Syn.*, 1839. No authority is given, and it is not mentioned by *Townsend* or *Nuttall*. Like the *Sialia*, it can only be a straggler.

25. **Icterus pustulatus**, *Licht.*, "California," *Bonap.*, *Notes Delatt.*, 1853, probably for Nicaragua. Some of these species may be looked for along the Colorado River.

26. **Icterus "californicus"** (*Lesson*), 1844; "California," *Bonap.*, *Consp.*, 1850. [This is a synonym of *I. pustulatus* (*Lawrence*).]

27. **Icterus icterocephalus** (*Linna.*), *Bonap.*, 1825, *Nuttall*, 1832–40. This well-known South American bird was only at first confounded with our *Xanthocephalus*, and should not have been credited to California by *Nuttall*.

28. **Xanthornus mexicanus** (*Briss.*), "Pacific coast of (North?) America," *Baird*, *List*, 1852. [Probably *Gymnomystax melanicterus* (*Vieill.*) of tropical America (*Allen*).]

29. **Trupialis militaris** (*Linna.*), "Monterey, Cal.," *Neboux*, *Voy. de la Venus*, 1855; "San Francisco," *Cutts*, in *Baird's N. A. Birds*, 1858.

---

\* *Chrysomitris yarrelli* (*Aud.*) and *Hypocanthus stanleyi* (*Aud.*) are now generally believed to be (if not hybrid cage-birds) from South America, and as he gives no authority for "Upper California," they may be omitted.

† *Petrochelidon macroura*, *Lawr.*, is quoted from "California" by *Cassin* (*Ill.*, p. 229), but has not lately been found west of Arizona. *Calamospiza bicolor*, *Townsend*, before reported doubtfully from "California," has been found as a rare bird near Tulare Lake, by *Mr. W. A. Cooper*.



Although not recently obtained in North America, it has thus been twice reported from California.

30. **Quiscalus major**, Vieill. "California," Gambel, 1847, but he obtained it only at Mazatlan. It may be looked for on the Lower Colorado.

31. **Quiscalus purpureus** (Bartr.). "Oregon," Aud., 1839, without authority. "California," Gambel, 1847, Newberry, 1857, who probably both mistook *Scolecophagus cyanocephalus* for it; no specimens are known from the west slope. *S. ferrugineus* (Gm.), "Oregon," Townsend, was obtained there by Peale, and is common in Alaska.

32. **Corvus ossifragus**, Wils. "Oregon," Townsend's List, Aud. Syn., 1839; "California," Woodhouse, 1853; all mistaking *C. americanus* var. *caurinus* for it.

33. **Cyanocitta beecheyi** (Vigors), 1829. "California," Botta in Eydoux's Voy. de la Favorite, 1839, but not known from the peninsula recently, or found north of Mexico.

34. **Cyanocitta ultramarina** (Temm.). Audubon (1839-40) and Nuttall (1840) confounded this Mexican species with *C. californica* (Vig.). The var. *arizonæ*, Ridgw., may reach California at the Lower Colorado River.

35. **Calocitta colliei** (Vigor), 1829, = *Pica bullocki*, Aud., 1831-42 (not of Wagler), Nuttall, 1840, both of whom described it as from the "Columbia River," but without good authority, while Nuttall denies ever having seen it there or in California. It probably does not even straggle north of Mazatlan, Mexico.

36. **Cyanocorax geoffroyi**, Bonap., 1850. "California." [This is a synonym of *Cyanocitta beecheyi* (Lawrence).]

37. **Sayornis fuscus** (Gmel.). "Oregon," Townsend's List, 1839, but it is not now known west of long. 100°. He may have mistaken *S. nigricans* for it, as that reaches Southern Oregon.\*

38. **Antrostomus? macromystax** (Wagl.), Cassin, p. 240. "California," from a label in Mus. Phil. Acad. A well-known Mexican species.

39. **Antrostomus nigrescens?** Cab., = *A. californianus*, Bonap., 1850, New Grenada. I think some of the larger tropical species of this family may stray into California, as I saw what appeared to be one as large as *A. carolinensis* in Ventura County in 1872, but could not obtain it, and heard no note.

40. **Picus lineatus**, Linn. "Oregon," Aud., 1839-41, from a specimen in Edinburgh "sent by Dr. Gairdner." Not known from North America, and was probably collected in South America.

---

\* *Saurophagus bairdi*, Gamb., 1847, has been attributed to California, but was given by the author as from the Gulf Region of Mexico. It is South American, and has not recently been reported from Mexico.

41. **Dryotomus delattri**, Bonap., 1854. Doubtless labelled "California" by error for Nicaragua.

42. **Campephilus imperialis** (Gould), 1832. "California," Aud., 1839, Nutt., 1840, Cassin (figured), 1855. Although Nuttall states that Townsend shot a specimen in the Rocky Mountains, near the head of the Colorado River, it is not mentioned in Townsend's List, nor figured by Audubon. Townsend, however, states that he shot, but lost, a Woodpecker resembling this on Malade River (now in Idaho). Cassin gives reasons for supposing it may also occur in California, and it is to be looked for in Arizona.

43. **Melanerpes erythrocephalus** (Linn.). "California," Gambel, 1847, Baird in Ornith. of Cal., 1870. Dr. Gambel does not seem to have preserved any specimens, and as no later collectors have found it west of Salt Lake City ("one seen," Ridgway), its occurrence in California must be merely accidental.

44. **Ceryle americana** (Gmel.). "Colorado River," Coues, 1866. As Dr. Coues only thought he saw this species along the river, and as such a large bird could scarcely escape the many collectors who have been at Fort Yuma and along the Gila River (whose clear waters are better suited for it than the muddy Colorado), we may doubt its occurrence until specimens are obtained, its range in Texas not being north of lat. 30°.

45. **Haliaeetus pelagicus** (Pull.), 1831. "Aleutian Islands." Although not obtained by late collectors in that region, its occurrence as a frequent visitor from Kamtschatka is more probable even than that of *H. albicilla* to Greenland, and it may be looked for at least as far south as lat. 50° on our coast.

46. **Syrnium nebulosum** (Forst.). "California," Woodhouse, 1853. The birds seen by him in Arizona, also, were probably *S. occidentale*, Xantus, 1859. See Baird in Orn. of Cal., 1870, p. 431. There are several instances besides this in which southern species of Owls do not extend across the continent, although those of the arctic regions, being mostly circumpolar, are common to both sides. Thus *Nyctale tengmalmi* might have been included in the Orn. of Cal. instead of this, it having been found in Oregon by Townsend.

47. **Catharista atrata** (Bartr.). "Columbia River," Douglass in Fann. Bor. Am., 1831, Aud., 1839, Peale, 1848, Cassin, Ill., 1853, 1858. It is possible that these references were to the young of *Rhinogryphus aura*, although an actual specimen seems to be alluded to. Dr. Gambel found it quite common about the Gulf of California, but does not add California as in other cases, nor has it been detected, though very likely to be, along the Colorado.

48. **Sarcorhamphus gryphus** (Linn.). "Southwestern States," Bonap., 1828 - 33; "Rocky Mountains," 1832 - 40, quoting Lewis and Clarke's "bustards," and the bill and talons brought by them to Peale's Museum. These were, however, probably those of *Pseudogryphus califor-*

*nianus*, and from the Lower Columbia River. The Condor is doubtless confined to South America, though quite able to visit our latitude.

49. **Melopelia leucoptera** (*Lin.*). This species, common in Arizona, has not been collected in California, but in 1853 I saw white-winged wild Pigeons, which I then had no doubt were this species, in the San Francisco market. As they are often caged in Mexico, these may have come from there, but it is also very probable that they may wander so far, like the little Ground Dove, which has been shot at San Francisco.

50. **Ortyx fasciatus**, *Natterer* MSS. "California," Gould, 1843, but it is not now known north of Colima, Mex., and not at all likely to be found north of that point.

51. **Lophortyx elegans** (*Lesson*), 1831. "Upper California," Nuttall, 1840, who, however, did not see it. Found at Mazatlan, but not much to the northward. *L. douglassi*, *Vig.*, is probably young of *L. californicus*.

52. **Eupsychortyx cristatus** (*Lin.*) = *O. neoxenus*, *Vig.*, 1830, Aud., 1839 - 42, Nutt., 1840. "Northwest coast of America," Beechey; "California," Audubon. This species is not given in any recent lists of birds of western tropical America. Beechey's specimen may have been a cage-bird, or obtained in Costa Rica.

53. **Gambetta flavipes** (*Gmel.*). "Oregon," Townsend, 1839; "California," Woodhouse, 1853, Newberry, 1857. Although some undoubtedly occur for a considerable distance west of the Rocky Mountains and in Alaska, it is a curious fact that no specimen seems to have been collected in California, Nevada, or Arizona, nor have I seen it, while the larger species is abundant.

54. **Hæmatopus ater**, *Vieill.*, = *H. townsendi*, Aud., 1839. "Oregon," Aud., not Townsend. Doubtless collected in South America, but, like other shore birds, may be also more or less common to the coast of North America.

55. **Numenius rufiventris**, *Vig.*, 1828. "Pacific coast of North America." The name would apply well to a common variety of *N. longirostris*; but it is not yet quoted as a synonym of that species, though there seems to be no other species on the coast to which it is referable.\*

56. **Grus americana**, *Forst.* "Oregon and California," Townsend and Audubon, 1839. The error arose from confounding *G. canadensis* with this, which is not now known to go west of long. 100°.

57. **Audubonia occidentalis** (*Aud.*). "California," Gambel, 1847, "to Columbia River"; Newberry, 1857. No specimens were obtained, and they no doubt mistook the large var. "*californica*" of *Arlea egretta* for it.

58. **Platalea ajaja**, *Lin.* = "*P. mexicana*,"? Willoughby. "California to San Francisco," Gambel, 1847. Not seen since then north of the Gulf of

---

\* *Actodromus "bairdii"*, Cassin, 1858, not of Schlegel, was confounded by him with *A. bairdi*, Coes, 1861, and is still doubtful as a West-coast bird, as is the more northern and Alaskan *Actiturus bartramius* (*Wils.*).

California, and not very likely to straggle so far north, although *Tantalus loculator* visits San Francisco Bay and vicinity every year. Could he have been deceived by hearing *Spatula clypeata* called "Spoon-bill"?

59. *Anser segetum* (Gmel.). "Probably on Northwest coast in winter," Nuttall, 1832. The evidence for the occurrence of this species anywhere in North America is very slight, although it is quoted by Swainson and Richardson from Hearne. [Doubtless *A. canadensis* (Coues).]

60. *Anas obscura*, Gmel. "Oregon," Aud., 1839, Townsend's List, "California," Woodhouse, 1853. This, probably a melanistic form of the Mallard, has not recently been found west of Utah. (See Henshaw, Rept., 1875.)

61. *Dafila urophasiana* (Vig.), 1829. "Northwest coast." This South American species has not been confirmed as from North America, and, if found, occurs only as a straggler.

62. *Larus belcheri*, Vig., 1829. "Pacific coast of North America." Though confounded by some with *L. heermanni*, this species is probably limited to South America, and the range of the two species does not meet.

63. *Chrococephalus atricilla* (Linn.). "Colorado River," Coues, 1868, but not yet obtained on Pacific side north of Cape Saint Lucas.

64. *Sterna antillarum* (Less.). In exactly the same category as the last.

65. *Hydrochelidon nigrum* (Linn.). "Oregon," Towns., 1839; "California," Heermann, 1858. According to Coues (Mon. Laridæ), the common American bird is *H. lariformis* (Linn.), so that the *H. nigrum* is not likely to occur, except as a straggler, in the West, where all I have seen were the common kind. [This is probably what Townsend and Heermann meant by "*nigrum*" (Coues). Saunders (Proc. Zool. Soc. 1876, p. 642) makes them identical (Lawrence).]

66. *Xema furcatum* (Neboze), 1840. "Coast of California." As this remarkable species has not been confirmed from the West coast, the locality may well be doubted, like too many recorded by the same author. [Recently stated by Salvin to be an inhabitant of the Gallapagos (Lawrence).]

67. *Graculus carbo* (Linn.). "Nootka Sound," Nutt., 1834, "Oregon," Towns., 1839. Not confirmed by collections from the West coast, and no doubt confounded with other species.

68. *Sula bassana* (Linn.). "Northwest coast of America," Nutt., 1834.

69. *Sula fiber* (Linn.) = *fusca*, Vieill. "Off coast of California from San Francisco south," Newberry, 1857. The species seen were probably *S. piscator* (Linn.), and *S. cyanops*, Sund., which have been obtained from the West Mexican coast, but not farther north, and I have not seen them along the coast north of lat. 30° during several voyages.

70. *Diomedea chlororhyncha* (Gmel.). "Off the coast of Oregon,"

Towns., 1839, Aud., 1839-44. Audubon's figured type proves to be *D. culminata*, Gould, and was probably obtained too far from our coast to be included in its Avifauna, as none have been found lately alongshore or in sight of land. Townsend does not include it in his list of Oregon birds.

71. *Diomedea fuliginosa*, Gmel., = *D. fusca*, Aud., 1839-44. Included in Townsend's List, but not recently confirmed, and as he does not mention *D. nigripes*, Aud., he may have referred to the type of that species. The South Pacific *D. exulans* and *Daption capensis* should also be excluded.

72. *Æstrelata hæsitata* (Kuhl.). "California," Lawrence, 1853, by error for *Proifinus cinereus* (Gmel.). No record of the former from the Pacific.

73. *Puffinus obscurus* (Gmel.). "Northwest coast of America," Nuttall, 1834. No late record of its occurrence in the Pacific.

74. *Podiceps minor* (Gmel.). "Oregon," Townsend, 1839. Given by Nuttall as North American, but not lately obtained, being a common European species, and confounded by Townsend with either *P. cornutus* or *P. auritus*.

75. *Podiceps dominicus* (Linn.). "California," Gambel, 1847. Probably not obtained north of the Gulf, and not confirmed as living north of lat. 32°.

---

#### REMARKS ON *SELASPHORUS ALLENI*, HENSHAW.

BY D. G. ELLIOT.

I HAD commenced an article in reference to the two forms of the *Selasphorus rufus* of authors, as observed in California and Mexico, when the July number of the Bulletin of the Nuttall Ornithological Club reached me, containing an interesting paper by Mr. Henshaw, on the California bird, which he describes as new under the name of *S. alleni*. That there are two well-defined *species*, as the term is usually understood nowadays, I have for a long time been well satisfied in my own mind, and the peculiar shape of the lateral rectrices would seem to be sufficient to establish the specific differences of the two birds. Mr. Henshaw has done good service in pointing these out; but unfortunately he has conferred a new name upon the wrong bird, for it is the southern form that requires to be designated, and not the northern, or to be perhaps more exact, it is the red-backed bird with the broad tail-feathers, and not the green-backed one with the narrow tail-feathers, that is in need of a name.

To prove this it would be necessary to go back in the history of the species, and commence at the beginning. Gmelin seems to be the first author who conferred a Latin name upon the Rufous-throated Hummer, which he did in his "Systema Naturæ" (1788), Vol. I, p. 497, sp. 57, and described it as follows: "Trochilus rufus . . . rostrum pedesque nigri; colli pennæ laterales nonnullæ elongatæ mobiles; tectrices alarum obscure virescentes; rectrices splendide rufæ *acuminatæ*, linea media longitudinali et apice nigris; *cauda cuneata*." Moreover, he gives the habitat as "in sinu Americæ Natka," and quotes as his synonyms the Ruffed Honey-Sucker of Pennant's "Arctic Zoology," Vol. II, p. 290, No. 177, and the Ruff-necked Humming-Bird of Latham's "Synopsis," Vol. II, p. 785, No. 56, t. 35, whose specimen, as Latham informs us, came from Nootka Sound. Now, as it is well known that the southern bird with the broad rectrices has a wide dispersion, going far to the north on the Pacific coast, it might be said, "How are we to know that the specimen from Nootka Sound was not this species, and that it was the one called *alleni* by Mr. Henshaw?" Fortunately this can be satisfactorily determined, and all doubts removed, by turning to the "Fauna Boreali Americana" (Birds), and on page 324 we find that Swainson, in his article on the *Trochilus (Selasphorus) rufus* of Gmelin, makes the following statement: "The discovery of this superb species in the cold and inhospitable regions of Nootka Sound is due to our great navigator, Captain Cook, while to Dr. Latham belongs the honor of first making it known to science. *By a singular chance we have at this moment before us one of the identical specimens, in perfect preservation, collected by the naturalists of that expedition*; it was presented by the late Sir Joseph Banks to Mr. Bullock, and was purchased by us at a very high price at the dispersion of that collector's museum by public auction." In his description of the *form* of this bird, he says: "The *tail*, although short, is more cuneated than rounded, the two middle pairs being longest, *all are narrowed and obtusely pointed at their extremities, but the two outer pairs are particularly narrow*." It will thus be seen, I think, that the species described by Gmelin from Nootka Sound was, without doubt, the bird with narrow rectrices, as Swainson's specimen was a typical one, if indeed it may not have been the original type; and he was too keen a naturalist not to have noticed the peculiar notch in the rectrices next the median pair, observed in the bird with the broad tail-feathers. He also speaks of the throat as being equally

brilliant with that of *T. moschilus* (!) (which, so far as I have seen, is never the case with the other species), and it has, he says, more of a *red* than an *orange* gloss," and the tints are "exquisitely splendid"; a perfectly accurate description of the California bird, but not of the other, which has the gorget *orange*, and not at all brilliant. To come a little nearer to our own time, we have Audubon, who, in his "Birds of America" (8vo edition, Vol. IV, p. 202), thus describes the *Selasphorus rufus* as he knew it from the specimens collected on the Blue Mountains of the Columbia River and at Nootka Sound by Messrs. Nuttall and Townsend: "Tail rather long, broad, graduated, the lateral feathers four and a half twelfths of an inch shorter than the central; the latter are extremely broad, measuring four and a half twelfths across, and the rest gradually diminish to the lateral, which are *very narrow*, all obtusely pointed." Not a word, it will be noticed, is said of the notch on the first rectrices from the central ones. The throat is also stated to be "splendent fire-red," etc. Baird, in the "Birds of North America" (1860), p. 134, in his description of the *S. rufus*, says that "the tail is strongly cuneate; the outer feather .40 of an inch shorter than the middle, which projects .14 of an inch beyond the rest. *The outer feather is very narrow, not exceeding .11 of an inch in width*; the rest widen and lengthen rapidly to the central one, which is very broad (.35 of an inch); the central feathers are all ovate-acuminate. The entire throat, including a short ruff on the side of the neck (about .40 of an inch long), is *metallic red*, of the same shade as in the Ruby-throat, although with brassy reflections in some lights." Gould, in his "Monograph of the *Trochilidae*," has apparently confused the two species together, but he makes no mention of the *notched* rectrices, but states they are all of a "broad lanceolate form," and his figures would seem to be taken from the California bird. I might go on and multiply the instances where writers in their descriptions of *S. rufus* have spoken only of the birds with the narrow rectrices, although, as in Mr. Gould's case, they may have had both Californian and Mexican specimens before them, but, regarding them as one species, they have always selected for their descriptions the specimens with the brilliant throats (as being in more perfect plumage, as they supposed), rather than the duller-throated examples, and so these last have escaped receiving a distinctive name, as they deserved. But I think enough has been said to show that authors generally, and the older ones especially,

always described the bird with all the lateral rectrices narrow, and destitute of any notch.

I have not said anything, in comparing the species, about the color of the back, as I consider this is not altogether a satisfactory character by which to distinguish the birds, though Mr. Henshaw makes it one of his principal ones. Latham, in his original description, states that the "crown" was "glossy green," and also that there was a "greenish gloss between the wings." In the first place, the females of both are entirely green in their upper surface, and this is not always pure green, as I have specimens now before me, collected by Boucard at Oaxaca, Mexico, in which the back is a yellowish-bronze, precisely like young males in my collection from California of the other form, collected by Dr. Heermann. Again, I have young males, also collected by Boucard at Oaxaca, which have the back of such a curious reddish-cinnamon that it is difficult to say what color it exactly is; and Mr. Henshaw says, in his article (p. 55), that "in California, however, where the *S. rufus* occurs in its typical condition, that is, with an unmixed rufous back, specimens are not uncommonly found which exhibit a strong approach to the coloration of *S. alleni*"; although, as he farther says, "they never appear to pass beyond a certain point." It is, however, indisputable, that the two species do vary in the amount of green upon their upper surface, and also that at times they approach each other in coloration so nearly that, were there no other differences existing, it would be impossible to separate them. For this reason I do not place much reliance upon the amount of green on the back as a *specific character*. But there are other differences, I think, not mentioned by Mr. Henshaw, to be observed in the females, by which this sex of the two species can be distinguished. The female of the Mexican\* species has the rectrices broad.

In addition to the superior width of its rectrices, the Mexican bird has the lateral tail-feathers, for more than a third of their length in the central portion, jet black, the base light rufous, and the tips white, so that when the tail is closed, and looked at from

---

\* I use the terms *Mexican* and *Californian* to designate the birds with broad and narrow rectrices respectively, for the term *rufous* has been so misapplied that I cannot employ it at present without risk of adding to the confusion. At the same time the bird called *alleni* is not restricted to California, as I have already shown, but goes as far north at least as Nootka Sound, and may in winter pass into Lower California, perhaps into Mexico.



beneath, it appears all black, tipped with white. In the other species the rufous of the basal portion is more extended, and the blackish bar narrower, and does not occupy all the space between the tips of the under-coverts and the white tips, at least on the lateral feathers. I do not think that the females have any metallic feathers on the throat; those mentioned as females, with these feathers, being usually young males. These last generally have the median rectrices cinnamon, the tips only being metallic green. As, therefore, it will be necessary to bestow a new name upon the bird with the metallic-orange throat and notched rectrices next to the central pair, the synonymy of the species will stand somewhat as below.

The descriptions are taken from specimens in my collection from California and Mexico respectively.

#### **Selasphorus rufus.**

*Trochilus rufus*, GMEL., Syst. Nat., Vol. I, p. 497. (1788.)

*Ruff-necked Humming-Bird*, LATH., Gen. Syn., Vol. II, p. 785, pl. 35. (1781.)

*Trochilus (Selasphorus) rufus*, SWAINS., Faun. Bor. Amer., Vol. II, p. 324. (1831.)

*Selasphorus rufus*, AUD., B. Amer., Svo ed. Vol. IV, p. 200. BAIRD, B. Amer. (1860), p. 134. GOULD, Mon. Troch., Vol. III, pl. 137 (partim).


*Schusphorus ruber*, BOX., Consp. Gen. Av., p. 82.

*Oreismya soisii*, LESS., Hist. Nat. Ois. Monch., p. 190, pls. 66, 67.

*Selasphorus allenii*, HENSHAW, Bull. Nutt. Ornith. Club, Vol. II, p. 54 (1877).

*Habitat.* In summer the Pacific coast of America from California to Nootka Sound. In winter — ?

*Male.* Top of head and back bronzy-green, dullest on the forehead. Sides of the head, rump, flank, abdomen, and under tail-coverts rufous. A gorget of metallic feathers, covering all



the throat and extending on to the sides of the neck, brilliant coppery-red, with brassy reflections in certain lights. Upper part of breast white. Wings purplish-brown. Tail short, cuneate,

---

\* Figures reprinted from Mr. Henshaw's article (this volume, p. 53), with change of names.

all the feathers acutely pointed. Median rectrices rather broad, lateral ones narrowing rapidly to the outermost, which is extremely narrow. Bill straight, black.

*Female.* Above entirely green, with a slight cinnamon shade on the rump. Under surface grayish-white, with a slight brownish tinge on the breast. Tail-feathers rufous at base, then a narrow subterminal bar of violaceous-black, and tipped with white.

The next species I propose to call

**Selasphorus henshawi.**

*Trochilus rufus*, HENSHAW, Bull. Nutt. Ornith. Club, Vol. II, p. 53 (1877).

*Habitat.* Mexico, northward along the Pacific coast to Sitka.

*Male.* Top of the head metallic-green, rest of upper parts cinnamon, but some specimens have green feathers intermixed with the rufous on the back. Throat metallic-orange, not brilliant as in the other species. Breast, and the centre of the abdomen, white; flanks and under tail-coverts rufous. Tail rufous, tipped with dark brown; feathers pointed at tip, median pair broad, lateral ones growing narrower to the outermost, which is the most attenuated. *On the inner web near the tip of the rectrices next the central pair is a conspicuous well-developed notch.* Bill black. Total length,  $3\frac{3}{4}$  inches; wing,  $1\frac{1}{2}$ ; tail,  $1\frac{3}{8}$ ; culmen,  $\frac{5}{8}$ .

*Female.* Entire upper parts shining grass-green, dullest on the crown. Throat white, spotted with brown. Under parts white; washed with rufous on the breast and flanks. Under tail-coverts buff. Median rectrices green; lateral ones rufous at base, then a band of metallic-green, succeeded by a subterminal broad black bar, and tips white. Bill black. Length,  $3\frac{1}{2}$  inches; wing,  $1\frac{7}{8}$ ; tail,  $1\frac{1}{2}$ ; culmen,  $\frac{5}{8}$ . Young males similar to the females, with a few metallic spots on the throat.

---

THE YELLOW-THROATED WARBLER (*DENDRÆCA*  
*DOMINICA*).

BY WILLIAM BREWSTER.

It is indeed surprising that a bird so generally distributed throughout the Southern States as the above-named species should be so little known. In "History of North American Birds" (Vol. I, p. 241), Dr. Brewer prefaces his account of its habits by the remark that its history "is very imperfectly known," and then proceeds to draw upon the meagre and conflicting descriptions given by Wilson, Audubon, and Nuttall. Although I cannot myself claim an ac-

quaintance of very long standing with this beautiful little species, still for five or six weeks during the past spring scarcely a day passed that I did not see one or more individuals. I first met with them at Mellonville, Florida, where, on March 14, I shot two specimens, both females, in the pine woods near the town. They were associated with Pine Warblers, Nuthatches, and Woodpeckers. During a trip up the Wekiva River, March 19 to 23 inclusive, I heard at frequent intervals a Warbler that I did not recognize singing in the cypresses, but from the impenetrable nature of the swamps, and the great height of the trees, I was unable to get even a glimpse of the bird. A week later, while descending the St. John's River by steamer, I again constantly heard, both from the cypress swamps and the open piny woods, the notes of this, to me, unknown species, and although I felt almost certain of its identity, it was not until I reached St. Mary's, Georgia, that I proved to my satisfaction that my suspicions were correct. There, from the 6th of April to the 4th of May, I enjoyed abundant opportunities of studying its habits, for it was everywhere, in suitable localities, if not one of the most abundant, at least a generally distributed species. At the time of my arrival the males were in full song and mating. A few individuals haunted the moss-hung live-oaks that shaded the village streets, but the open piny woods were their favorite abode. There, with the Summer Redbird (*Pyrranga astiva*), the Pine Warbler (*Dendroica pinus*), the Brown-headed Nuthatch (*Sitta pusilla*), and a variety of Woodpeckers, they frequented the beautiful Southern pines. Indeed, so great was their attachment to this tree that, with the exception of those heard in the cypress swamps of the Upper St. John's, and the few that inhabited the oaks in the town, I do not remember to have seen one in any other tree. So marked and unvarying was this preference, that on more than one occasion I made use of the notes of this bird to guide me out of some bewildering thicket, feeling sure that beyond where it was singing I should find the more open pine-clad country.

Nearly all the authors who have written on the Yellow-throated Warbler from personal observation compare his movements along the branches to those of the Black-and-white Creeper (*Mniotilta varia*). At first I was inclined to the same opinion, but after my eagerness to secure specimens had somewhat abated, through success in collecting them, I felt more at leisure to watch the pretty little birds before taking their innocent lives, and, having spent many hours in

carefully studying their habits, I became convinced of the error of my earlier impressions. Their movements are much slower than those of the *Mniotilta*, and there is less of that crouching, creeping motion. They do, indeed, spend much of their time searching the larger branches for food, but it is much more in the manner of the Pine Warbler, and their motion is rather a hopping than a creeping one. I have never seen them ascend the trees from the roots to the top-most branches, as Audubon relates, but I occasionally observed one clinging against the main trunk for a moment, to seize an insect, as will the Bluebird (*Sialia sialis*) and many of the Warblers. Their hunting-ground is for the most part, however, among the higher branches, and a considerable part of their time is spent at the extremities of the limbs, searching for food among the pine needles. Their bright yellow throats, brought out by contrast with the dark evergreen foliage, give them a certain resemblance to the Blackburnian Warbler (*Dendroica blackburniae*). The males are not very persistent singers. I rarely heard them during the warm hours of the day, even when pairing was almost their sole occupation. Their song is very pretty; it may be nearly imitated by the syllables *Twsee-twsee-twsee*, *twsee-see*, the last two rising and terminating abruptly. It most nearly resembles that of the Nashville Warbler (*Helminthophaga ruficapilla*), beginning in almost the same way, but ending differently, and, indeed, throughout the notes are much sweeter. Both sexes utter a chirp similar to that of other Warblers, but sharper.

By the middle of April there was a marked decrease in the number of Yellow-throated Warblers about St. Mary's. This was partly owing to my having shot many for specimens, but not entirely to this, for extended researches over new ground convinced me that the greater number had passed on, probably to the northward. A few, however, still remained; perhaps on an average one pair to every hundred acres of pine forest. While collecting near St. Mary's, April 18, I was in the act of shooting a female when I noticed that she was gathering material for building, and, tracing her flight, I was fortunate enough to discover her half-completed nest. Visiting the spot at frequent intervals, I invariably found both birds feeding among the pines in the vicinity, although the nest, as far as I could judge, seemed finished. At length, May 2, a friend, ascending the tree, found the female sitting. She remained on the nest until he nearly touched it, although the limb shook

violently under his weight. When she did finally leave it she sailed down into a smaller tree a few rods off, where she remained a silent and seemingly unconcerned spectator of what followed. The nest and its contents being safely lowered to the ground, I shot both the female and her mate. The latter was singing, as usual, a short distance off, and apparently took no more interest than the female in the destruction of their mutual hopes. Embryos of small size had already formed in the eggs, so that incubation must have begun three or four days previously. This nest was placed at the height of about thirty-five feet from the ground, on the stout horizontal branch of a Southern pine, one of a thinly scattered grove or belt that stretched along the edge of a densely wooded hummock. It was set flatly on the limb, — not saddled to it, — nearly midway between the juncture with the main trunk and the extremity of the twigs, and was attached to the rough bark by silky fibres. It is composed externally of a few short twigs and strips of bark bound together by Spanish moss (*Tillandsia usneoides*) and a silky down from plants. The lining consists of a few hair-like filaments of moss and soft cottony vegetable down. The whole structure is neatly and firmly compacted, though essentially simple in appearance, and, from the nature of the component materials, of a grayish inconspicuous color. In size, shape, and general formation it very nearly resembles nests of the Black-throated Green Warbler (*Dendroica virens*) in my collection. It measures externally 2.80 inches in diameter by 1.70 in depth; internally, 1.77 inches in diameter by 1.30 in depth. The eggs, four in number, measure .69 by .53 of an inch. They are quite regularly ovate, with fine dottings of pale lilac scattered thinly and evenly over a grayish-white ground-color. A few spots or blotches of burnt sienna occur about the large ends, while occasional irregular penlike lines of dark brown diversify the remaining surface.

Upon referring to published accounts of the nesting of this Warbler, I find the statements by the different authors most conflicting, and the authenticity of many of the specimens open to gravest doubt. Both Mr. Maynard (in "Birds of Florida," Part II, page 61) and Dr. Coues ("Birds of the Northwest," page 67) base their descriptions upon alleged specimens sent to the Smithsonian Institution by Mr. Norwood C. Giles, of Wilmington, N. C. Dr. Brewer refers to these specimens as "eggs supposed to be of this species," and Dr. Coues describes the nest as "built in a large mass

of Spanish moss," and as "composed chiefly of that material. A part of the moss which hung from an oak bough, two feet downward and a foot across, was caught up and closely woven together with a little fibrous substance and much plant-down, to form a swinging bed for the nest, with a lateral entrance which will admit the hand. Inside is the nest proper, of the usual dimensions, very neatly wrought of the moss, with a smooth even border, and lined with plant-down and a few fine grasses." From the great difference in the position and structure of the nest, there seems little reason to doubt that Mr. Giles was mistaken in his identification. Nuttall's account of "its curious fabric, suspended to a kind of rope which hangs from tree to tree," is manifestly fabulous, while Audubon's description is, to say the least, very vague and unsatisfactory, though, as far as it goes, it certainly most nearly approximates to the specimen before me.

From the number of individuals I saw near Savannah, Ga., May 5 (I heard in Bonaventure Cemetery four different males singing at one time), I am led to believe that the Yellow-throated Warbler breeds more abundantly in Northern Georgia (and perhaps in South and North Carolina) than farther south.

---

### Recent Literature.

D'HAMONVILLE'S ENUMERATION OF THE BIRDS OF EUROPE.\*—The writer is indebted to his friend and correspondent, Baron D'Hamonville, for a copy of his recent Catalogue of the Birds of Europe, which deserves more than a mere mention on account of the admirably comprehensive manner in which it has been prepared. While it enumerates every form that has been claimed to have been taken within the limits of Europe, it is careful to designate in a significant manner the writer's mode of dissent. Thus, names believed to have no specific significance, or to represent a race rather than a species, are marked with a Greek minuscule; those whose presence is questioned are given with an interrogation sign, and those whose occurrence is exceptional are also designated. Thus divided, the Baron makes the whole number 658, as follows:—

---

\* Catalogue des Oiseaux d'Europe, ou énumération des espèces et races d'oiseaux dont la présence, soit habituelle soit fortuite, a été dûment constatée dans les limites géographiques de l'Europe, par J. C. L. T. D'Hamonville. Svo. pp. 74. Paris: 1876.

Species resident or of regular migration . . . . .	425
Species of fortuitous occurrence . . . . .	156
Species whose capture in Europe is disputed . . . . .	22
European races . . . . .	55
	658

The number of accidental visitors is surprisingly large, being about one fourth of the whole Catalogue, and is very nearly equally divided between Africa, Asia, and North America. These do not include those whose presence is disputed, though the number of the latter should probably be enlarged. Thus *Haliaeetus leucocephalus* is accepted without dispute, though its right to be included is generally questioned.

Appended is a list of the portion of the visitors to Europe from North America, with the region where taken : —

<i>Nauclerus furcatus</i> , England.	<i>Tringa bonapartei</i> , England and France.
<i>Accipiter atricapillus</i> , Scotland.	<i>Tringa pusilla</i> , England.
<i>Scops asio</i> , England.	<i>Tringites rufescens</i> , England, France, Heligoland.
<i>Picus villosus</i> , England.	<i>Gambetta flavipes</i> , England.
<i>Picus pubescens</i> , England.	<i>Rhyacophilus solitarius</i> , Scotland.
<i>Coccyzus americanus</i> , England.	<i>Actitis macularia</i> , Western Europe.
<i>Coccyzus erythrophthalmus</i> , Italy (Lucca).	<i>Bartramia longicauda</i> , Central Europe.
<i>Alcedo alcyon</i> , Ireland.	<i>Symphemia semipalmata</i> , Sweden, France.
<i>Agelæus phœniceus</i> , England.	<i>Botaurus lentiginosus</i> , Germany, England.
<i>Loxia leucoptera</i> , England.	<i>Puffinus obscurus</i> , France, England.
<i>Anthus ludovicianus</i> , Heligoland.	<i>Puffinus fuliginosus</i> , France, England.
<i>Turdus fuscescens</i> , Pomerania.	<i>Thalassidroma leucorhoa</i> , Western Europe.
<i>Turdus pallasi</i> , Switzerland.	<i>Chrocephalus atricula</i> , France, England.
<i>Turdus swainsoni</i> (not given).	<i>Chrocephalus philadelphia</i> , England.
<i>Harporhynchus rufus</i> , Heligoland.	<i>Xema sabinei</i> , Northern Europe.
<i>Regulus calendula</i> , England.	<i>Mareca americana</i> , England.
<i>Dendroica virens</i> , Heligoland.	<i>Querquedula discors</i> , France.
<i>Progne purpurea</i> , England.	<i>Aix sponsa</i> , England, France, Germany.
<i>Hirundo bicolor</i> , England.	<i>Fulix collaris</i> , England.
<i>Ectopistes migratorius</i> , Russia, Norway, England.	<i>Fulix affinis</i> , England.
<i>Charadrius virginicus</i> , Heligoland.	<i>Clangula albeola</i> , England.
<i>Charadrius fulvus</i> , Malta.	<i>Eidemia perspicillata</i> , Western Europe.
<i>Egialitis vocifera</i> , England.	<i>Mergus cucullatus</i> , France, England.
<i>Numenius hudsonicus</i> , Iceland, England.	<i>Uria columba</i> , Greece (Von Henglin).
<i>Numenius borealis</i> , Scotland.	<i>Simorhynchus psittaculus</i> , Sweden.
<i>Macrorhamphus griseus</i> , Northern Europe.	
<i>Tringa maculata</i> , England.	

T. M. B.

MERRIAM'S "REVIEW OF THE BIRDS OF CONNECTICUT."\* — Through

\* A Review of the Birds of Connecticut, with Remarks on their Habits. By C. Hart Merriam. Transactions of the Connecticut Academy, Vol. IV, pp. —, 1877.

the kindness of the author we have received advance sheets of the above-cited paper, shortly to be issued under the auspices of the Connecticut Academy, as the first of a proposed series on the Fauna of that State. Since the appearance of Linsley's "Catalogue of the Birds of Connecticut" in 1843, no detailed enumeration of the birds of that State has been published. Hence the advent of Mr. Merriam's paper must be hailed with interest by all engaged in the study of New England Ornithology. The author gives in all two hundred and ninety-two species. Of these he arranges under special lists in tabular form, one hundred and thirty-five as "summer residents"; twenty which probably breed occasionally, but are not known to do so; forty-one resident species; ninety migrants; sixty-seven winter visitants; thirty-one irregular summer visitants; forty-six rare accidental visitants; and nineteen rare and irregular visitants. Following these is a tabulated analysis of Linsley's Catalogue, in which he eliminates sixty-three species given by that author on apparently insufficient authority, though many of these are afterwards included by Mr. Merriam upon more recent and tangible evidence. The Avifauna of Connecticut, though essentially Alleghanian in character, has been long known to embrace many Carolinian forms, but the relative extent of this "tinge," geographically as well as specifically, has been considerably developed by Mr. Merriam's careful researches. Thus, upon unimpeachable authority are given as birds of Connecticut, *Dendroica dominica*, *Lophophanes bicolor*, *Oporornis formosus*, *Cardinalis virginianus*, *Empidonax acadicus*, and *Centurus carolinus*, while, singularly enough, several species known only in the more northern New England States as spring and fall migrants have been found breeding. In the careful elaboration of interesting details culled from personal experience and the note-books of well-known and trustworthy field collectors, this paper is most rich. Indeed, if we may be permitted to qualify otherwise undiluted praise by a little censure, we should say that a judicious summarizing of data and incident would divest this paper of a great deal of unnecessary cumberdom. Still, it is perhaps better to err in this direction than in the other and too common one, and Mr. Merriam certainly deserves much credit for his arduous labors. — W. B.

---

### General Notes.

NOTE ON *DORICHA ENICURA* (*Vicill.*). — About two years ago Mr. H. W. Henshaw submitted some birds to me for determination, among which was the female of a species of Humming-Bird obtained by him in Arizona, which I considered to be *Doricha enicura*, and it is so recorded in United States Geographical Survey W. of 100th Meridian, Vol. V, Chap. III. On a re-examination lately made, I find it was not properly referred, and a comparison with the female of *Calothorax lucifer* (Sw.) shows it to be that



species, which must now be included in our Fauna in place of *D. enicura*. At the time of my first examination I had no specimen of the female of *C. lucifer*, and was misled by the general similarity of coloring of the two species, especially the distribution of colors in the tail, both having it rufous at base, black in the middle, and white at the end; *D. enicura* has the tail-feathers narrower, and the rufous on their bases rather more in extent. The females of *Calothorax pulchra* and of *Myrtis fannia* closely resemble the same sex of the species spoken of above in general plumage, and in having their tails of the same pattern of coloration. In many allied forms among the *Trochilidae*, the females resemble each other so much that a satisfactory determination of them is quite difficult, except with authentic examples of the different species to compare with. — GEO. N. LAWRENCE, *New York City*.

‘ OCCURRENCE OF THE BLACK VULTURE OR CARRION CROW IN OHIO. — On or about December 20, 1876, I came upon three individuals of this species (*Cathartes atratus*, Less.), feeding on the carcass of a hog, in a wooded ravine near Madisonville; one of them I shot at and wounded, but lost sight of it in the woods, and the other two remained in the immediate vicinity long enough to give me an excellent opportunity to observe their peculiarities of form and flight, although I could not approach within gunshot of them. On January 1, 1877, however, I found a specimen that had been killed a few days previous, in the same locality, by Mr. Edwin Leonard, of Madisonville, under circumstances rendering it probable that it was the one I had wounded; its skin is now in my collection.

The occurrence of this bird in Ohio, or in fact anywhere in the Mississippi Valley north of the Ohio River, has heretofore rested solely on Audubon's account of its range, which has been quoted by all subsequent writers; and, being essentially a southern species, its capture here, at a time when the Ohio River was frozen over and the ground covered with several inches of snow, seems worthy of remark. I have identified this species here satisfactorily to myself, on two previous occasions, both in winter, but have never seen the “Turkey Buzzard” (*C. aura*) at that season, although it is quite common during the summer. — FRANK W. LANGDON.

OCCURRENCE OF THE WESTERN NONPAREIL AND BERLANDIER'S WREN AT FORT BROWN, TEXAS. — Dr. J. C. Merrill, U. S. A., in a recent letter to the writer, says: “I have recently (April 23 and 24, 1877) taken two fine males of *Cyanospiza versicolor*, a bird new to our Fauna, although included in Baird, Brewer, and Ridgway's ‘History of North American Birds.’ I have also heard and seen several others. They frequent mesquite chaparral, and betray themselves by their notes, which somewhat resemble those of *C. cyanea*. Berlandier's Wren (*Thryothorus ludovicianus* var. *berlandieri*), also new, but included in Baird, Brewer, and Ridgway's work, I find to be a rather common visitant.” Dr. Merrill also states that

he had just found a beautiful nest of *Spermophila moreleti*, which proves to be a rather abundant species. — J. A. ALLEN, *Cambridge, Mass.*

A CUCKOO'S EGG IN A CEDAR-BIRD'S NEST. — On July 19, 1874, while collecting in a piece of swampy second growth about four miles from Utica, N. Y., I discovered a nest in a small sapling about eight feet from the ground, which, on examination, I found to contain four eggs of the Cedar-Bird (*Ampelis cedrorum*), and one egg of the Cuckoo. From the damp situation, from the appearance of the egg itself, and from the fact of my having found the nest of that bird in the same piece of wood the previous year (1873), I am led to believe it was that of the Black-billed species (*Coccyzus erythrophthalmus*). The nest was deserted and apparently had been for some time, as all the eggs were addled, nor could I see that incubation had begun; certainly it had not in the Cuckoo's egg. I can find no mention in print, nor have I ever before heard, of such a case. — EGBERT BAGG, JR., *Utica, N. Y.*

[The laying of our American Cuckoos in the nests of other birds is doubtless not so rare an occurrence as has been supposed. Two instances have been observed recently in this immediate vicinity, and I have heard of others. Mr. A. M. Frazar, of Watertown, Mass., informs me of one instance where the Yellow-billed Cuckoo (*Coccyzus americanus*) deposited an egg in a Robin's nest, and another case of the laying of the same species in the nest of a Wood Thrush. In the March number of the "Oölogist" (Vol. III, p. 3, published at Utica, N. Y.) an anonymous writer records (since Mr. Bagg's note was received for publication) the finding of two eggs of the "Cuckoo" (species not given) in the nest of a "Redbird" at Gambier, Ohio. — J. A. ALLEN.]

OCCURRENCE FOR THE FIRST TIME IN ENGLAND OF THE ROBIN (*Turdus migratorius*). — Mr. J. E. Harting records the first occurrence in England of this species in an article in "The Field" (published in London, Eng.), of December 23, 1876, and also in the "Zoölogist" for January, of which paper he is editor. The bird was taken alive, owing to its exhausted state, when reaching land at Dover, during April or May.

Mr. Harting is a well-known authority on stragglers, having published in 1872 a Handbook of British Birds, in which was recorded 212 specimens of American birds, belonging to 42 different species. Since that time some of these have been discredited, and others added, until at the present time the list embraces 220 instances of the occasional appearance in Great Britain of North American birds. Of the species referred to in this list, there are five birds of prey, fourteen *Passeres* and *Picariæ*, one *Columba*, fourteen *Grallatores*, and eight *Natatores*. The prevailing winds of the Atlantic being westerly probably accounts for the greater abundance of American stragglers in Europe than the contrary.—H. B. BAILEY, *Orange, N. J.*

# INDEX.

- ACCENTOR auricapillus**, 31.  
**Accipiter atricapillus**, 107.  
**Actitis macularia**, 107.  
**Ægialitis vocifera**, 107.  
     " *wilsonia*, 13, 46.  
**Ægiothus canescens**, 14, 46.  
     " *exilipes*, 51.  
     " *rufescens*, 51.  
**Ægithina leucoptera**, 90.  
**Æstelata hesitata**, 97.  
**Agelaius phoeniceus**, 107.  
**Aix sponsa**, 107.  
**Akhurst, J.**, capture of the Egyptian Goose on Long Island, N. Y., 52.  
**Alcedo alexon**, 107.  
**Allen, J. A.**, note on breeding-dress of Phalaropes, 41; nest and eggs of the Alaskan Wren, 82; on Cuckoos laying in nests of other birds, 111.  
**Ammodromus caudatus**, 27, 28.  
**Anas obscura**, 96.  
**Anatide**, neotropical, notice of Selater and Salvin's paper on, 24.  
**Anser gambeli**, 14, 46.  
     " *segetum*, 96.  
**Anthus coronata**, 32.  
     " *l'herminieri*, 33.  
     " *ludovicianus*, 17, 107.  
**Antrostomus macromystax**, 93.  
     " *nigrescens*, 93.  
**Aramus scolopaceus**, 23.  
**Archibuteo ferrugineus**, 26.  
**Audubonia occidentalis**, 95.  
**BAGG, E., Jr.**, Cuckoo's egg in a Cedar-Bird's nest, 110.  
**Bailey, H. B.**, occurrence of Leconte's Bunting in Iowa, 26; occurrence of *Passerculus princeps* in New York, 78; occurrence for the first time of the Robin in England, 110.  
**Bartramia longicauda**, 107.  
**Bernicla hutchinsi**, 14, 46.  
     " *leucopsis*, 17.  
**Botaurus lentiginosus**, 107.  
**Brachyotus cassini**, 14.  
**Brewer, T. M.**, a defence of his Catalogue of New England Birds, 44; a new bird to Massachusetts, 78; a new form of *Sturnia* to New England, 78; breeding of Leach's Petrel on the coast of Maine, 80.  
**Brewster, Wm.**, the Black-and-Yellow Warbler, 1; northern range of the Sharp-tailed Finch, 28; two undescribed nests of Californian birds, 37; an undescribed Hybrid between two North American Grouse, 66; the Yellow-throated Warbler, 102.  
**Brown, J. A. H.**, on *Ægiothus exilipes* in Europe, 51.  
**Brown, N. C.**, notes on birds new to the fauna of Maine, etc., 27.  
**Bubo virginianus**, 80.  
**Bunting**, Chestnut-collared, 78.  
     " *Leconte's*, 26.  
     " *MacCown's*, 52.  
**Burroughs, J.**, notice of his "Wake-Robin," 48.  
**Buteo cooperi**, 89.  
**Buzzard, Ferruginous**, 26.  
**CALAMOSPIZA bicolor**, 92.  
**Calocitta collicie**, 93.  
**Cardinalis virginianus**, 108.  
**Carpodacus amplus**, 60 - 66.  
     " *frontalis*, 60 - 66.  
     " *purpureus*, var. *californicus*, 37.  
**Catharista atrata**, 94.  
**Cathartes atratus**, 109.  
**Centurus carolinus**, 108.  
**Ceryle americana**, 94.  
**Charadrius fulvus**, 107.  
     " *virginicus*, 107.  
**Chat, Yellow-breasted**, 16.  
**Chenalopex aegyptius**, 52.  
**Chrocephalus atricilla**, 96, 107.  
     " *philadelphia*, 107.  
**Chrysomitris yarelli**, 92.  
**Chrysopoga typica**, 92.  
**Cistothorus stellaris**, 15.  
**Clangula albeola**, 107.  
**Coale, H. K.**, MacCown's Longspur in Illinois, 52; *Juncos oregonus* in Illinois, 82; notes on *Nyctale acadica*, 83.  
**Coccothraustes ferreirostris**, 92.  
**Coccyzus americanus**, 107, 108.  
     " *erythrophthalmus*, 107, 108.  
**Collocalia ludovicianus**, 90.  
**Collurio ludovicianus** var. *excubitoroides*, 21.  
**Contopus borealis**, 16.  
**Conurus carolinensis**, 50.

- Cooper, J. G., notice of his "New Facts relating to Californian Ornithology," 76; on Seventy-five Doubtful West-Coast Birds, 88.
- Coot, 17.
- Corvus americanus*, 14, 45.  
" *ossifragus*, 13, 46, 93.
- Cory, C. B., the Black Gyr-Falcon in Massachusetts, 27.
- Coturniculus lecontei*, 26.  
" *passerinus*, 16, 47.
- Cooper, E., note on *Podiceps dominicus*, 26; Eastward Range of the Ferruginous Buzzard, 26; corrections of nomenclature in the genus *Siturus*, 29; Western Range of *Conurus carolinensis*, 50; note on the Cinnamon Teal, 51; a pigeon (*Leptoptila albifrons*) new to the United States Fauna, 82; *Melopelia leucoptera* in Colorado, 83.
- Cowbird, Common, 85.  
" Dwarf, 85.
- Crow, Common, 14, 45.  
" Fish, 13, 46.
- Cuckoo, Black-billed, 110.  
" Yellow-billed, 110.
- Cupidonia cupidini-columbianus*, 67.  
" *cupido*, 66.  
" var. *pallidicinctus*, 52.
- Curlew, Long-billed, 17.
- Cyanocitta beecheyi*, 93.  
" *ultramarina*, 93.
- Cyanocorax geoffroyi*, 93.
- Cyanospiza versicolor*, 109.
- DAFILA *nrophasiana*, 96.
- D'Hamonville, Baron, notice of his "Catalogue des Oiseaux d'Europe," 106.
- Deane, R., unusual abundance of the Snowy Owl in New England, 9; note on the Barnacle Goose, 18; occurrence of the Sooty Tern in Massachusetts, 27; probable breeding of the Acadian Owl in Massachusetts, 84.
- Dendroica auduboni*, 27.  
" *blackburniæ*, 48, 104.  
" *cærula*, 14, 21, 48.  
" *coronata*, 31.  
" *dominica*, 102, 108.  
" *maculosa*, 1.  
" *montana*, 90.  
" *striata*, 90.  
" *virens*, 105, 107.
- Demigretta ludoviciana*, 51.
- Diomedea chlororhyncha*, 96.  
" *fuliginosa*, 97.
- Doricha euicura*, 108.
- Dove, Carolina, 15.  
" White-crowned, 82.  
" White-winged, 83.
- Dryotomus delatris*, 94.
- Dury, C., Fecundity of the Carolina Wren, 50.
- ECTOPISTES migratorius*, 107.
- Elliot, D. G., remarks on *Scelaphorus alleni*, 97.
- Emberiza atricapilla*, 92.
- Empidonax acadica*, 108.
- Enicocichla*, see *Henicocichla*.
- Eupsychortyx cristatus*, 95.
- FALCO columbarius*, 79.  
" *sacer* var. *labrador*, 27.
- Falconidae, notices of Ridgway's papers on, 70-73.
- Finch, Californian Purple, 37.  
" Sharp-tailed, 27, 28.
- Flycatcher, Great-crested, 16.  
" Olive-sided, 16.  
" Townsend's, 77.
- Fox, W. H., capture of the Philadelphia Vireo in New Hampshire, 78.
- Frazar, A. M., Audubon's Warbler in Massachusetts, 27; persistency at nest-building in a House-Wren, 78; the Mottled Owl as a Fisherman, 80.
- Fringilla maculata*, 92.
- Fulica americana*, 17.
- Fulix affinis*, 107.  
" *collaris*, 107.
- GALLINULE, Florida, 83.  
" Purple, 83.
- Gambetta flavipes*, 95, 107.
- Gannet, Booby, 14, 46.
- Garrod, A. H., notice of papers by, 23.
- Geothlypis philadelphia*, 16, 17.  
" *trichas*, 89.  
" *velatus*, 89.
- Gnateatcher*, Blue-Gray, 14, 20, 49.
- Goose, Barnacle, 18.  
" Egyptian, 52.  
" Gambel's, 14, 46.  
" Hutchins's, 14, 46.
- Grackle, Boat-tailed, 12, 45.  
" Long-tailed, 85.
- Graculus carbo*, 96.
- Grant's Catalogue of the Birds of Malta and Gozo, notice of, 70.
- Grebe, Saint Domingo, 26.
- Grouse, Hybrid, 66.  
" Pinnated, 66.  
" Sharp-tailed, 66.
- Grns americana*, 95.
- Guadaloupe Island, birds of, discussed with reference to the present genesis of species, 58.
- Gulls, Skua or Jager, 23.
- Gyr-Falcon, Black, 25, 27.
- Gyr-Falcons, 25.
- HÆMATOPUS *ater*, 95.
- Haliaeetus leucocephalus*, 107.  
" *pelagicus*, 94.
- Harporhynchus rufus*, 107.  
" var. *longirostris*, 90.

- Hawk, Pigeon, 79.  
 " Sharp-shinned, 14.
- Helminthophaga celata*, 21.  
 " *chrysoptera*, 16, 47.  
 " *lawrencei*, 19.  
 " *leucobronchialis*, 79.  
 " *pinus*, 16.
- Helmitherus vernivornis*, 16, 21.
- Henicocichla auropillus*, 31.  
 " *ludovicianus*, 33.  
 " *major*, 33.  
 " *motacilla*, 33.  
 " *novaboracensis*, 32.  
 " *sulphurascens*, 33.
- Henshaw, H. W., description of a new species of Humming-Bird from California, 53.
- Heron, Louisiana, 51.
- Herrick, H., capture of a second specimen of *Helminthophaga lawrencei*, 19.
- Hierofalco candicans*, 25.  
 " *islandicus*, 25.  
 " *labrador*, 25.
- Hirundo bicolor*, 107.
- Humming-Bird, Green-backed, 53.  
 " Rufous-backed, 53.
- Hydrochelidon lariformis*, 34, 96.  
 " *nigrum*, 96.
- Hypocanthus stanleyi*, 92.
- Hypocolius ampelinus*, 91.
- ICTERIA virens, 16.
- Icterus baltimore*, 92.  
 " *californicus*, 92.  
 " *pustulatus*, 92.  
 " *spurius* var. *affinis*, 85.
- JUNCO *annectens* 60-66.  
 " *hyemalis*, 17.  
 " *insularis*, 60-66.  
 " *oregonus*, 82.
- KINGLET, Golden-crested, 17.
- LAGOPUS *albus*, 14, 46.
- Lamb, W. F., on nest and eggs of Townsend's Flycatcher, 77.
- Langdon, F. W., Black Vulture in Ohio, 109.
- Lanius elegans*, 91.  
 " *lahtora*, 91.
- Larus belcheri*, 96.
- Lawrence, G. N., occurrence of the Barnacle Goose (*Bernicla leucopsis*) on Long Island, N. Y., 18; note on *Cupidonia cupido* var. *pallidicinctus*, 52; note on *Doricha enicura*, 108.
- Leptoptila albifrons*, 82.
- Lobipes hyperboreus*, 38, 42.
- Longspur, MacCown's, 52.
- Lophophanes bicolor*, 108.
- Lophortyx elegans*, 95.
- Loxia leucoptera*, 103.
- MACRORHAMPHUS *griseus*, 107.
- Mareca americana*, 107.
- McCanley, C. A. II, notice of his Notes on Texan Ornithology, 76.
- Melanerpes erythrocephalus*, 94.
- Melospelia leucoptera*, 83, 95.
- Mergus cucullatus*, 107.
- Merriam, C. H., his "Review of Birds of Connecticut" noticed, 107.
- Merrill, J. C., a Humming-Bird new to the Fauna of the United States, 26; notes on *Molothrus aeneus*, 85; Western Nonparcil and Berlandier's Wren in Texas, 109.
- Micropalama himantopus*, 17, 48.
- Milvulus forficatus*, 21.
- Minot, H. D., notice of his "Land-Birds and Game-Birds of New England," 49.
- Molothrus aeneus*, 85.  
 " *pecoris*, 85.  
 " " var. *obscurus*, 85.
- Motacilla auropilla*, 31.  
 " *canadensis*, 31.  
 " *fluvialis*, 32.  
 " *leucoptera*, 90.  
 " *naevia*, 32.  
 " *novaboracensis*, 32.  
 " *tigrina*, 32.
- Murdock, J., the Pigeon-Hawk at sea, 79.
- Myiadestes townsendi*, 77.
- Myiarchus crinitus*, 16.
- Myiodioctes minutus*, 14, 46.  
 " *mitratus*, 16, 21.
- Nauclerus furcatus*, 107.
- Nelson, E. W., a contribution to the biography of Wilson's Phalarope, 38; the Louisiana Heron in Indiana, 51; notice of his "Birds of Northeastern Illinois," 68.
- Nettion crecca*, 13, 46.
- Nisus fusens*, 14.  
 " *gmdlachti*, 71.
- Nonparcil, Western, 109.
- Numenius borealis*, 107.  
 " *hudsonicus*, 107.  
 " *rufiventris*, 95.
- Nyctale acadica*, 14, 15, 83, 84.
- Nyctea scandiaca*, 9.
- ŒDEMA perspicillata*, 107.
- Onychotes gruberi*, 89.
- Oporornis formosus*, 108.
- Oriole, Baltimore, 92.  
 " Bullock's, 86.  
 " Hooded, 86, 87.  
 " Orchard, 86, 87.
- Ortyx fasciatus*, 95.  
 " *virginianus*, 15.
- Otus wilsonianus*, 14.
- Owl, Acadian, 14, 15, 83, 84.  
 " Barn, 28.  
 " Great-horned, 80.  
 " Hawk, 78.  
 " Long-eared, 14.

- Owl, Mottled, 80.  
 " Short-eared, 14.  
 " Snowy, 9.  
 PARROT, Carolina, 50.  
 Parus carolinensis, 90.  
 Passerculus princeps, 27, 78.  
 Pedicocetes phasianellus var. columbianus, 66.  
 Pelecanus trachyrhynchus, 22.  
 Pelican, White, 22.  
 Perissoglossa tigrina, 16.  
 Petrel, Leach's, 28, 80.  
 Penecea ruficeps, 37.  
 Phalarope, Northern, 38, 42.  
 " Red, 38, 42.  
 " Wilson's, 38.  
 Phalaropus fulicarius, 42.  
 " lobatus, 38.  
 Philomachus pugnax, 83.  
 Picoides americanus, 16.  
 " arcticus, 16.  
 Picus lineatus, 93.  
 " pubescens, 107.  
 " villosus, 14, 45, 107.  
 Pipilo consobrinus, 60, 66.  
 " maculatus, 60-66.  
 Platalea ajaja, 95.  
 Plectrophanes maccowni, 52, 92.  
 " ornatus, 78.  
 Plotus anhinga, 23.  
 Plover, Wilson's, 13, 46.  
 Podiceps dominicensis, 26, 97.  
 " minor, 97.  
 Polioptila cerulea, 14, 20, 47.  
 Polyborus cheriway, 61-66.  
 " lutosus, 60-66.  
 " tharus, 60-66.  
 Porphyrio martinica, 83.  
 Porzana jamaicensis, 22.  
 " noveboracensis, 17.  
 Procellariidae, notice of Salvin's paper on, 69.  
 Progne purpurea, 107.  
 Puffinus obscurus, 97, 107.  
 " fuliginosus, 107.  
 Purdie, H. A., distribution of New England Birds, a reply to Dr. T. M. Brewer, 11; notice of a few birds of rare or accidental occurrence in New England, 20.  
 Pyrrhula erythrogastra, 21, 107.  
 Pyrgisoma bicarvatus, 91.  
 Pyrrhophena rickei, 26.  
 Pyrrhula inornata, 91.  
 QUAIL, Common, 15.  
 Querquedula cyanoptera, 51, 75.  
 " discors, 107.  
 Quiscalus major, 12, 45, 93.  
 " purpureus, 93.  
 RAIL, Black, 22.  
 " King, 22.  
 " Yellow, 17, 22.  
 Rallus elegans, 22.  
 " longirostris, 22.  
 Ramphopsis flamigerus, 92.  
 Recent ornithological articles in American journals, 73.  
 Redbird, Summer, 21.  
 Regulus calendula, 59, 61, 107.  
 " obscurus, 59, 61.  
 " satrapa, 17.  
 Ridgway, R., on Geographical Variation in *Turdus migratorius*, 8; the Birds of Guadalupe Island discussed with reference to the present genesis of species, 58; notice of his "Studies of the American Falconidae," 70, 73.  
 Roberts, T. S., notes on the breeding of the Black Tern, 34.  
 Robin, 8, 12, 14, 110.  
 Ruff, 83.  
 Rhyacophilus solitarius, 107.  
 SALPICETES guadalupensis, 60-66.  
 " obsoletus, 60-66.  
 Saltator rufiventris, 92.  
 Salvin, O., notice of his paper on *Procellariidae*, 69.  
 Sandpiper, Baird's, 28.  
 " Stilt, 17, 48.  
 Sarcorhamphus gryphus, 94.  
 Saunders, H., notice of his papers on *Stercorariinae* and *Sterninae*, 23, 24.  
 Sayornis fuscus, 93.  
 Selater, P. L., and Salvin, O., notice of their paper on Neotropical *Anatidae*, 24.  
 Scops asio, 80, 107.  
 Seiurus, see Siurus.  
 Selasphorus alleni, 53, 97, 101.  
 " henschawi, 102.  
 " rufus, 53, 97, 101.  
 Setophaga ruticilla, 90.  
 Shrike, White-rumped, 21.  
 Sialia sialis, 89.  
 Simorhynchus psittacula, 107.  
 Sitta pusilla, 99.  
 Siurus, corrections of nomenclature in the genus, 29.  
 Siurus auricapillus, 31, 90.  
 " ludovicianus, 16, 33.  
 " motacilla, 33.  
 " naevius, 32.  
 " sulphurascens, 33.  
 " tenuirostris, 33.  
 Snowbird, Eastern, 17.  
 " Oregon, 82.  
 Sparrow, Field, 15.  
 " Ipswich, 27, 78.  
 " Rufous-crowned, 37.  
 " Yellow-winged, 16, 47.  
 Spermophila moreleti, 103.  
 Steganopus wilsoni, 38.  
 Stelgidopteryx serripennis, 21.  
 Sterna antillarum, 96.

- Sterna fuliginosa*, 22, 27, 44, 75.  
 " *plumbea*, 34.  
*Strix flammea* var. *pratineola*, 28.  
*Sula bassana*, 96.  
 " *fiber*, 14, 46, 96.  
 Swallow, Rough-winged, 21.  
*Sylvia anthoides*, 33.  
 " *aurocapilla*, 31.  
 " *leucoptera*, 90.  
 " *novaboracensis*, 32.  
 " *tigrina*, 32.  
*Syrnium nebulosum*, 94.  
 TEAL, Cinnamon, 51.  
 " European, 13, 46.  
 Tern, Black, 34.  
 " Sooty, 22, 27.  
 Terns, notice of Saunders's paper on, 24.  
*Thalassidroma leachi*, 28.  
 " *leucorrhoa*, 80, 107.  
 Titlark, 17.  
*Tinnunculus sparverius*, 112.  
*Trichas delatfieldi*, 51.  
*Tringa bairdi*, 28.  
 " *bonapartei*, 107.  
 " *lobata*, 38.  
 " *maculata*, 107.  
 " *pusilla*, 107.  
*Tringites rufescens*, 107.  
*Troglodytes aëdon*, 78.  
 " *parvulus* var. *alascensis*, 82.  
 Trotter, S., capture of a second specimen of *Helminthophaga leucobronchialis*, 79.  
*Trupialis militaris*, 92.  
*Thryomanes bewicki*, 60 - 66.  
 " *brevicauda*, 60 - 66.  
*Thryothorus ludovicianus*, 40.  
 " " var. *berlandieri*, 109.  
 Thrush, Golden-crowned, 30, 32.  
 " Long-billed Water, 16, 30, 33.  
 " Short-billed Water, 30, 32.  
*Turdus aquaticus*, 32.  
 " *aurocapillus*, 32.  
 " *citreus*, 31.  
 " *coronatus*, 32.  
 " *flavirostris*, 89.  
 " *fuscescens*, 107.  
 " *migratorius*, 8, 9, 12, 14, 110.  
*Turdus minimus*, 31.  
 " *motacilla*, 33.  
 " *novaboracensis*, 32.  
 " *pallasi*, 107.  
 " *propinquus*, 9.  
 " *rufopaliatus*, 90.  
 " *swainsoni*, 107.  
 URIA *columba*, 107.  
 VENNOR, H. G., notice of his Canadian "Birds of Prey," 24.  
*Vireo agilis*, 90.  
 " *gilvus*, 15.  
 " " var. *swainsoni*, 90.  
 " *novaboracensis*, 15, 47.  
 " *philadelphia*, 78.  
 " *virescens*, 91.  
*Vireo*, Philadelphia, 78.  
 " Warbling, 15.  
 " White-eyed, 15.  
 WARBLER, Audubon's, 27.  
 " Black-and-Yellow, 1  
 " Blackburnian, 48, 104.  
 " Black-throated Green, 104.  
 " Blue, 14, 21, 48.  
 " Blue-winged, 16.  
 " Cape May, 16.  
 " Golden-winged, 16.  
 " Hooded, 16, 21.  
 " Mourning, 16.  
 " Orange crowned, 21.  
 " Pine, 99.  
 " Worm-eating, 16, 21.  
 " Yellow-throated, 102.  
 Wheaton, J. M., on the Ruf and Purple Gallinule in Ohio, 83.  
 Woodpecker, Black-backed Three toed, 16.  
 Woodpecker, Banded Three-toed, 16.  
 " Hairy, 14, 15, 45.  
 Wren, Alaskan, 82.  
 " Berlandier's, 109.  
 " Carolina, 50.  
 " House, 78.  
 " Short-billed Marsh, 15.  
 XANTHORUS *mexicanus*, 92.  
*Xema furcatum*, 96.  
 " *sabinei*, 107.  
 ZENÆDURA *carolinensis*, 15.  
*Zonotrichia coronata*, 92.







